# Length of Stay by Operation, Western Region, 1982

A Companion Volume To
Length of Stay
by Diagnosis,
Western Region, 1982

Commission on Professional and Hospital Activities
Ann Arbor, Michigan
October, 1983

# LENGTH OF STAY

Copyright 1983 by Commission on Professional and Hospital Activities (CPHA), Ann Arbor, Michigan. All rights reserved. Printed in the United States of America. This book, or parts thereof, may not be reproduced in any form without permission of the publisher.

*Library of Congress Catalog Card Number: 73-173193*

This volume is one of the books in a series entitled
Length of Stay . . .

. . . by Diagnosis, United States, 1982
. . . by Diagnosis, United States, North Central Region, 1982
. . . by Diagnosis, United States, Northeastern Region, 1982
. . . by Diagnosis, United States, Southern Region, 1982
. . . by Diagnosis, United States, Western Region, 1982
. . . by Diagnosis, Canada, 1982

. . . by Operation, United States, 1982
. . . by Operation, United States, North Central Region, 1982
. . . by Operation, United States, Northeastern Region, 1982
. . . by Operation, United States, Southern Region, 1982
. . . by Operation, United States, Western Region, 1982
. . . by Operation, Canada, 1982

Geriatric Length of Stay, by Diagnosis and Operation, United States, 1982
Pediatric Length of Stay, by Diagnosis and Operation, United States, 1982

The length of stay series began in 1963, and certain volumes are still available from the Commission on Professional and Hospital Activities.

# PREFACE

This book is one volume of a set of 14 books presenting statistics on hospital stay during 1982 in North America. The complete set is the latest in a series of publications on length of stay in hospitals that participate in the Professional Activity Study (PAS) of the Commission on Professional and Hospital Activities. Several organizations have contributed data to CPHA from other abstracting services. Some of those data were included in the 1982 series.

These statistics, which are also available on computer-readable magnetic tape, can provide a valuable perspective for physicians, administrators, utilization review authorities, and planners who make decisions based on length of hospitalization. They can also serve as an aid in establishing check points for "continued stay review" and in the implementation or revision of utilization review systems.

We extend our appreciation to the hospitals and state organizations that supported this study by making their data available.

## PROJECT STAFF
## 1982 LENGTH OF STAY BOOKS

Edward J. Kobrinski, Manager, Research Services
Carol A. Prevost, Research Assistant

# CONTENTS


**Preface** . . . . . iii
**Discussion of the Data** . . . . . 1
Source of the Data. . . . . 1
Data Quality. . . . . 2
Coverage . . . . . 3
Organization of Length of Stay Tables. . . . . 5
Definition of Multiple Diagnosis Patients . . . . . 5
Definition of Operated Patients . . . . . 6
Explanation of Statistics. . . . . 6
Defining the Extended Stay Case . . . . . 7
**Length of Stay Tables** . . . . . 15
Summary of All Patients. . . . . 16
Operation Groups 501-764 . . . . . 17
**Appendix: *ICD-9-CM* Codes for Selected Operation Groups** . . . . . 83
**Index to Length of Stay Tables** . . . . . 87

# DISCUSSION OF THE DATA

## Source of the Data

Data in the length of stay books were compiled from individual patient discharge records submitted by hospitals participating in the Professional Activity Study (PAS) of the Commission on Professional and Hospital Activities (CPHA). In addition, some books contain discharge data from hospitals participating in information systems other than PAS.

CPHA is an organization dedicated to the improvement of hospital and medical care through research, education, publishing, and systems development. Established in 1955 under grants from the W.K. Kellogg Foundation, CPHA is nonprofit, nongovernmental, noncommercial. CPHA is sponsored by the American College of Physicians, the American College of Surgeons, the American Hospital Association, and the Southwestern Michigan Hospital Council.

The following steps explain the PAS method of storing and retrieving medical data:

1. In each participating hospital, personnel transfer data from the medical record of each discharged patient into the PAS System via PAS+ microcomputer (floppy disk), PAS Case Abstract, or magnetic tape.

2. The disk, abstract, or tape is mailed to the center of PAS operations in Ann Arbor, Michigan, where it is entered into the computer system.

3. Data then undergo a comprehensive editing process and errors are corrected by CPHA and contributing hospitals.

4. The computer system continually adds data to the permanent history file. Participating hospitals are provided access to these data through interactive systems, reports, and in-house computers. The stored data are also available to researchers on a contractual basis.

5. Length of stay statistics from this file are published annually in this series of books.

## Data Quality

Data quality has always been and remains one of CPHA's major concerns. Although the hospital has basic responsibility for the quality of PAS data, CPHA shares that responsibility in its function as an extension of the hospital's medical information network. CPHA's position of leadership in health information processing systems and data usage necessitates a constant striving for data accuracy.

CPHA goes to considerable length to prevent and correct errors in data submission by training hospital medical record personnel in coding and abstracting methods, by monitoring submitted data, and by meticulously checking and editing the data for errors. Over 350 checks are performed during the processing of each record to ensure the reasonableness of the data received. Examples of discrepancies in the data that are detected by the audit are: hospital day of operative episode exceeding total stay, cesarean section with a normal delivery diagnosis, a sex-specific diagnosis or operation with the wrong sex recorded, incalculable age or stay, unrecorded or invalid diagnosis. When illogical, invalid, or missing data are detected, about 40 percent of the errors are corrected by the computer. The remainder are referred to the hospital staff through error messages generated by the computer.

In addition to the initial data editing, the computer system that produces the length of stay tables does numerous edits to eliminate uncorrected or unusual data. The following events noted on a record cause that record to be excluded:

1. death
2. stillborn
3. admitted from other short-term hospital
4. discharged to other short-term hospital
5. discharged against medical advice
6. length of stay greater than 99 days (included only in the percentile calculations)

These patient records were excluded because they do not represent typical short-term patients. Data that pass all the edits represent patients whose length of stay was solely determined and managed by the physician.

## Coverage

Data chosen for inclusion in the length of stay series are from general, non-federal, short-term hospitals in the United States. Short-term hospitals are defined as those in which the median stay is shorter than 30 days. The map indicates the boundaries of the regional books.

Most of the data used in the length of stay book series are from short-term, general hospitals participating in PAS through CPHA. In addition, the following organizations have sent data to CPHA: Colorado Hospital Association, Kansas Hospital Association, Kaiser Foundation Hospitals, Michigan Health Data Corporation, Missouri Health Data Corporation, The National Association of Children's Hospitals and Related Institutions, Inc. (NACHRI), Sisters of Mercy Health Corporation (SMHC), Utilization Information Service of the Hospital Association of New York State, and the Vermont Hospital Association. Several individual hospitals throughout the United States have also sent data to CPHA for special studies. These contributed data expand CPHA as a national health information center. The cooperation of these organizations is appreciated.

Hospitals participating in PAS from United States Territories and Puerto Rico were excluded from the series. Data from psychiatric hospitals were excluded because the psychiatric hospitals are not general hospitals. Data from psychiatric units within general hospitals were included. Pediatric hospitals were also included.

Characteristics of all the hospitals with data included in the length of stay books are displayed on the following page.

*Census Divisions and Regions in the United States*

**Bed Size, Region, and Census Division: Included Hospitals Compared with All Short-Term Nonfederal Hospitals[1] in the U.S. January-December 1982**

| Hospital Class (1) | Short-Term Nonfederal Hospitals U.S. Total[2] (2) | Number in Books (3) | Percent in Books (4) | Beds U.S. Total[2] (5) | Number in Books (6) | Percent in Books (7) |
|---|---|---|---|---|---|---|
| **Bed Size** | | | | | | |
| 6-24 beds | 254 | 35 | 13.8 | 4,814 | 696 | 14.5 |
| 25-49 | 971 | 180 | 18.5 | 35,155 | 6,841 | 19.5 |
| 50-99 | 1,432 | 318 | 22.2 | 103,270 | 23,365 | 22.6 |
| 100-199 | 1,374 | 357 | 26.0 | 193,551 | 50,983 | 26.3 |
| 200-299 | 712 | 227 | 31.9 | 172,913 | 55,670 | 32.2 |
| 300-399 | 423 | 136 | 32.2 | 144,133 | 46,500 | 32.3 |
| 400-499 | 267 | 94 | 35.2 | 117,869 | 41,567 | 35.3 |
| 500 or more | 328 | 84 | 25.6 | 226,214 | 55,946 | 24.7 |
| Total | 5,761 | 1,431 | 24.8 | 997,919 | 281,568 | 28.2 |
| **Region and Census Division** | | | | | | |
| Northeast | 828 | 283 | 34.2 | 213,258 | 67,701 | 31.7 |
| New England | 244 | 61 | 25.0 | 49,916 | 13,631 | 27.3 |
| Middle Atlantic | 584 | 222 | 38.0 | 163,342 | 54,070 | 33.1 |
| North Central | 1,694 | 571 | 33.7 | 295,316 | 110,053 | 37.3 |
| East North Central | 897 | 350 | 39.0 | 195,881 | 77,902 | 39.8 |
| West North Central | 797 | 221 | 27.7 | 99,435 | 32,151 | 32.3 |
| South | 2,162 | 306 | 14.2 | 340,530 | 61,715 | 18.1 |
| South Atlantic | 815 | 211 | 25.9 | 160,263 | 42,298 | 26.4 |
| East South Central | 500 | 30 | 6.0 | 73,359 | 7,375 | 10.1 |
| West South Central | 847 | 65 | 7.7 | 106,908 | 12,042 | 11.3 |
| West | 1,077 | 271 | 25.2 | 148,815 | 42,099 | 28.3 |
| Mountain | 369 | 92 | 24.9 | 41,978 | 14,587 | 34.7 |
| Pacific | 708 | 179 | 25.3 | 106,837 | 27,512 | 25.8 |
| Total | 5,761 | 1,431 | 24.8 | 997,919 | 281,568 | 28.2 |

The Pediatric book was compiled from the data of 1,428 hospitals. The Geriatric book was compiled from the data of 1,427 hospitals.

[1] Hospitals with average stay less than 30 days, excluding psychiatric hospitals,

[2] 1981 Annual Survey of Hospitals, magnetic tape, American Hospital Association.

## Organization of Length of Stay Tables

The length of stay tables are organized according to CPHA's cross classification, a patient classification system similar to Diagnosis Related Groups (DRGs). The operated classification system is called *ICD-9-CM List B: Hospital Procedure Groups.*[1] Within each procedure group, patients are further stratified by single or multiple diagnoses and age.

Patient data were assigned *ICD-9-CM List B* groups on the basis of the principal procedure. "Principal procedure" is an item in the Uniform Hospital Discharge Data Set (UHDDS)[2] and generally is that therapeutic procedure most related to the principal diagnosis. Procedure information for all patient data included in this book was coded using *ICD-9-CM.*

In *List B* there are 250 groups that contain at least one UHDDS Class 1 procedure. Fourteen of the 264 *List B* groups consist solely of Class 2, 3, or 4 procedures. This book displays data only for those 250 groups with Class 1 procedures.

### Definition of Multiple Diagnosis Patients

Patients are classified in the multiple diagnosis category if they had at least one valid secondary diagnosis in addition to the principal. The following codes are not considered valid secondary diagnoses for purposes of this classification:

1) Manifestation codes (conditions that evolved from underlying diseases [etiology] and are in italics in *ICD-9-CM Volume 1*[3])
2) Codes V27.0-V27.9 (outcome of delivery)
3) E codes (external causes of injury and poisoning)

[1] *ICD-9-CM List B: Hospital Procedure Groups* is based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM),* Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

[2] Additional information regarding UHDDS principal procedure and classes of procedures can be obtained in *UHDDS Classes of Procedures, ICD-9-CM,* Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

[3] *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM), Volume 1, Diseases: Tabular List,* Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

Definition of Operated Patients

Operated patients are patients with an *ICD-9-CM* procedure code that fell into a UHDDS category containing at least one Class 1 procedure. Class 1 procedures are defined as those procedures that carry an operative or anesthetic risk, require highly trained personnel, or require special facilities or equipment. The following *ICD-9-CM List B* codes contain a mixture of Class 1 and 2-4 procedures: 548, 557, 575, 576, 587, 609, 615, 619, 626, 630, 636, 646, 654, 666, 679, 682, 686, 690, 696, 706, 758, 760, 764. All the other *ICD-9-CM List B* groups contain only Class 1 procedures or they contain only Class 2-4 procedures and are excluded from this book.

## Explanation of Statistics

The *List B* diagnosis group titles heading the tables are followed in parentheses by the *ICD-9-CM* diagnosis codes included in the *List B* group.

Length of stay was calculated by the computer from admission and discharge dates by counting the day of admission but not the day of discharge. Patients discharged on the day of admission were counted as staying one day in the calculation of average stay and variance but as staying less than one day in calculation of the percentiles. Thus, if more than 5 percent of the patients in a cell (one line of data within a table) were discharged on the day of admission, the 5th percentile is shown as "<1."

Patients who were hospitalized 100 days or longer are not included in Number of Patients, Average Stay, or Variance. The percentiles shown in columns 5 through 11, however, do include these patients.

The variance displayed is an unbiased estimate of the mean squared deviation from the average. For each cell it is calculated as $\sum_{i=1}^{N} (X_i - \bar{X})^2/(N-1)$, where the $X_i$'s are the length of stay for each patient, $\bar{X}$ is the average stay in the cell, and N is the number of patients in the cell.

The 5th, 10th, 50th, 75th, 90th, 95th, and 99th percentiles of stay are displayed in days. If, for example, the 10th percentile for a group of patients is four, it means that 10 percent of the patients stayed four days or less. The 50th percentile is the median. Any percentile with a value of 100 days or more is listed as 99+.

Determining a stay percentile for a group of patients is done by arranging the individual patient stays from low to high. Counting up from the lowest stay to the point where half of the patients have been counted yields the value of the 50th percentile. Counting off one-tenth of the total patients gives the 10th percentile, and so on.

## Defining the Extended Stay Case

Concurrent review of a continuous hospital stay that equals or exceeds the period defined in the hospital's utilization review plan or memorandum of understanding with the utilization review authority is called extended stay review. This review, whether applied universally to all patients within the facility or focused on known or suspected problem areas, is designed to answer the following question: Is further hospitalization medically necessary?

Ideally, this question is posed at that point in the patient's hospitalization which is neither so early as to be annoyingly pointless nor so late that opportunity for cost containment has passed. The median (50th percentile) stay for similar patients, as reflected in current length of stay tables, represents an effort to arrive at a sensible compromise as to the correct day on which to initiate the review process.

To satisfy the requirements for extended stay review, a method of defining an extended stay must begin with a definition of factors that affect stay. Presented here is a technique for determining extended stay review dates according to the most important scheduled operation, whether the patient had single or multiple diagnoses, and the patient's age.

While the usual procedure for assigning the initial review date is based upon the admitting diagnosis (problem), there are circumstances when a table based upon the contemplated operation is much more meaningful. The latter situations involve diagnoses where two or more quite different operations (and different lengths of stay) could be appropriate managements. Examples would be benign prostatic hypertrophy (cystoscopy, transurethral resection [TUR] or other prostatectomy); primary malignancy of lung, suspected or previously diagnosed (bronchoscopy or resection of lung); and angina pectoris (coronary arteriography or bypass surgery). Many diagnoses, on the other hand, are usually managed by a single type of operation and the operated section of the diagnosis table and the operation table appear to be almost identical. Examples would be uncomplicated inguinal hernia (herniorrhaphy), acute appendicitis (appendectomy), and chronic cholecystitis with cholelithiasis (cholecystectomy).

An extended stay review date can be projected for each patient at the time of admission to the hospital by using one of the length of stay books published by CPHA. With the appropriate book, the six steps to determine a review date are as follows:

Step 1. Find the length of stay operation group (derived from the *ICD-9-CM List B: Hospital Procedure Groups*[1]) that contains the contemplated operation in one of the following ways:

a. Usually the diagnosis can be found in *Concurrent Review Screening,*[2] developed for admission certification and extended stay review purposes, in which diagnoses are listed alphabetically. The length of stay group for each diagnosis is given in column 4 where appropriate reference is made to the operation tables.

b. Procedures can also be found in the Alphabetic Index at the back of the length of stay books.

c. If an operation is not listed in the index to the length of stay tables, the code for the diagnosis can be found in Volume 3 of *ICD-9-CM.*[3] That code can then be located in *List B* or in the heading of the length of stay table.

Step 2. Find the appropriate table in the length of stay book. The group number precedes the title of each table.

Step 3. Find the appropriate part of the table for single or multiple diagnoses.

Step 4. Determine the patient's age on day of admission and locate the correct age group to find the appropriate cell.

Step 5. Find the 50th percentile of stay for similar patients at the point where the 50th percentile column (column 7) of the table intersects the horizontal line that includes the pertinent variables for the patient having an extended stay review date assigned.

Example: On 3 November a 57-year-old patient is admitted to a hospital for elective surgery, a transurethral resection. The admitting diagnosis is prostatic hyperplasia, and no secondary diagnoses are recorded. The following illustrations show the five steps for finding the date on which the patient becomes an extended stay case and must be reviewed.

[1] *ICD-9-CM List B: Hospital Procedure Groups,* Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

[2] *Concurrent Review Screening – Criteria for Hospital Admission and Assignment of Length of Stay,* Second Edition, Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

[3] *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM), Volume 3, Procedures: Tabular List and Alphabetic Index,* Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978.

Step 1. The length of stay group can be found in any of the following ways.

a. Prostatic hyperplasia (benign prostatic hypertrophy) is listed as a diagnosis in *Concurrent Review Screening.* Select the contemplated surgery (TUR). The length of stay table in the operation volume is 648.

| 1<br>ADMITTING DIAGNOSIS | 2<br>Suggested Admission Category | 3<br>Your Hospital's Admission Category | 4<br>Length of Stay Group |
|---|---|---|---|
| Pregnancy, undelivered | | | |
| pre-eclampsia, during pregnancy | 3 | | 261 |
| Criteria for admission | | | |
| *1. Diastolic BP > 110* | | | |
| *2. Not controlled by outpatient Rx* | | | |
| *3.* | | | |
| *4.* | | | |
| premature labor, threatened | 1 | | 268 |
| premature rupture of membranes | 1 | | 268 |
| premature separation of placenta | 1 | | 265 |
| Pre-infarction syndrome | 1 | | 142 |
| Prepuce, redundant or phimosis | 2 | | 239 |
| Proctitis | 4 | | 211 |
| Prolonged labor, excludes uterine inertia | 1 | | 280 |
| **Prostatic hyperplasia** | 2 | | 235 |
| for: Cystoscopy | | | 636 |
| TUR | | | 648 |
| Other prostatectomy | | | 649 |
| Prostatitis | | | |
| acute | 3 | | 236 |
| Criteria for admission | | | |
| *1. Sepsis or fever > 102°, chills* | | | |
| *2. Abscess formation* | | | |

b. The Alphabetic Index at the back of the operation length of stay book also shows that TUR is included in operation group 648.

OPERATION INDEX (continued)

| Group Number | Title |
|---|---|
| 551 | Palatoplasty |
| 618 | Pancreatectomy |
| 618 | Pancreatotomy |
| | Paracentesis |
| 626 | Abdominal |
| 524 | Anterior chamber (eye) |
| 576 | Thoracic |
| 514 | Parathyroidectomy |
| 550 | Parotidectomy |
| 673 | Pelvic exenteration |
| 575 | Pericardiocentesis |
| 629 | Peritoneal dialysis |
| 628 | Peritoneoscopy |
| 557 | Pharyngoscopy |
| 556 | Pharyngotomy |
| 757 | Phlebography |
| 540 | Piercing, ear |
| 576 | Pleurocentesis |
| | Plication |
| 644 | Urethral, urethrovesical |
| 581 | Vena cava |
| 753 | Pneumoencephalography |
| 566 | Pneumonectomy |
| 659 | Preputiotomy |
| 608 | Proctectomy |
| 609 | Proctopexy |
| 619 | Proctosigmoidoscopy, except surgical approach |
| 609 | Surgical approach (transabdominal) |
| 607 | Proctotomy |
| 649 | Prostatectomy, except transurethral approach |
| 648 | Transurethral approach |
| 761 | Psychiatric testing and evaluation |
| 761 | Psychotherapy |
| | Puncture |
| 501 | Cranial |
| 512 | Lumbar |
| 501 | Skull, for drainage |
| 512 | Spinal |
| 758 | Pyelogram (intravenous) (retrograde) |
| 636 | Pyeloscopy |
| 632 | Pyelotomy |
| 590 | Pyloromyotomy |
| 596 | Pyloroplasty |

c. Transurethral resection can be found in Volume 3 of *ICD-9-CM* on page 408.




**Procedure** — *continued*
diagnostic NEC — *continued*
pleura 34.28
posterior chamber, eye 14.19
prostate 60.18
pulmonary 33.29
rectum 48.29
renal 55.29
respiratory 33.29
retina 14.19
retroperitoneum 59.29
salivary gland or duct 26.19
sclera 12.29
scrotum 61.19
seminal vesicle 60.19
sigmoid 45.28
sinus, nasal 22.19
skin 86.19
skull 01.19
soft tissue 83.29
spermatic cord 63.09
sphincter of Oddi 51.19
spine, spinal (canal) (cord) (meninges) (structure) 03.39
spleen 41.39
stomach 44.19
subcutaneous tissue 86.19
sympathetic nerve 05.19
tarsus 09.19
tendon NEC 83.29
extraocular 15.09
testicle 62.19
thalamus 01.18
thoracic duct 40.19
thorax 34.28
thymus 07.19
thyroid gland 06.19
tongue 25.09
tonsils 28.19
tooth 24.19
trachea 31.49
tunica vaginalis 61.19
ureter 56.39
urethra 58.29
uterus and supporting structures 68.19
uvula 27.29
vagina 70.29
vas deferens 63.09
vesical 57.39
vessel (blood) (any site) 38.29
vitreous 14.19
vulva 71.19
fistulizing, sclera NEC 12.69
miscellaneous (nonoperative) NEC 99.99
respiratory (nonoperative) NEC 93.99
surgical — *see* Operation
**Proctectasis** 96.22
**Proctectomy** (partial) *(see also* Resection, rectum) 48.69
abdominoperineal 48.5
**Proctectomy** (partial) — *continued*
complete (Miles) (Rankin) 48.5
pull-through 48.49
**Proctoclysis** 96.37
**Proctolysis** 48.99
**Proctopexy** (Delorme) 48.76
abdominal (Ripstein) 48.75
**Proctoplasty** 48.79
**Proctorrhaphy** 48.71
**Proctoscopy** 48.23
through stoma (artificial) 48.22
transabdominal approach 48.21
**Proctosigmoidectomy** *(see also* Resection, rectum) 48.69
**Proctosigmoidopexy** 48.76
**Proctosigmoidoscopy** 48.23
through stoma (artificial) 48.22
transabdominal approach 48.21
**Proctostomy** 48.1
**Proctotomy** (decompression) (linear) 48.0
**Production** — *see also* Formation
atrial septal defect 35.42
subcutaneous tunnel for esophageal anastomosis 42.86
with anastomosis — *see* Anastomosis, esophagus, antesternal
**Prognathic recession** 76.64
**Prophylaxis, dental** (scaling) (polishing) 96.54
**Prostatectomy** (complete) (partial) NEC 60.69
loop 60.2
perineal 60.62
radical (any approach) 60.5
retropubic (punch) (transcapsular) 60.4
suprapubic (punch) (transvesical) 60.3
transcapsular NEC 60.69
retropubic 60.4
**transurethral** 60.2
transvesical punch (suprapubic) 60.3
**Prostatocystotomy** 60.0
**Prostatolithotomy** 60.0
**Prostatotomy** (perineal) 60.0
**Prostatovesiculectomy** 60.5
**Protection** (of)
individual from his surroundings 99.84
surroundings from individual 99.84
**Psychoanalysis** 94.31
**Psychodrama** 94.43
**Psychotherapy** NEC 94.39
biofeedback 94.39
exploratory verbal 94.37
group 94.44
for psychosexual dysfunctions 94.41
play 94.36
psychosexual dysfunctions 94.34
supportive verbal 94.38
**Ptyalectasis** 26.91
**Ptyalithotomy** 26.0
**Ptyalolithotomy** 26.0

Step 2. Find the table for group 648 in the length of stay book.

**648: Transurethral prostatectomy (60.2)**

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 33 | 6.7 | 12 | 2 | 3 | 6 | 8 | 11 | 15 | 18 |
| 20-34 | 39 | 6.4 | 19 | 2 | 3 | 6 | 8 | 11 | 13 | 27 |
| 35-49 | 428 | 6.6 | 9 | 3 | 4 | 6 | 8 | 10 | 13 | 18 |
| 50-64 | 7705 | 7.0 | 10 | 4 | 4 | 6 | 8 | 10 | 12 | 18 |
| 65+ | 16192 | 8.1 | 18 | 4 | 5 | 7 | 9 | 13 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 9.6 | 33 | 3 | 4 | 8 | 13 | 16 | 23 | 30 |
| 20-34 | 78 | 13.3 | 270 | 2 | 4 | 8 | 14 | 27 | 49 | 94 |
| 35-49 | 679 | 8.8 | 45 | 3 | 4 | 7 | 10 | 15 | 20 | 36 |
| 50-64 | 13172 | 9.7 | 47 | 4 | 5 | 8 | 11 | 16 | 21 | 40 |
| 65+ | 42559 | 12.1 | 77 | 4 | 5 | 9 | 15 | 22 | 29 | 49 |
| **TOT. SINGLE DX** | 24397 | 7.8 | 17 | 4 | 4 | 7 | 9 | 12 | 15 | 22 |
| **" MULTIPLE DX** | 56542 | 11.5 | 72 | 4 | 5 | 9 | 14 | 21 | 28 | 47 |
| TOTAL 0-19 YRS | 87 | 8.5 | 28 | 2 | 4 | 7 | 10 | 15 | 19 | 29 |
| 20-34 | 117 | 11.0 | 197 | 2 | 3 | 7 | 11 | 21 | 32 | 90 |
| 35-49 | 1107 | 7.9 | 33 | 3 | 4 | 7 | 9 | 13 | 17 | 29 |
| 50-64 | 20877 | 8.7 | 35 | 4 | 4 | 7 | 10 | 14 | 18 | 34 |
| 65+ | 58751 | 11.0 | 65 | 4 | 5 | 9 | 13 | 20 | 26 | 44 |
| **GRAND TOTAL** | **80939** | **10.4** | **58** | **4** | **5** | **8** | **12** | **18** | **24** | **43** |

Step 3. There are no secondary diagnoses recorded, so attention is directed to the top five lines of the table (1. Single Dx).

**648: Transurethral prostatectomy (60.2)**

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 33 | 6.7 | 12 | 2 | 3 | 6 | 8 | 11 | 15 | 18 |
| 20-34 | 39 | 6.4 | 19 | 2 | 3 | 6 | 8 | 11 | 13 | 27 |
| 35-49 | 428 | 6.6 | 9 | 3 | 4 | 6 | 8 | 10 | 13 | 18 |
| 50-64 | 7705 | 7.0 | 10 | 4 | 4 | 6 | 8 | 10 | 12 | 18 |
| 65+ | 16192 | 8.1 | 18 | 4 | 5 | 7 | 9 | 13 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 9.6 | 33 | 3 | 4 | 8 | 13 | 16 | 23 | 30 |
| 20-34 | 78 | 13.3 | 270 | 2 | 4 | 8 | 14 | 27 | 49 | 94 |
| 35-49 | 679 | 8.8 | 45 | 3 | 4 | 7 | 10 | 15 | 20 | 36 |
| 50-64 | 13172 | 9.7 | 47 | 4 | 5 | 8 | 11 | 16 | 21 | 40 |
| 65+ | 42559 | 12.1 | 77 | 4 | 5 | 9 | 15 | 22 | 29 | 49 |
| **TOT. SINGLE DX** | 24397 | 7.8 | 17 | 4 | 4 | 7 | 9 | 12 | 15 | 22 |
| **" MULTIPLE DX** | 56542 | 11.5 | 72 | 4 | 5 | 9 | 14 | 21 | 28 | 47 |
| TOTAL 0-19 YRS | 87 | 8.5 | 28 | 2 | 4 | 7 | 10 | 15 | 19 | 29 |
| 20-34 | 117 | 11.0 | 197 | 2 | 3 | 7 | 11 | 21 | 32 | 90 |
| 35-49 | 1107 | 7.9 | 33 | 3 | 4 | 7 | 9 | 13 | 17 | 29 |
| 50-64 | 20877 | 8.7 | 35 | 4 | 4 | 7 | 10 | 14 | 18 | 34 |
| 65+ | 58751 | 11.0 | 65 | 4 | 5 | 9 | 13 | 20 | 26 | 44 |
| **GRAND TOTAL** | **80939** | **10.4** | **58** | **4** | **5** | **8** | **12** | **18** | **24** | **43** |

Step 4. The patient is 57 years old and therefore falls in the age group 50-64.

**648: Transurethral prostatectomy (60.2)**

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 33 | 6.7 | 12 | 2 | 3 | 6 | 8 | 11 | 15 | 18 |
| 20-34 | 39 | 6.4 | 19 | 2 | 3 | 6 | 8 | 11 | 13 | 27 |
| 35-49 | 428 | 6.6 | 9 | 3 | 4 | 6 | 8 | 10 | 13 | 18 |
| 50-64 | 7705 | 7.0 | 10 | 4 | 4 | 6 | 8 | 10 | 12 | 18 |
| 65+ | 16192 | 8.1 | 18 | 4 | 5 | 7 | 9 | 13 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 9.6 | 33 | 3 | 4 | 8 | 13 | 16 | 23 | 30 |
| 20-34 | 78 | 13.3 | 270 | 2 | 4 | 8 | 14 | 27 | 49 | 94 |
| 35-49 | 679 | 8.8 | 45 | 3 | 4 | 7 | 10 | 15 | 20 | 36 |
| 50-64 | 13172 | 9.7 | 47 | 4 | 5 | 8 | 11 | 16 | 21 | 40 |
| 65+ | 42559 | 12.1 | 77 | 4 | 5 | 9 | 15 | 22 | 29 | 49 |
| **TOT. SINGLE DX** | 24397 | 7.8 | 17 | 4 | 4 | 7 | 9 | 12 | 15 | 22 |
| **" MULTIPLE DX** | 56542 | 11.5 | 72 | 4 | 5 | 9 | 14 | 21 | 28 | 47 |
| TOTAL 0-19 YRS | 87 | 8.5 | 28 | 2 | 4 | 7 | 10 | 15 | 19 | 29 |
| 20-34 | 117 | 11.0 | 197 | 2 | 3 | 7 | 11 | 21 | 32 | 90 |
| 35-49 | 1107 | 7.9 | 33 | 3 | 4 | 7 | 9 | 13 | 17 | 29 |
| 50-64 | 20877 | 8.7 | 35 | 4 | 4 | 7 | 10 | 14 | 18 | 34 |
| 65+ | 58751 | 11.0 | 65 | 4 | 5 | 9 | 13 | 20 | 26 | 44 |
| **GRAND TOTAL** | **80939** | **10.4** | **58** | **4** | **5** | **8** | **12** | **18** | **24** | **43** |

Step 5. The 50th percentile day is the date assigned for the initial extended stay review. Column 7 shows the sixth day after admission to be the day that this patient comes up for extended stay review.

**648: Transurethral prostatectomy (60.2)**

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 33 | 6.7 | 12 | 2 | 3 | 6 | 8 | 11 | 15 | 18 |
| 20-34 | 39 | 6.4 | 19 | 2 | 3 | 6 | 8 | 11 | 13 | 27 |
| 35-49 | 428 | 6.6 | 9 | 3 | 4 | 6 | 8 | 10 | 13 | 18 |
| 50-64 | 7705 | 7.0 | 10 | 4 | 4 | 6 | 8 | 10 | 12 | 18 |
| 65+ | 16192 | 8.1 | 18 | 4 | 5 | 7 | 9 | 13 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 9.6 | 33 | 3 | 4 | 8 | 13 | 16 | 23 | 30 |
| 20-34 | 78 | 13.3 | 270 | 2 | 4 | 8 | 14 | 27 | 49 | 94 |
| 35-49 | 679 | 8.8 | 45 | 3 | 4 | 7 | 10 | 15 | 20 | 36 |
| 50-64 | 13172 | 9.7 | 47 | 4 | 5 | 8 | 11 | 16 | 21 | 40 |
| 65+ | 42559 | 12.1 | 77 | 4 | 5 | 9 | 15 | 22 | 29 | 49 |
| **TOT. SINGLE DX** | 24397 | 7.8 | 17 | 4 | 4 | 7 | 9 | 12 | 15 | 22 |
| **" MULTIPLE DX** | 56542 | 11.5 | 72 | 4 | 5 | 9 | 14 | 21 | 28 | 47 |
| TOTAL 0-19 YRS | 87 | 8.5 | 28 | 2 | 4 | 7 | 10 | 15 | 19 | 29 |
| 20-34 | 117 | 11.0 | 197 | 2 | 3 | 7 | 11 | 21 | 32 | 90 |
| 35-49 | 1107 | 7.9 | 33 | 3 | 4 | 7 | 9 | 13 | 17 | 29 |
| 50-64 | 20877 | 8.7 | 35 | 4 | 4 | 7 | 10 | 14 | 18 | 34 |
| 65+ | 58751 | 11.0 | 65 | 4 | 5 | 9 | 13 | 20 | 26 | 44 |
| **GRAND TOTAL** | **80939** | **10.4** | **58** | **4** | **5** | **8** | **12** | **18** | **24** | **43** |

The patient was admitted on 3 November; therefore the sixth day will fall on 9 November. Prior to or on 9 November the groups responsible for conducting extended stay review will make a finding as to whether further stay is medically necessary.

In the above example, the stay percentile used to define extended stay was the 50th percentile (column 7). If, based on the initial review, the patient's stay in the hospital was approved for extension beyond the 50th percentile, the date for the second extended stay review is determined in accordance with the hospital's plan, e.g., the 75th percentile.

Subsequent extended stay review dates should be set by the physician advisor or the utilization review committee as necessary based on the patient's medical condition and the attending physician's evaluation of the patient's progress.

# Length of Stay Tables

## SUMMARY OF ALL PATIENTS IN OPERATION GROUPS

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 73298 | 2.6 | 9 | <1 | <1 | 2 | 3 | 5 | 7 | 14 |
| 20-34 | 149187 | 2.9 | 8 | <1 | 1 | 2 | 4 | 5 | 7 | 13 |
| 35-49 | 62667 | 4.0 | 15 | <1 | 1 | 3 | 5 | 8 | 11 | 19 |
| 50-64 | 57134 | 4.6 | 20 | <1 | 1 | 3 | 6 | 10 | 12 | 21 |
| 65+ | 58932 | 5.3 | 29 | 1 | 1 | 3 | 7 | 12 | 15 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 69952 | 4.6 | 54 | <1 | 1 | 3 | 5 | 9 | 15 | 43 |
| 20-34 | 191034 | 4.3 | 28 | 1 | 1 | 3 | 5 | 7 | 11 | 28 |
| 35-49 | 99405 | 6.3 | 48 | 1 | 1 | 5 | 7 | 12 | 17 | 36 |
| 50-64 | 127110 | 7.7 | 69 | 1 | 2 | 5 | 10 | 16 | 22 | 44 |
| 65+ | 211075 | 9.8 | 94 | 2 | 2 | 7 | 12 | 20 | 28 | 52 |
| **TOT. SINGLE DX** | 401218 | 3.6 | 15 | <1 | 1 | 2 | 4 | 7 | 10 | 19 |
| **" MULTIPLE DX** | 698576 | 6.9 | 66 | 1 | 1 | 4 | 8 | 15 | 21 | 44 |
| TOTAL 0-19 YRS | 143250 | 3.6 | 32 | <1 | 1 | 2 | 4 | 7 | 10 | 30 |
| 20-34 | 340221 | 3.7 | 19 | 1 | 1 | 3 | 4 | 7 | 9 | 22 |
| 35-49 | 162072 | 5.4 | 37 | <1 | 1 | 4 | 7 | 10 | 15 | 31 |
| 50-64 | 184244 | 6.8 | 55 | 1 | 1 | 5 | 8 | 14 | 19 | 39 |
| 65+ | 270007 | 8.8 | 83 | 1 | 2 | 6 | 11 | 18 | 25 | 49 |
| **GRAND TOTAL** | **1099794** | **5.7** | **50** | **1** | **1** | **3** | **7** | **12** | **18** | **37** |

## 501: Incision, excision of skull, brain, and cerebral meninges (01.01-01.15; 01.3-01.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 67 | 8.5 | 47 | 1 | 1 | 7 | 11 | 16 | 21 | 34 |
| 20-34 | 86 | 11.7 | 95 | 4 | 5 | 9 | 14 | 19 | 21 | 67 |
| 35-49 | 110 | 11.0 | 49 | 2 | 3 | 10 | 14 | 21 | 26 | 34 |
| 50-64 | 147 | 12.0 | 59 | 3 | 5 | 11 | 15 | 19 | 26 | 57 |
| 65+ | 137 | 14.1 | 95 | 4 | 6 | 11 | 18 | 26 | 31 | 46 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 156 | 17.5 | 313 | 1 | 3 | 13 | 23 | 55 | 94 | 99+ |
| 20-34 | 144 | 19.7 | 272 | 4 | 6 | 15 | 24 | 48 | 66 | 99+ |
| 35-49 | 171 | 20.9 | 343 | 4 | 7 | 14 | 27 | 55 | 78 | 99+ |
| 50-64 | 262 | 21.5 | 314 | 5 | 7 | 16 | 27 | 49 | 69 | 99+ |
| 65+ | 407 | 25.6 | 345 | 7 | 8 | 21 | 35 | 55 | 75 | 99+ |
| **TOT. SINGLE DX** | 547 | 11.8 | 73 | 2 | 4 | 10 | 15 | 21 | 27 | 41 |
| **" MULTIPLE DX** | 1140 | 22.1 | 332 | 4 | 6 | 16 | 29 | 53 | 74 | 99+ |
| TOTAL 0-19 YRS | 223 | 14.8 | 250 | 1 | 2 | 10 | 20 | 42 | 64 | 99+ |
| 20-34 | 230 | 16.7 | 221 | 4 | 6 | 12 | 20 | 34 | 57 | 99+ |
| 35-49 | 281 | 17.0 | 251 | 3 | 5 | 12 | 21 | 38 | 60 | 99+ |
| 50-64 | 409 | 18.0 | 243 | 5 | 6 | 13 | 22 | 42 | 56 | 99+ |
| 65+ | 544 | 22.7 | 307 | 5 | 7 | 17 | 29 | 49 | 66 | 99+ |
| **GRAND TOTAL** | **1687** | **18.8** | **271** | **3** | **5** | **14** | **23** | **43** | **61** | **99+** |

## 502: Craniotomy and craniectomy (01.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 36 | 9.6 | 85 | 2 | 3 | 7 | 10 | 24 | 30 | 47 |
| 20-34 | 55 | 10.7 | 126 | 2 | 3 | 9 | 12 | 17 | 22 | 81 |
| 35-49 | 40 | 11.3 | 69 | 3 | 4 | 9 | 13 | 29 | 36 | 99+ |
| 50-64 | 58 | 12.4 | 59 | 5 | 6 | 10 | 15 | 22 | 31 | 40 |
| 65+ | 49 | 12.1 | 87 | 3 | 4 | 11 | 14 | 18 | 26 | 51 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 90 | 15.0 | 275 | 3 | 4 | 9 | 17 | 40 | 68 | 99+ |
| 20-34 | 129 | 20.6 | 421 | 4 | 5 | 13 | 25 | 65 | 87 | 99+ |
| 35-49 | 107 | 20.1 | 430 | 3 | 4 | 11 | 28 | 59 | 66 | 98 |
| 50-64 | 120 | 22.0 | 250 | 7 | 8 | 17 | 30 | 50 | 66 | 99+ |
| 65+ | 190 | 23.3 | 324 | 5 | 7 | 17 | 30 | 50 | 63 | 96 |
| **TOT. SINGLE DX** | 238 | 11.3 | 86 | 3 | 4 | 9 | 13 | 20 | 30 | 55 |
| **" MULTIPLE DX** | 636 | 20.8 | 346 | 4 | 5 | 14 | 28 | 54 | 68 | 99+ |
| TOTAL 0-19 YRS | 126 | 13.5 | 226 | 3 | 3 | 8 | 16 | 32 | 55 | 99+ |
| 20-34 | 184 | 17.6 | 352 | 3 | 5 | 11 | 20 | 54 | 72 | 99+ |
| 35-49 | 147 | 17.7 | 347 | 3 | 4 | 10 | 21 | 48 | 64 | 99+ |
| 50-64 | 178 | 18.9 | 208 | 6 | 7 | 14 | 24 | 43 | 61 | 99+ |
| 65+ | 239 | 21.0 | 296 | 4 | 6 | 16 | 27 | 46 | 58 | 94 |
| **GRAND TOTAL** | **874** | **18.2** | **292** | **4** | **5** | **12** | **23** | **44** | **64** | **99+** |

## 503: Cranioplasty (02.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 96 | 5.0 | 8 | 1 | 2 | 5 | 6 | 8 | 8 | 18 |
| 20-34 | 44 | 5.4 | 10 | 2 | 2 | 5 | 8 | 10 | 11 | 13 |
| 35-49 | 25 | 6.5 | 10 | 3 | 3 | 6 | 8 | 10 | 12 | 17 |
| 50-64 | 18 | 9.1 | 56 | 3 | 3 | 7 | 10 | 22 | 29 | 31 |
| 65+ | 9 | 7.1 | 22 | 3 | 3 | 5 | 10 | 15 | 15 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 129 | 7.1 | 39 | 2 | 3 | 6 | 8 | 14 | 19 | 63 |
| 20-34 | 105 | 12.3 | 276 | 3 | 3 | 6 | 12 | 43 | 74 | 99+ |
| 35-49 | 44 | 11.3 | 213 | 3 | 3 | 7 | 13 | 21 | 35 | 88 |
| 50-64 | 41 | 11.0 | 73 | 3 | 4 | 7 | 16 | 25 | 28 | 35 |
| 65+ | 11 | 11.4 | 110 | 3 | 4 | 8 | 14 | 59 | 94 | 99+ |
| **TOT. SINGLE DX** | 192 | 5.8 | 15 | 2 | 2 | 5 | 7 | 10 | 12 | 24 |
| **" MULTIPLE DX** | 330 | 9.9 | 148 | 3 | 3 | 6 | 11 | 21 | 42 | 99+ |
| TOTAL 0-19 YRS | 225 | 6.2 | 27 | 2 | 2 | 5 | 7 | 11 | 16 | 33 |
| 20-34 | 149 | 10.2 | 207 | 2 | 3 | 6 | 10 | 24 | 67 | 99+ |
| 35-49 | 69 | 9.6 | 144 | 3 | 3 | 6 | 10 | 17 | 26 | 80 |
| 50-64 | 59 | 10.4 | 68 | 3 | 4 | 7 | 15 | 25 | 28 | 35 |
| 65+ | 20 | 9.5 | 72 | 3 | 3 | 8 | 13 | 26 | 67 | 99+ |
| **GRAND TOTAL** | **522** | **8.4** | **103** | **2** | **3** | **6** | **8** | **17** | **26** | **91** |

## 504: Ventricular shunt, extracranial (02.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 33 | 7.0 | 23 | 2 | 3 | 6 | 8 | 12 | 12 | 28 |
| 20-34 | 19 | 9.2 | 14 | 4 | 4 | 8 | 13 | 15 | 15 | 15 |
| 35-49 | 10 | 7.9 | 7 | 5 | 5 | 7 | 11 | 12 | 13 | 13 |
| 50-64 | 19 | 8.5 | 29 | 3 | 4 | 8 | 9 | 18 | 21 | 23 |
| 65+ | 33 | 12.8 | 178 | 3 | 4 | 10 | 14 | 18 | 49 | 69 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 90 | 19.9 | 452 | 4 | 5 | 9 | 27 | 63 | 78 | 99+ |
| 20-34 | 37 | 22.1 | 553 | 6 | 6 | 10 | 33 | 56 | 77 | 99 |
| 35-49 | 27 | 17.3 | 322 | 4 | 5 | 10 | 19 | 39 | 59 | 82 |
| 50-64 | 43 | 21.5 | 201 | 5 | 7 | 18 | 29 | 42 | 59 | 99+ |
| 65+ | 125 | 27.3 | 301 | 7 | 8 | 24 | 37 | 56 | 68 | 99+ |
| **TOT. SINGLE DX** | 114 | 9.4 | 70 | 3 | 4 | 8 | 11 | 15 | 18 | 59 |
| **" MULTIPLE DX** | 322 | 23.0 | 369 | 5 | 6 | 17 | 32 | 55 | 70 | 99+ |
| TOTAL 0-19 YRS | 123 | 16.4 | 369 | 3 | 4 | 9 | 23 | 45 | 71 | 99+ |
| 20-34 | 56 | 17.7 | 405 | 4 | 6 | 10 | 16 | 50 | 61 | 98 |
| 35-49 | 37 | 14.8 | 253 | 5 | 5 | 9 | 14 | 37 | 49 | 82 |
| 50-64 | 62 | 17.5 | 184 | 4 | 5 | 14 | 27 | 38 | 47 | 96 |
| 65+ | 158 | 24.3 | 310 | 5 | 7 | 21 | 33 | 52 | 66 | 99 |
| **GRAND TOTAL** | **436** | **19.5** | **327** | **4** | **5** | **13** | **26** | **46** | **64** | **99+** |

## 505: Miscellaneous operations on brain and cerebral meninges (02.1-02.2; 02.4-02.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 56 | 6.1 | 21 | 1 | 2 | 5 | 8 | 13 | 14 | 24 |
| 20-34 | 14 | 6.6 | 31 | 2 | 2 | 5 | 8 | 16 | 21 | 22 |
| 35-49 | 18 | 7.6 | 12 | 4 | 4 | 6 | 12 | 12 | 13 | 14 |
| 50-64 | 17 | 11.4 | 242 | 1 | 2 | 6 | 10 | 28 | 52 | 64 |
| 65+ | 20 | 8.5 | 65 | 1 | 1 | 5 | 12 | 23 | 25 | 27 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 231 | 11.8 | 237 | 2 | 3 | 6 | 13 | 32 | 55 | 99+ |
| 20-34 | 106 | 19.0 | 381 | 4 | 5 | 11 | 27 | 57 | 91 | 99+ |
| 35-49 | 53 | 11.8 | 104 | 2 | 3 | 8 | 15 | 28 | 39 | 98 |
| 50-64 | 51 | 15.5 | 149 | 4 | 4 | 11 | 23 | 36 | 42 | 44 |
| 65+ | 87 | 23.7 | 401 | 3 | 3 | 16 | 31 | 60 | 66 | 80 |
| **TOT. SINGLE DX** | 125 | 7.5 | 59 | 1 | 2 | 5 | 9 | 15 | 22 | 38 |
| **" MULTIPLE DX** | 528 | 15.6 | 291 | 2 | 3 | 9 | 21 | 43 | 62 | 99+ |
| TOTAL 0-19 YRS | 287 | 10.7 | 201 | 2 | 2 | 6 | 11 | 27 | 48 | 99 |
| 20-34 | 120 | 17.5 | 356 | 3 | 4 | 10 | 25 | 51 | 89 | 99+ |
| 35-49 | 71 | 10.7 | 84 | 3 | 3 | 8 | 13 | 25 | 38 | 88 |
| 50-64 | 68 | 14.5 | 173 | 3 | 3 | 9 | 21 | 36 | 42 | 60 |
| 65+ | 107 | 20.9 | 373 | 2 | 3 | 15 | 28 | 55 | 65 | 79 |
| **GRAND TOTAL** | **653** | **14.0** | **256** | **2** | **3** | **8** | **17** | **37** | **57** | **99+** |

## 506: Exploration and decompression of spinal canal structures (03.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 18 | 9.3 | 16 | 4 | 4 | 9 | 12 | 15 | 17 | 18 |
| 20-34 | 183 | 9.4 | 33 | 4 | 4 | 8 | 11 | 15 | 18 | 32 |
| 35-49 | 332 | 9.6 | 34 | 4 | 5 | 8 | 11 | 16 | 19 | 28 |
| 50-64 | 358 | 10.2 | 40 | 5 | 6 | 9 | 12 | 15 | 18 | 30 |
| 65+ | 194 | 12.3 | 43 | 5 | 6 | 10 | 14 | 21 | 27 | 37 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 16.6 | 166 | 3 | 5 | 13 | 32 | 83 | 99+ | 99+ |
| 20-34 | 192 | 15.3 | 195 | 4 | 5 | 11 | 18 | 30 | 49 | 86 |
| 35-49 | 350 | 12.8 | 90 | 4 | 5 | 10 | 15 | 23 | 30 | 57 |
| 50-64 | 601 | 15.7 | 166 | 5 | 6 | 12 | 17 | 32 | 44 | 73 |
| 65+ | 583 | 19.1 | 209 | 6 | 8 | 14 | 23 | 37 | 47 | 79 |
| **TOT. SINGLE DX** | 1085 | 10.2 | 39 | 4 | 5 | 9 | 12 | 16 | 21 | 31 |
| **" MULTIPLE DX** | 1742 | 16.2 | 174 | 5 | 6 | 12 | 19 | 32 | 45 | 78 |
| TOTAL 0-19 YRS | 34 | 12.7 | 98 | 4 | 5 | 9 | 17 | 33 | 83 | 99+ |
| 20-34 | 375 | 12.4 | 125 | 4 | 5 | 9 | 14 | 23 | 33 | 69 |
| 35-49 | 682 | 11.3 | 66 | 4 | 5 | 9 | 13 | 19 | 26 | 48 |
| 50-64 | 959 | 13.6 | 126 | 5 | 6 | 10 | 15 | 24 | 36 | 67 |
| 65+ | 777 | 17.4 | 177 | 6 | 7 | 13 | 21 | 33 | 44 | 76 |
| **GRAND TOTAL** | **2827** | **13.9** | **130** | **5** | **6** | **11** | **16** | **26** | **36** | **71** |

## 507: Operations on spinal cord structures, except exploration (03.1-03.2; 03.32; 03.4-03.8; 03.93-03.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 32 | 7.5 | 27 | <1 | 1 | 7 | 11 | 16 | 18 | 19 |
| 20-34 | 80 | 6.6 | 53 | 1 | 1 | 3 | 10 | 16 | 23 | 33 |
| 35-49 | 91 | 6.0 | 28 | 1 | 1 | 3 | 9 | 14 | 16 | 23 |
| 50-64 | 53 | 7.3 | 41 | 1 | 1 | 4 | 13 | 17 | 20 | 22 |
| 65+ | 34 | 10.0 | 60 | 1 | 2 | 9 | 12 | 23 | 28 | 29 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 51 | 13.7 | 184 | 2 | 3 | 8 | 20 | 40 | 65 | 99+ |
| 20-34 | 137 | 13.9 | 199 | 1 | 2 | 10 | 19 | 35 | 45 | 75 |
| 35-49 | 128 | 11.4 | 111 | 2 | 2 | 8 | 15 | 24 | 35 | 54 |
| 50-64 | 146 | 13.0 | 133 | 2 | 2 | 10 | 18 | 28 | 34 | 54 |
| 65+ | 94 | 18.5 | 242 | 2 | 4 | 15 | 24 | 42 | 53 | 99+ |
| **TOT. SINGLE DX** | 290 | 7.0 | 42 | 1 | 1 | 4 | 11 | 16 | 20 | 27 |
| **" MULTIPLE DX** | 556 | 13.8 | 172 | 1 | 2 | 10 | 19 | 32 | 43 | 90 |
| TOTAL 0-19 YRS | 83 | 11.3 | 132 | 1 | 2 | 8 | 14 | 30 | 42 | 99+ |
| 20-34 | 217 | 11.2 | 157 | 1 | 1 | 7 | 15 | 29 | 37 | 64 |
| 35-49 | 219 | 9.1 | 84 | 1 | 2 | 7 | 12 | 20 | 25 | 51 |
| 50-64 | 199 | 11.5 | 115 | 1 | 2 | 8 | 16 | 25 | 30 | 54 |
| 65+ | 128 | 16.2 | 208 | 2 | 3 | 13 | 23 | 37 | 47 | 99+ |
| **GRAND TOTAL** | **846** | **11.5** | **138** | **1** | **2** | **8** | **15** | **26** | **37** | **68** |

## 508: Excision or destruction of lesion of cranial or peripheral nerve (04.0-04.12)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 35 | 2.3 | 4 | <1 | <1 | 2 | 3 | 4 | 8 | 11 |
| 20-34 | 202 | 2.8 | 8 | <1 | <1 | 2 | 3 | 6 | 9 | 14 |
| 35-49 | 229 | 2.9 | 9 | <1 | <1 | 2 | 3 | 6 | 9 | 18 |
| 50-64 | 318 | 2.8 | 6 | <1 | <1 | 2 | 3 | 6 | 9 | 12 |
| 65+ | 161 | 3.1 | 8 | <1 | <1 | 2 | 3 | 7 | 10 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 30 | 5.0 | 31 | <1 | <1 | 3 | 7 | 10 | 11 | 29 |
| 20-34 | 156 | 4.8 | 23 | <1 | 1 | 3 | 6 | 12 | 15 | 22 |
| 35-49 | 216 | 5.8 | 53 | <1 | 1 | 3 | 8 | 13 | 17 | 31 |
| 50-64 | 252 | 7.0 | 74 | 1 | 1 | 5 | 9 | 15 | 22 | 67 |
| 65+ | 136 | 7.2 | 91 | 1 | 1 | 4 | 8 | 15 | 28 | 44 |
| **TOT. SINGLE DX** | 945 | 2.9 | 8 | <1 | <1 | 2 | 3 | 6 | 9 | 14 |
| **" MULTIPLE DX** | 790 | 6.2 | 61 | 1 | 1 | 4 | 8 | 13 | 19 | 47 |
| TOTAL 0-19 YRS | 65 | 3.5 | 19 | <1 | <1 | 2 | 3 | 9 | 11 | 26 |
| 20-34 | 358 | 3.7 | 16 | <1 | <1 | 2 | 4 | 9 | 12 | 21 |
| 35-49 | 445 | 4.3 | 33 | <1 | <1 | 2 | 4 | 10 | 14 | 29 |
| 50-64 | 570 | 4.7 | 41 | <1 | 1 | 3 | 6 | 10 | 14 | 49 |
| 65+ | 297 | 5.0 | 51 | <1 | 1 | 3 | 5 | 10 | 15 | 38 |
| **GRAND TOTAL** | **1735** | **4.4** | **35** | **<1** | **1** | **2** | **5** | **10** | **13** | **29** |

## 509: Miscellaneous operations on cranial and other peripheral nerves (04.2-04.7; 04.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 132 | 2.2 | 5 | <1 | <1 | 2 | 2 | 4 | 5 | 12 |
| 20-34 | 836 | 1.9 | 1 | <1 | <1 | 2 | 2 | 3 | 5 | 7 |
| 35-49 | 1071 | 1.9 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 7 |
| 50-64 | 947 | 2.1 | 4 | <1 | <1 | 2 | 2 | 3 | 5 | 9 |
| 65+ | 594 | 2.2 | 2 | <1 | <1 | 2 | 3 | 4 | 5 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 157 | 3.0 | 37 | <1 | 1 | 2 | 3 | 5 | 7 | 15 |
| 20-34 | 540 | 2.7 | 9 | <1 | 1 | 2 | 3 | 5 | 7 | 16 |
| 35-49 | 514 | 3.4 | 21 | <1 | 1 | 2 | 3 | 6 | 9 | 25 |
| 50-64 | 637 | 3.3 | 13 | <1 | 1 | 2 | 4 | 6 | 10 | 18 |
| 65+ | 496 | 4.6 | 52 | 1 | 1 | 2 | 4 | 9 | 15 | 54 |
| **TOT. SINGLE DX** | 3580 | 2.0 | 3 | <1 | <1 | 2 | 2 | 3 | 4 | 8 |
| **" MULTIPLE DX** | 2344 | 3.4 | 25 | <1 | 1 | 2 | 3 | 6 | 10 | 23 |
| TOTAL 0-19 YRS | 289 | 2.6 | 23 | <1 | <1 | 2 | 3 | 4 | 6 | 14 |
| 20-34 | 1376 | 2.2 | 5 | <1 | <1 | 2 | 3 | 4 | 5 | 10 |
| 35-49 | 1585 | 2.4 | 8 | <1 | <1 | 2 | 3 | 4 | 5 | 12 |
| 50-64 | 1584 | 2.6 | 8 | <1 | <1 | 2 | 3 | 4 | 7 | 14 |
| 65+ | 1090 | 3.3 | 27 | <1 | 1 | 2 | 3 | 6 | 9 | 23 |
| **GRAND TOTAL** | **5924** | **2.6** | **12** | **<1** | **<1** | **2** | **3** | **4** | **6** | **15** |

## 510: Sympathectomy (05.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 20-34 | 23 | 4.3 | 6 | <1 | 1 | 4 | 6 | 7 | 9 | 10 |
| 35-49 | 18 | 5.5 | 18 | <1 | 1 | 5 | 7 | 13 | 15 | 16 |
| 50-64 | 16 | 6.8 | 37 | <1 | <1 | 5 | 10 | 16 | 21 | 23 |
| 65+ | 22 | 10.5 | 56 | 2 | 5 | 9 | 12 | 20 | 27 | 36 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 4.5 | 13 | 2 | 2 | 4 | 7 | 7 | 7 | 7 |
| 20-34 | 19 | 5.1 | 4 | 2 | 3 | 5 | 6 | 8 | 9 | 10 |
| 35-49 | 34 | 11.4 | 119 | 2 | 3 | 8 | 12 | 29 | 39 | 47 |
| 50-64 | 45 | 12.6 | 171 | 5 | 6 | 9 | 13 | 25 | 29 | 87 |
| 65+ | 130 | 15.5 | 179 | 4 | 5 | 12 | 18 | 29 | 48 | 88 |
| **TOT. SINGLE DX** | 80 | 6.8 | 34 | <1 | 1 | 5 | 9 | 13 | 18 | 32 |
| **" MULTIPLE DX** | 230 | 13.4 | 161 | 3 | 4 | 9 | 16 | 27 | 40 | 85 |
| TOTAL 0-19 YRS | 3 | 4.7 | 6 | 2 | 2 | 5 | 6 | 7 | 7 | 7 |
| 20-34 | 42 | 4.6 | 5 | 1 | 1 | 5 | 6 | 8 | 9 | 10 |
| 35-49 | 52 | 9.3 | 91 | 1 | 2 | 7 | 11 | 21 | 31 | 47 |
| 50-64 | 61 | 11.1 | 142 | 1 | 4 | 8 | 12 | 21 | 26 | 81 |
| 65+ | 152 | 14.8 | 164 | 4 | 5 | 11 | 18 | 28 | 46 | 85 |
| **GRAND TOTAL** | **310** | **11.7** | **137** | **1** | **3** | **8** | **14** | **23** | **36** | **78** |

## 511: Miscellaneous operations on sympathetic nerves and ganglia (05.0-05.11; 05.3-05.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 2.5 | 1 | 2 | 2 | 2 | 3 | 3 | 3 | 3 |
| 20-34 | 15 | 6.9 | 40 | 1 | 1 | 5 | 10 | 17 | 21 | 22 |
| 35-49 | 14 | 5.5 | 19 | <1 | 1 | 3 | 9 | 11 | 14 | 15 |
| 50-64 | 19 | 5.4 | 63 | <1 | <1 | 3 | 6 | 8 | 24 | 37 |
| 65+ | 5 | 4.4 | 16 | 1 | 1 | 3 | 6 | 11 | 11 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 1.1 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| 20-34 | 16 | 6.9 | 67 | <1 | 1 | 3 | 9 | 23 | 23 | 23 |
| 35-49 | 39 | 9.2 | 93 | <1 | 1 | 5 | 15 | 26 | 29 | 33 |
| 50-64 | 32 | 8.1 | 95 | 1 | 2 | 6 | 10 | 16 | 18 | 56 |
| 65+ | 36 | 10.4 | 91 | 2 | 2 | 8 | 14 | 23 | 25 | 49 |
| **TOT. SINGLE DX** | 55 | 5.6 | 38 | <1 | 1 | 3 | 8 | 11 | 16 | 36 |
| **" MULTIPLE DX** | 130 | 8.6 | 88 | 1 | 1 | 5 | 11 | 23 | 26 | 50 |
| TOTAL 0-19 YRS | 9 | 1.4 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 31 | 6.9 | 52 | 1 | 1 | 3 | 10 | 21 | 23 | 23 |
| 35-49 | 53 | 8.2 | 76 | <1 | 1 | 4 | 10 | 25 | 27 | 33 |
| 50-64 | 51 | 7.1 | 84 | <1 | 1 | 5 | 8 | 13 | 18 | 56 |
| 65+ | 41 | 9.7 | 86 | 1 | 2 | 7 | 13 | 22 | 24 | 49 |
| **GRAND TOTAL** | **185** | **7.7** | **74** | **1** | **1** | **5** | **10** | **21** | **25** | **45** |

## 512: Miscellaneous diagnostic and therapeutic procedures on nervous system (01.18-01.19; 03.31; 03.39; 03.91-03.92; 04.19; 04.8; 05.19)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 513: Thyroidectomy (06.2-06.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 69 | 3.3 | 1 | 1 | 2 | 3 | 4 | 5 | 5 | 7 |
| 20-34 | 548 | 3.6 | 1 | 2 | 2 | 4 | 4 | 5 | 6 | 8 |
| 35-49 | 548 | 3.6 | 2 | 2 | 2 | 3 | 4 | 5 | 6 | 8 |
| 50-64 | 438 | 3.7 | 1 | 2 | 2 | 4 | 4 | 5 | 6 | 9 |
| 65+ | 200 | 4.3 | 4 | 2 | 3 | 4 | 5 | 6 | 7 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 30 | 5.5 | 10 | 2 | 3 | 4 | 6 | 10 | 14 | 16 |
| 20-34 | 238 | 4.9 | 44 | 2 | 3 | 4 | 5 | 7 | 9 | 19 |
| 35-49 | 284 | 4.6 | 7 | 2 | 3 | 4 | 5 | 7 | 9 | 17 |
| 50-64 | 311 | 4.8 | 8 | 2 | 3 | 4 | 5 | 8 | 10 | 18 |
| 65+ | 204 | 6.3 | 39 | 2 | 3 | 5 | 6 | 13 | 16 | 21 |
| **TOT. SINGLE DX** | 1803 | 3.7 | 2 | 2 | 2 | 4 | 4 | 5 | 6 | 8 |
| **" MULTIPLE DX** | 1067 | 5.1 | 22 | 2 | 3 | 4 | 5 | 8 | 11 | 19 |
| TOTAL 0-19 YRS | 99 | 3.9 | 5 | 1 | 2 | 4 | 4 | 6 | 8 | 15 |
| 20-34 | 786 | 4.0 | 15 | 2 | 2 | 4 | 4 | 5 | 7 | 10 |
| 35-49 | 832 | 3.9 | 4 | 2 | 2 | 4 | 4 | 6 | 7 | 12 |
| 50-64 | 749 | 4.2 | 5 | 2 | 3 | 4 | 5 | 6 | 8 | 13 |
| 65+ | 404 | 5.3 | 23 | 2 | 3 | 4 | 5 | 8 | 13 | 19 |
| **GRAND TOTAL** | **2870** | **4.2** | **10** | **2** | **2** | **4** | **5** | **6** | **8** | **16** |

## 514: Operations on parathyroid and thyroid, except thyroidectomy (06.01-06.13; 06.7-06.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 138 | 1.9 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 20-34 | 109 | 2.9 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 75 | 3.7 | 5 | 1 | 2 | 3 | 5 | 6 | 9 | 13 |
| 50-64 | 72 | 3.8 | 3 | 2 | 2 | 4 | 4 | 6 | 7 | 11 |
| 65+ | 32 | 4.4 | 3 | 1 | 2 | 4 | 6 | 7 | 7 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 26 | 3.2 | 9 | <1 | 1 | 2 | 3 | 7 | 10 | 14 |
| 20-34 | 69 | 6.7 | 64 | 1 | 1 | 4 | 7 | 14 | 19 | 48 |
| 35-49 | 71 | 7.0 | 52 | 2 | 2 | 5 | 8 | 11 | 26 | 38 |
| 50-64 | 121 | 7.3 | 35 | 3 | 3 | 5 | 9 | 15 | 23 | 99+ |
| 65+ | 106 | 10.3 | 156 | 3 | 3 | 6 | 10 | 25 | 33 | 71 |
| **TOT. SINGLE DX** | 426 | 3.0 | 4 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| **" MULTIPLE DX** | 393 | 7.7 | 77 | 1 | 2 | 5 | 9 | 15 | 27 | 53 |
| TOTAL 0-19 YRS | 164 | 2.1 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 178 | 4.3 | 30 | 1 | 1 | 3 | 5 | 8 | 11 | 31 |
| 35-49 | 146 | 5.3 | 31 | 1 | 2 | 4 | 6 | 9 | 12 | 36 |
| 50-64 | 193 | 6.0 | 27 | 2 | 3 | 4 | 7 | 11 | 17 | 71 |
| 65+ | 138 | 9.0 | 127 | 2 | 3 | 6 | 9 | 18 | 30 | 58 |
| **GRAND TOTAL** | **819** | **5.2** | **44** | **1** | **1** | **4** | **6** | **10** | **15** | **39** |

## 515: Operations on endocrine glands, except parathyroid and thyroid (07.00-07.17; 07.2-07.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 8.1 | 26 | 2 | 3 | 8 | 11 | 15 | 19 | 21 |
| 20-34 | 39 | 7.6 | 9 | 4 | 4 | 7 | 9 | 11 | 12 | 20 |
| 35-49 | 38 | 9.6 | 50 | 5 | 5 | 8 | 11 | 12 | 23 | 44 |
| 50-64 | 33 | 9.3 | 51 | 4 | 5 | 8 | 9 | 11 | 20 | 45 |
| 65+ | 14 | 12.1 | 61 | 3 | 6 | 9 | 16 | 19 | 31 | 35 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 11.6 | 51 | 3 | 5 | 8 | 16 | 23 | 25 | 25 |
| 20-34 | 44 | 9.4 | 26 | 3 | 5 | 8 | 11 | 19 | 24 | 99+ |
| 35-49 | 56 | 16.8 | 195 | 5 | 6 | 11 | 19 | 40 | 49 | 67 |
| 50-64 | 54 | 13.1 | 48 | 6 | 7 | 12 | 15 | 23 | 30 | 34 |
| 65+ | 41 | 19.5 | 270 | 6 | 7 | 14 | 26 | 49 | 91 | 99+ |
| **TOT. SINGLE DX** | 143 | 9.0 | 38 | 3 | 4 | 8 | 10 | 12 | 18 | 44 |
| **" MULTIPLE DX** | 209 | 14.5 | 139 | 5 | 6 | 11 | 18 | 28 | 45 | 99+ |
| TOTAL 0-19 YRS | 33 | 9.5 | 39 | 2 | 3 | 8 | 13 | 20 | 23 | 25 |
| 20-34 | 83 | 8.6 | 19 | 4 | 5 | 8 | 10 | 13 | 20 | 74 |
| 35-49 | 94 | 13.9 | 148 | 5 | 6 | 9 | 15 | 28 | 46 | 64 |
| 50-64 | 87 | 11.7 | 52 | 5 | 6 | 9 | 13 | 22 | 28 | 41 |
| 65+ | 55 | 17.6 | 226 | 5 | 7 | 12 | 24 | 37 | 79 | 99+ |
| **GRAND TOTAL** | **352** | **12.3** | **104** | **4** | **5** | **9** | **14** | **24** | **34** | **86** |

## 516: Miscellaneous diagnostic and therapeutic procedures on endocrine system (06.19; 07.19)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 517: Excision or destruction of lesion or tissue of eyelid (08.11; 08.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 106 | 1.2 | | ≤1 | ≤1 | <1 | 1 | 2 | 2 | 5 |
| 20-34 | 14 | 1.6 | | ≤1 | ≤1 | 1 | 2 | 3 | 3 | 3 |
| 35-49 | 21 | 1.3 | | ≤1 | ≤1 | 1 | 2 | 2 | 2 | 2 |
| 50-64 | 38 | 2.3 | 3 | ≤1 | ≤1 | 2 | 2 | 5 | 7 | 9 |
| 65+ | 68 | 2.2 | 6 | ≤1 | ≤1 | 2 | 2 | 3 | 4 | 19 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 11 | 2.0 | 9 | ≤1 | ≤1 | <1 | 1 | 6 | 11 | 11 |
| 20-34 | 9 | 1.7 | 1 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 4 |
| 35-49 | 26 | 5.3 | 230 | ≤1 | ≤1 | 2 | 3 | 8 | 24 | 79 |
| 50-64 | 33 | 3.6 | 31 | ≤1 | ≤1 | 2 | 3 | 9 | 11 | 31 |
| 65+ | 76 | 4.8 | 62 | ≤1 | ≤1 | 2 | 5 | 9 | 19 | 44 |
| **TOT. SINGLE DX** | 247 | 1.7 | 3 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 9 |
| **" MULTIPLE DX** | 155 | 4.3 | 76 | ≤1 | ≤1 | 2 | 3 | 9 | 11 | 46 |
| TOTAL 0-19 YRS | 117 | 1.3 | 1 | ≤1 | ≤1 | <1 | 1 | 2 | 2 | 8 |
| 20-34 | 23 | 1.6 | 1 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 4 |
| 35-49 | 47 | 3.6 | 129 | ≤1 | ≤1 | 1 | 2 | 4 | 8 | 79 |
| 50-64 | 71 | 2.9 | 17 | ≤1 | ≤1 | 2 | 3 | 6 | 9 | 27 |
| 65+ | 144 | 3.6 | 38 | ≤1 | ≤1 | 2 | 3 | 8 | 10 | 40 |
| **GRAND TOTAL** | **402** | **2.7** | **33** | **≤1** | **≤1** | **1** | **2** | **4** | **8** | **30** |

## 518: Miscellaneous operations on eyelid (08.0; 08.3-08.8; 08.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 123 | 1.5 | | ≤1 | ≤1 | 1 | 2 | 2 | 3 | 4 |
| 20-34 | 49 | 2.2 | 5 | ≤1 | ≤1 | 1 | 2 | 5 | 7 | 13 |
| 35-49 | 70 | 1.4 | | ≤1 | ≤1 | 1 | 2 | 2 | 3 | 4 |
| 50-64 | 161 | 1.6 | 1 | ≤1 | ≤1 | 1 | 2 | 2 | 3 | 6 |
| 65+ | 214 | 1.6 | | ≤1 | ≤1 | 1 | 2 | 2 | 3 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 154 | 2.3 | 5 | ≤1 | 1 | 1 | 3 | 5 | 7 | 11 |
| 20-34 | 184 | 3.4 | 26 | 1 | 1 | 2 | 4 | 7 | 9 | 27 |
| 35-49 | 121 | 3.7 | 32 | ≤1 | ≤1 | 2 | 4 | 7 | 11 | 37 |
| 50-64 | 169 | 3.1 | 29 | ≤1 | ≤1 | 2 | 3 | 5 | 10 | 35 |
| 65+ | 300 | 4.1 | 41 | ≤1 | 1 | 2 | 4 | 10 | 13 | 27 |
| **TOT. SINGLE DX** | 617 | 1.6 | 1 | ≤1 | ≤1 | 1 | 2 | 2 | 3 | 7 |
| **" MULTIPLE DX** | 928 | 3.4 | 29 | ≤1 | 1 | 2 | 3 | 7 | 11 | 26 |
| TOTAL 0-19 YRS | 277 | 2.0 | 3 | ≤1 | ≤1 | 1 | 2 | 4 | 6 | 10 |
| 20-34 | 233 | 3.2 | 22 | ≤1 | 1 | 2 | 4 | 6 | 9 | 26 |
| 35-49 | 191 | 2.9 | 22 | ≤1 | ≤1 | 2 | 3 | 6 | 8 | 28 |
| 50-64 | 330 | 2.4 | 16 | ≤1 | ≤1 | 1 | 2 | 4 | 6 | 19 |
| 65+ | 514 | 3.1 | 26 | ≤1 | ≤1 | 2 | 3 | 6 | 11 | 21 |
| **GRAND TOTAL** | **1545** | **2.7** | **19** | **≤1** | **≤1** | **2** | **2** | **5** | **9** | **20** |

## 519: Operations on lacrimal system (09.0-09.12; 09.2-09.3; 09.5-09.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 27 | 1.5 | | ≤1 | <1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 35 | 2.1 | | ≤1 | 1 | 2 | 2 | 3 | 4 | 4 |
| 35-49 | 36 | 1.9 | | ≤1 | 1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 58 | 2.2 | | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 65+ | 72 | 2.4 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 26 | 2.3 | 4 | ≤1 | ≤1 | 1 | 3 | 7 | 7 | 7 |
| 20-34 | 31 | 2.4 | 2 | ≤1 | ≤1 | 2 | 3 | 5 | 6 | 6 |
| 35-49 | 26 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 36 | 2.6 | 3 | 1 | 1 | 2 | 3 | 5 | 7 | 8 |
| 65+ | 78 | 3.4 | 66 | 1 | 1 | 2 | 3 | 3 | 5 | 56 |
| **TOT. SINGLE DX** | 228 | 2.1 | 1 | <1 | 1 | 2 | 3 | 3 | 4 | 5 |
| **" MULTIPLE DX** | 197 | 2.7 | 28 | <1 | 1 | 2 | 3 | 4 | 6 | 11 |
| TOTAL 0-19 YRS | 53 | 1.9 | 3 | ≤1 | <1 | 1 | 2 | 4 | 6 | 7 |
| 20-34 | 66 | 2.2 | 1 | ≤1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 35-49 | 62 | 1.9 | | 1 | 1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 94 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 65+ | 150 | 2.9 | 35 | 1 | 1 | 2 | 3 | 4 | 5 | 14 |
| **GRAND TOTAL** | **425** | **2.4** | **13** | **<1** | **1** | **2** | **3** | **3** | **5** | **7** |

## 520: Operations on conjunctiva (10.0-10.21; 10.3-10.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 11 | 1.4 | | ≤1 | ≤1 | 1 | 2 | 2 | 2 | 2 |
| 20-34 | 5 | 1.8 | 1 | ≤1 | ≤1 | 1 | 2 | 4 | 4 | 4 |
| 35-49 | 11 | 2.5 | 6 | ≤1 | ≤1 | 2 | 2 | 7 | 9 | 9 |
| 50-64 | 11 | 1.7 | | ≤1 | ≤1 | 2 | 2 | 3 | 4 | 4 |
| 65+ | 9 | 2.1 | 3 | ≤1 | ≤1 | <1 | 3 | 6 | 6 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 2.1 | 4 | ≤1 | ≤1 | 1 | 2 | 4 | 8 | 9 |
| 20-34 | 7 | 3.0 | 3 | ≤1 | ≤1 | 3 | 4 | 6 | 6 | 6 |
| 35-49 | 6 | 5.2 | 41 | 1 | 1 | 3 | 5 | 17 | 18 | 18 |
| 50-64 | 9 | 4.7 | 39 | 1 | 1 | 2 | 4 | 16 | 20 | 20 |
| 65+ | 20 | 4.6 | 51 | ≤1 | 1 | 2 | 5 | 8 | 21 | 33 |
| **TOT. SINGLE DX** | 47 | 1.9 | 3 | ≤1 | ≤1 | 1 | 2 | 4 | 5 | 9 |
| **" MULTIPLE DX** | 56 | 3.8 | 30 | ≤1 | 1 | 2 | 4 | 9 | 15 | 32 |
| TOTAL 0-19 YRS | 25 | 1.8 | 3 | ≤1 | ≤1 | 1 | 2 | 2 | 5 | 9 |
| 20-34 | 12 | 2.5 | 3 | ≤1 | ≤1 | 2 | 4 | 5 | 6 | 6 |
| 35-49 | 17 | 3.4 | 19 | ≤1 | ≤1 | 2 | 3 | 8 | 15 | 18 |
| 50-64 | 20 | 3.1 | 19 | ≤1 | ≤1 | 2 | 2 | 6 | 14 | 20 |
| 65+ | 29 | 3.8 | 37 | ≤1 | ≤1 | 2 | 5 | 8 | 10 | 33 |
| **GRAND TOTAL** | **103** | **2.9** | **18** | **≤1** | **≤1** | **2** | **3** | **6** | **9** | **26** |

## 521: Corneal transplant (11.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 27 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 20-34 | 77 | 3.1 | 1 | 1 | 2 | 3 | 4 | 5 | 5 | 8 |
| 35-49 | 62 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 9 |
| 50-64 | 45 | 3.0 | 3 | 1 | 1 | 3 | 4 | 5 | 5 | 12 |
| 65+ | 151 | 3.3 | 4 | 1 | 1 | 3 | 4 | 5 | 7 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 4.2 | 7 | 1 | 2 | 4 | 5 | 7 | 10 | 13 |
| 20-34 | 41 | 3.9 | 4 | 1 | 2 | 3 | 5 | 7 | 8 | 10 |
| 35-49 | 38 | 4.4 | 20 | 1 | 1 | 3 | 4 | 9 | 14 | 25 |
| 50-64 | 105 | 4.4 | 16 | 1 | 2 | 3 | 5 | 8 | 11 | 24 |
| 65+ | 289 | 3.9 | 5 | 1 | 2 | 4 | 5 | 6 | 8 | 15 |
| **TOT. SINGLE DX** | 362 | 3.2 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 492 | 4.1 | 9 | 1 | 2 | 3 | 5 | 7 | 9 | 16 |
| TOTAL 0-19 YRS | 46 | 3.5 | 5 | 1 | 1 | 3 | 4 | 6 | 7 | 13 |
| 20-34 | 118 | 3.4 | 3 | 1 | 2 | 3 | 4 | 5 | 7 | 10 |
| 35-49 | 100 | 3.6 | 9 | 1 | 1 | 3 | 4 | 5 | 8 | 20 |
| 50-64 | 150 | 4.0 | 13 | 1 | 1 | 3 | 4 | 7 | 11 | 22 |
| 65+ | 440 | 3.7 | 5 | 1 | 2 | 3 | 5 | 6 | 8 | 15 |
| **GRAND TOTAL** | **854** | **3.7** | **7** | **1** | **2** | **3** | **4** | **6** | **8** | **15** |

## 522: Miscellaneous operations on cornea (11.0-11.22; 11.3-11.5; 11.7-11.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 27 | 3.1 | 11 | <1 | <1 | 2 | 4 | 7 | 9 | 17 |
| 20-34 | 60 | 1.7 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 35-49 | 55 | 2.1 | 2 | <1 | <1 | 1 | 3 | 4 | 5 | 8 |
| 50-64 | 68 | 1.6 | | <1 | <1 | 1 | 2 | 3 | 4 | 4 |
| 65+ | 48 | 1.9 | 2 | <1 | <1 | 2 | 2 | 3 | 5 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 87 | 3.6 | 10 | 1 | 1 | 3 | 5 | 7 | 8 | 20 |
| 20-34 | 59 | 3.8 | 10 | <1 | 1 | 3 | 5 | 8 | 9 | 18 |
| 35-49 | 41 | 3.4 | 7 | <1 | 1 | 3 | 5 | 6 | 8 | 15 |
| 50-64 | 48 | 6.3 | 140 | 1 | 1 | 2 | 5 | 14 | 29 | 73 |
| 65+ | 72 | 4.0 | 16 | <1 | 1 | 2 | 5 | 9 | 14 | 19 |
| **TOT. SINGLE DX** | 258 | 1.9 | 3 | <1 | <1 | 1 | 2 | 4 | 5 | 8 |
| **" MULTIPLE DX** | 307 | 4.1 | 32 | <1 | 1 | 3 | 5 | 8 | 12 | 27 |
| TOTAL 0-19 YRS | 114 | 3.5 | 11 | <1 | 1 | 3 | 4 | 7 | 8 | 20 |
| 20-34 | 119 | 2.7 | 7 | <1 | <1 | 2 | 3 | 5 | 8 | 15 |
| 35-49 | 96 | 2.6 | 5 | <1 | <1 | 2 | 4 | 5 | 7 | 13 |
| 50-64 | 116 | 3.5 | 63 | <1 | <1 | 2 | 3 | 5 | 12 | 47 |
| 65+ | 120 | 3.2 | 12 | <1 | <1 | 2 | 3 | 8 | 11 | 17 |
| **GRAND TOTAL** | **565** | **3.1** | **20** | **<1** | **<1** | **2** | **4** | **6** | **8** | **19** |

## 523: Operations for regulation of intraocular pressure (12.5-12.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 74 | 2.0 | 1 | <1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 20-34 | 22 | 2.9 | 2 | 1 | 1 | 3 | 3 | 5 | 7 | 8 |
| 35-49 | 41 | 2.8 | 2 | 1 | 1 | 2 | 3 | 5 | 6 | 10 |
| 50-64 | 119 | 3.1 | 9 | 1 | 1 | 2 | 3 | 6 | 8 | 19 |
| 65+ | 233 | 2.8 | 5 | 1 | 1 | 2 | 3 | 4 | 7 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 38 | 3.9 | 58 | <1 | 1 | 2 | 3 | 6 | 13 | 47 |
| 20-34 | 23 | 3.6 | 7 | <1 | 1 | 3 | 5 | 7 | 11 | 12 |
| 35-49 | 23 | 4.2 | 16 | <1 | 1 | 3 | 4 | 11 | 15 | 16 |
| 50-64 | 91 | 3.9 | 6 | 1 | 2 | 3 | 5 | 8 | 9 | 12 |
| 65+ | 217 | 4.1 | 17 | 1 | 1 | 3 | 5 | 9 | 12 | 23 |
| **TOT. SINGLE DX** | 489 | 2.8 | 6 | 1 | 1 | 2 | 3 | 5 | 6 | 13 |
| **" MULTIPLE DX** | 392 | 4.0 | 18 | 1 | 1 | 3 | 5 | 9 | 10 | 19 |
| TOTAL 0-19 YRS | 112 | 2.7 | 21 | <1 | 1 | 2 | 3 | 4 | 5 | 27 |
| 20-34 | 45 | 3.2 | 5 | 1 | 1 | 3 | 4 | 6 | 8 | 12 |
| 35-49 | 64 | 3.3 | 8 | 1 | 1 | 2 | 4 | 5 | 10 | 16 |
| 50-64 | 210 | 3.4 | 9 | 1 | 1 | 2 | 4 | 7 | 9 | 13 |
| 65+ | 450 | 3.4 | 12 | 1 | 1 | 2 | 4 | 7 | 9 | 18 |
| **GRAND TOTAL** | **881** | **3.3** | **12** | **1** | **1** | **2** | **4** | **6** | **9** | **16** |

## 524: Other operations on iris, ciliary body, sclera and anterior chamber (12.00-12.22; 12.3-12.4; 12.8-12.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 22 | 2.9 | 4 | 1 | 1 | 2 | 4 | 5 | 7 | 9 |
| 20-34 | 27 | 2.7 | 3 | <1 | 1 | 2 | 3 | 5 | 6 | 8 |
| 35-49 | 35 | 2.3 | 2 | <1 | 1 | 2 | 3 | 5 | 5 | 6 |
| 50-64 | 77 | 2.4 | 3 | <1 | <1 | 2 | 3 | 5 | 6 | 10 |
| 65+ | 139 | 2.5 | 5 | <1 | 1 | 2 | 3 | 5 | 6 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 40 | 4.5 | 9 | <1 | 1 | 4 | 6 | 9 | 11 | 11 |
| 20-34 | 42 | 4.1 | 12 | <1 | 1 | 3 | 5 | 9 | 12 | 15 |
| 35-49 | 26 | 3.7 | 7 | 1 | 1 | 3 | 5 | 7 | 10 | 12 |
| 50-64 | 72 | 3.4 | 8 | 1 | 1 | 3 | 4 | 6 | 12 | 15 |
| 65+ | 161 | 3.1 | 6 | 1 | 1 | 2 | 4 | 6 | 9 | 12 |
| **TOT. SINGLE DX** | 300 | 2.5 | 4 | <1 | 1 | 2 | 3 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 341 | 3.5 | 8 | 1 | 1 | 2 | 4 | 8 | 10 | 15 |
| TOTAL 0-19 YRS | 62 | 3.9 | 8 | 1 | 1 | 3 | 5 | 8 | 10 | 11 |
| 20-34 | 69 | 3.6 | 9 | 1 | 1 | 3 | 5 | 8 | 10 | 15 |
| 35-49 | 61 | 2.9 | 5 | 1 | 1 | 2 | 4 | 6 | 7 | 12 |
| 50-64 | 149 | 2.9 | 6 | <1 | 1 | 2 | 3 | 5 | 8 | 15 |
| 65+ | 300 | 2.8 | 6 | <1 | 1 | 2 | 3 | 5 | 8 | 12 |
| **GRAND TOTAL** | **641** | **3.0** | **7** | **<1** | **1** | **2** | **4** | **6** | **8** | **13** |

## 525: Intracapsular extraction of lens (13.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 17 | 1.9 | | <1 | 1 | 2 | 2 | 2 | 4 | 5 |
| 20-34 | 9 | 2.2 | | 2 | 2 | 2 | 3 | 3 | 3 | 3 |
| 35-49 | 196 | 2.3 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 8 |
| 50-64 | 1201 | 2.2 | | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 65+ | 4458 | 2.4 | 2 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 15 | 3.1 | 5 | 1 | 1 | 2 | 4 | 6 | 9 | 10 |
| 20-34 | 22 | 4.3 | 29 | <1 | 1 | 3 | 4 | 8 | 16 | 27 |
| 35-49 | 123 | 3.0 | 4 | 1 | 1 | 2 | 3 | 5 | 8 | 11 |
| 50-64 | 892 | 2.8 | 6 | 1 | 2 | 2 | 3 | 4 | 6 | 12 |
| 65+ | 4170 | 2.9 | 10 | 1 | 2 | 2 | 3 | 4 | 6 | 15 |
| **TOT. SINGLE DX** | 5881 | 2.3 | 2 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| **" MULTIPLE DX** | 5222 | 2.9 | 10 | 1 | 2 | 2 | 3 | 4 | 6 | 14 |
| TOTAL 0-19 YRS | 32 | 2.5 | 3 | 1 | 1 | 2 | 2 | 5 | 6 | 10 |
| 20-34 | 31 | 3.7 | 22 | 1 | 1 | 3 | 4 | 6 | 9 | 27 |
| 35-49 | 319 | 2.5 | 3 | 1 | 1 | 2 | 3 | 4 | 5 | 10 |
| 50-64 | 2093 | 2.5 | 3 | 1 | 1 | 2 | 3 | 3 | 4 | 9 |
| 65+ | 8628 | 2.6 | 6 | 1 | 1 | 2 | 3 | 4 | 5 | 11 |
| **GRAND TOTAL** | **11103** | **2.6** | **6** | **1** | **1** | **2** | **3** | **4** | **5** | **10** |

## 526: Other lens extraction (13.2-13.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 97 | 2.0 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 9 |
| 20-34 | 49 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 3 | 4 |
| 35-49 | 311 | 1.8 | | 1 | 1 | 2 | 2 | 2 | 3 | 4 |
| 50-64 | 1648 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 65+ | 6460 | 2.1 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 49 | 2.3 | 2 | <1 | 1 | 2 | 3 | 4 | 7 | 8 |
| 20-34 | 62 | 2.5 | 2 | <1 | 1 | 2 | 3 | 4 | 6 | 8 |
| 35-49 | 197 | 2.6 | 22 | 1 | 1 | 2 | 2 | 3 | 5 | 12 |
| 50-64 | 1091 | 2.3 | 2 | 1 | 1 | 2 | 2 | 3 | 4 | 8 |
| 65+ | 5186 | 2.5 | 4 | 1 | 1 | 2 | 3 | 3 | 5 | 11 |
| **TOT. SINGLE DX** | 8565 | 2.0 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| **" MULTIPLE DX** | 6585 | 2.5 | 5 | 1 | 1 | 2 | 2 | 3 | 5 | 10 |
| TOTAL 0-19 YRS | 146 | 2.1 | 2 | <1 | 1 | 2 | 2 | 3 | 4 | 8 |
| 20-34 | 111 | 2.2 | 2 | <1 | 1 | 2 | 2 | 3 | 5 | 8 |
| 35-49 | 508 | 2.1 | 9 | 1 | 1 | 2 | 2 | 3 | 3 | 7 |
| 50-64 | 2739 | 2.1 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 7 |
| 65+ | 11646 | 2.3 | 3 | 1 | 1 | 2 | 2 | 3 | 4 | 8 |
| **GRAND TOTAL** | **15150** | **2.2** | **3** | **1** | **1** | **2** | **2** | **3** | **4** | **8** |

## 527: Miscellaneous operations on lens (13.0; 13.7-13.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 1.3 | | <1 | <1 | 1 | 2 | 2 | 2 | 2 |
| 20-34 | 2 | 2.5 | 1 | 2 | 2 | 2 | 3 | 3 | 3 | 3 |
| 35-49 | 32 | 2.0 | | <1 | 1 | 2 | 2 | 3 | 3 | 6 |
| 50-64 | 230 | 2.0 | 3 | 1 | 1 | 2 | 2 | 2 | 3 | 7 |
| 65+ | 792 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 1.5 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 8 | 2.3 | 1 | <1 | <1 | 2 | 3 | 4 | 4 | 4 |
| 35-49 | 22 | 2.5 | 5 | <1 | 1 | 2 | 3 | 4 | 7 | 12 |
| 50-64 | 148 | 2.2 | 2 | <1 | 1 | 2 | 2 | 3 | 5 | 9 |
| 65+ | 635 | 2.5 | 5 | 1 | 1 | 2 | 3 | 3 | 5 | 13 |
| **TOT. SINGLE DX** | 1060 | 2.0 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 6 |
| **" MULTIPLE DX** | 817 | 2.5 | 5 | 1 | 1 | 2 | 3 | 3 | 5 | 12 |
| TOTAL 0-19 YRS | 8 | 1.4 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 10 | 2.3 | 1 | <1 | <1 | 2 | 3 | 4 | 4 | 4 |
| 35-49 | 54 | 2.2 | 3 | <1 | 1 | 2 | 2 | 3 | 4 | 12 |
| 50-64 | 378 | 2.1 | 3 | <1 | 1 | 2 | 2 | 3 | 4 | 8 |
| 65+ | 1427 | 2.2 | 3 | 1 | 1 | 2 | 2 | 3 | 4 | 8 |
| **GRAND TOTAL** | **1877** | **2.2** | **3** | **1** | **1** | **2** | **2** | **3** | **4** | **8** |

## 528: Repair of retina (14.3-14.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 48 | 4.1 | 5 | 1 | 1 | 4 | 5 | 8 | 9 | 11 |
| 20-34 | 133 | 4.5 | 4 | 2 | 2 | 4 | 5 | 7 | 9 | 12 |
| 35-49 | 129 | 4.4 | 5 | 1 | 2 | 4 | 6 | 8 | 9 | 13 |
| 50-64 | 455 | 4.5 | 7 | 1 | 2 | 4 | 5 | 7 | 9 | 14 |
| 65+ | 431 | 4.5 | 6 | 2 | 2 | 4 | 5 | 7 | 9 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 49 | 5.2 | 14 | 2 | 2 | 4 | 6 | 8 | 11 | 22 |
| 20-34 | 101 | 4.7 | 7 | 1 | 2 | 4 | 6 | 8 | 10 | 15 |
| 35-49 | 100 | 5.1 | 8 | 1 | 2 | 4 | 6 | 8 | 12 | 16 |
| 50-64 | 385 | 5.3 | 11 | 2 | 2 | 4 | 6 | 10 | 12 | 17 |
| 65+ | 483 | 5.3 | 14 | 2 | 2 | 4 | 7 | 10 | 12 | 23 |
| **TOT. SINGLE DX** | 1196 | 4.5 | 7 | 1 | 2 | 4 | 5 | 7 | 9 | 14 |
| **" MULTIPLE DX** | 1118 | 5.2 | 12 | 2 | 2 | 4 | 6 | 9 | 12 | 18 |
| TOTAL 0-19 YRS | 97 | 4.7 | 10 | 1 | 2 | 4 | 5 | 8 | 9 | 22 |
| 20-34 | 234 | 4.6 | 6 | 2 | 2 | 4 | 6 | 8 | 9 | 12 |
| 35-49 | 229 | 4.7 | 7 | 1 | 2 | 4 | 6 | 8 | 9 | 15 |
| 50-64 | 840 | 4.8 | 10 | 1 | 2 | 4 | 6 | 9 | 10 | 17 |
| 65+ | 914 | 5.0 | 11 | 2 | 2 | 4 | 6 | 8 | 11 | 18 |
| **GRAND TOTAL** | **2314** | **4.8** | **10** | **2** | **2** | **4** | **6** | **8** | **10** | **17** |

## 529: Miscellaneous operations on retina, choroid, and vitreous (14.0-14.11; 14.2; 14.6-14.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 23 | 2.7 | 5 | <1 | 1 | 2 | 3 | 4 | 7 | 12 |
| 20-34 | 28 | 3.4 | 3 | 1 | 1 | 4 | 5 | 5 | 6 | 8 |
| 35-49 | 22 | 3.4 | 2 | 1 | 1 | 3 | 4 | 6 | 7 | 7 |
| 50-64 | 58 | 3.8 | 8 | 1 | 1 | 3 | 5 | 8 | 10 | 15 |
| 65+ | 123 | 3.4 | 4 | 1 | 1 | 3 | 4 | 7 | 8 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 36 | 3.9 | 7 | <1 | 1 | 3 | 5 | 8 | 10 | 12 |
| 20-34 | 111 | 4.6 | 24 | 1 | 2 | 4 | 5 | 7 | 9 | 31 |
| 35-49 | 100 | 3.7 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 11 |
| 50-64 | 200 | 4.2 | 21 | 1 | 1 | 3 | 5 | 7 | 10 | 20 |
| 65+ | 316 | 4.1 | 13 | 1 | 1 | 3 | 5 | 7 | 9 | 19 |
| **TOT. SINGLE DX** | 254 | 3.4 | 5 | 1 | 1 | 3 | 4 | 6 | 8 | 13 |
| **" MULTIPLE DX** | 763 | 4.1 | 16 | 1 | 1 | 3 | 5 | 7 | 9 | 18 |
| TOTAL 0-19 YRS | 59 | 3.4 | 7 | <1 | 1 | 3 | 4 | 7 | 10 | 12 |
| 20-34 | 139 | 4.3 | 20 | 1 | 2 | 4 | 5 | 7 | 8 | 23 |
| 35-49 | 122 | 3.7 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 11 |
| 50-64 | 258 | 4.1 | 19 | 1 | 1 | 3 | 5 | 7 | 10 | 15 |
| 65+ | 439 | 3.9 | 11 | 1 | 1 | 3 | 5 | 7 | 9 | 17 |
| **GRAND TOTAL** | **1017** | **4.0** | **13** | **1** | **1** | **3** | **5** | **7** | **9** | **16** |

## 530: Resection and recession of ocular muscle (15.1-15.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 893 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 2 | 5 |
| 20-34 | 180 | 1.7 | | <1 | 1 | 2 | 2 | 2 | 3 | 4 |
| 35-49 | 86 | 1.6 | | <1 | <1 | 2 | 2 | 2 | 2 | 3 |
| 50-64 | 45 | 1.8 | | <1 | 1 | 2 | 2 | 2 | 3 | 5 |
| 65+ | 37 | 1.9 | | <1 | <1 | 2 | 2 | 3 | 3 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 206 | 1.7 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| 20-34 | 68 | 2.0 | 2 | <1 | 1 | 2 | 2 | 3 | 4 | 11 |
| 35-49 | 62 | 2.0 | 3 | <1 | 1 | 2 | 2 | 2 | 3 | 14 |
| 50-64 | 45 | 2.3 | 3 | 1 | 1 | 2 | 2 | 3 | 5 | 13 |
| 65+ | 45 | 2.2 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 8 |
| **TOT. SINGLE DX** | 1241 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 4 |
| **" MULTIPLE DX** | 426 | 1.9 | 2 | <1 | <1 | 2 | 2 | 3 | 3 | 8 |
| TOTAL 0-19 YRS | 1099 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 7 |
| 20-34 | 248 | 1.8 | 1 | <1 | 1 | 2 | 2 | 2 | 3 | 5 |
| 35-49 | 148 | 1.8 | 2 | <1 | <1 | 2 | 2 | 2 | 3 | 4 |
| 50-64 | 90 | 2.1 | 2 | 1 | 1 | 2 | 2 | 3 | 4 | 11 |
| 65+ | 82 | 2.0 | 1 | <1 | 1 | 2 | 2 | 3 | 3 | 7 |
| **GRAND TOTAL** | **1667** | **1.6** | **1** | **<1** | **<1** | **2** | **2** | **2** | **3** | **7** |

## 531: Miscellaneous operations on ocular muscle (15.01; 15.7-15.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 1.8 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 3 |
| 20-34 | 3 | 1.3 | | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 35-49 | 2 | 1.5 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 50-64 | 1 | 1.0 | | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 1.3 | | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 20-34 | 5 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| 35-49 | 2 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 50-64 | 3 | 15.0 | 547 | 1 | 1 | 2 | 32 | 42 | 42 | 42 |
| 65+ | 1 | 1.0 | | <1 | <1 | <1 | <1 | <1 | <1 | <1 |
| **TOT. SINGLE DX** | 10 | 1.5 | 1 | 1 | 1 | 1 | 2 | 3 | 3 | 3 |
| **" MULTIPLE DX** | 14 | 4.4 | 118 | <1 | <1 | 1 | 2 | 7 | 34 | 42 |
| TOTAL 0-19 YRS | 7 | 1.6 | 1 | 1 | 1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 8 | 1.3 | | <1 | <1 | 1 | 2 | 2 | 2 | 2 |
| 35-49 | 4 | 2.3 | 1 | 1 | 1 | 2 | 3 | 3 | 3 | 3 |
| 50-64 | 4 | 11.5 | 414 | 1 | 1 | 2 | 22 | 42 | 42 | 42 |
| 65+ | 1 | 1.0 | | <1 | <1 | <1 | <1 | <1 | <1 | <1 |
| **GRAND TOTAL** | **24** | **3.2** | **69** | **<1** | **<1** | **1** | **2** | **3** | **15** | **42** |

## 532: Operations on orbit and eyeball (16.0-16.1; 16.22-16.23; 16.3-16.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 63 | 2.6 | 5 | <1 | 1 | 2 | 3 | 5 | 6 | 14 |
| 20-34 | 61 | 2.7 | 3 | 1 | 1 | 2 | 3 | 5 | 8 | 9 |
| 35-49 | 42 | 2.9 | 5 | 1 | 1 | 3 | 3 | 4 | 4 | 17 |
| 50-64 | 55 | 2.4 | 1 | <1 | 1 | 2 | 3 | 4 | 4 | 10 |
| 65+ | 81 | 2.9 | 3 | 1 | 1 | 2 | 4 | 5 | 5 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 59 | 4.3 | 23 | 1 | 1 | 3 | 5 | 9 | 11 | 32 |
| 20-34 | 121 | 4.2 | 21 | 1 | 1 | 3 | 5 | 9 | 12 | 30 |
| 35-49 | 48 | 4.5 | 35 | 1 | 1 | 3 | 5 | 8 | 10 | 41 |
| 50-64 | 84 | 5.0 | 45 | 1 | 2 | 3 | 5 | 10 | 15 | 44 |
| 65+ | 115 | 4.5 | 13 | 1 | 1 | 3 | 6 | 9 | 14 | 48 |
| **TOT. SINGLE DX** | 302 | 2.7 | 4 | 1 | 1 | 2 | 3 | 4 | 5 | 11 |
| **" MULTIPLE DX** | 427 | 4.5 | 26 | 1 | 1 | 3 | 5 | 9 | 12 | 34 |
| TOTAL 0-19 YRS | 122 | 3.4 | 15 | <1 | 1 | 2 | 4 | 6 | 10 | 20 |
| 20-34 | 182 | 3.7 | 16 | 1 | 1 | 2 | 4 | 8 | 10 | 24 |
| 35-49 | 90 | 3.7 | 22 | 1 | 1 | 3 | 4 | 6 | 9 | 31 |
| 50-64 | 139 | 4.0 | 30 | 1 | 1 | 3 | 4 | 7 | 10 | 33 |
| 65+ | 196 | 3.8 | 10 | 1 | 1 | 3 | 4 | 8 | 11 | 19 |
| **GRAND TOTAL** | **729** | **3.7** | **18** | **1** | **1** | **3** | **4** | **7** | **10** | **24** |

## 533: Misc. diagnostic and therapeutic procedures on eye (08.19; 08.91-08.93; 09.19; 09.4; 10.29; 11.29; 12.29; 14.19; 15.09; 16.21; 16.29)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 534: Operations on external ear (18.02-18.09; 18.12; 18.2-18.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 262 | 1.6 | | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| 20-34 | 83 | 2.0 | 4 | <1 | <1 | 1 | 2 | 4 | 7 | 11 |
| 35-49 | 51 | 1.8 | 2 | <1 | <1 | 1 | 2 | 3 | 5 | 9 |
| 50-64 | 61 | 2.0 | 2 | <1 | <1 | 2 | 2 | 4 | 5 | 10 |
| 65+ | 79 | 2.8 | 11 | <1 | <1 | 2 | 3 | 5 | 10 | 20 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 169 | 2.8 | 15 | <1 | 1 | 2 | 3 | 5 | 7 | 18 |
| 20-34 | 76 | 4.9 | 99 | <1 | 1 | 2 | 5 | 10 | 13 | 67 |
| 35-49 | 47 | 4.9 | 31 | 1 | 1 | 2 | 7 | 13 | 18 | 23 |
| 50-64 | 66 | 3.9 | 19 | <1 | 1 | 2 | 4 | 11 | 13 | 21 |
| 65+ | 95 | 9.7 | 200 | 1 | 1 | 5 | 10 | 24 | 40 | 74 |
| **TOT. SINGLE DX** | 536 | 1.9 | 3 | <1 | <1 | 1 | 2 | 3 | 5 | 10 |
| **" MULTIPLE DX** | 453 | 5.0 | 76 | <1 | 1 | 2 | 5 | 11 | 17 | 47 |
| TOTAL 0-19 YRS | 431 | 2.1 | 7 | <1 | <1 | 1 | 2 | 4 | 5 | 8 |
| 20-34 | 159 | 3.3 | 51 | <1 | <1 | 2 | 3 | 7 | 11 | 15 |
| 35-49 | 98 | 3.3 | 19 | <1 | <1 | 2 | 3 | 9 | 13 | 23 |
| 50-64 | 127 | 3.0 | 12 | <1 | <1 | 2 | 3 | 7 | 11 | 19 |
| 65+ | 174 | 6.5 | 126 | <1 | <1 | 3 | 6 | 16 | 25 | 65 |
| **GRAND TOTAL** | **989** | **3.3** | **39** | **<1** | **<1** | **2** | **3** | **7** | **11** | **27** |

## 535: Stapes mobilization and stapedectomy (19.0-19.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 46 | 1.9 | | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 20-34 | 208 | 1.7 | | 1 | 1 | 1 | 2 | 3 | 3 | 4 |
| 35-49 | 289 | 1.7 | | 1 | 1 | 1 | 2 | 3 | 3 | 5 |
| 50-64 | 346 | 1.6 | | 1 | 1 | 1 | 2 | 3 | 3 | 4 |
| 65+ | 159 | 1.8 | | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 22 | 2.0 | | <1 | 1 | 2 | 3 | 3 | 4 | 4 |
| 20-34 | 64 | 1.8 | | 1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 35-49 | 86 | 2.2 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 50-64 | 93 | 2.6 | 13 | 1 | 1 | 2 | 2 | 4 | 6 | 25 |
| 65+ | 49 | 2.4 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 7 |
| **TOT. SINGLE DX** | 1048 | 1.7 | 1 | 1 | 1 | 1 | 2 | 3 | 3 | 4 |
| **" MULTIPLE DX** | 314 | 2.2 | 5 | 1 | 1 | 2 | 3 | 3 | 4 | 8 |
| TOTAL 0-19 YRS | 68 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 20-34 | 272 | 1.7 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 35-49 | 375 | 1.8 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 50-64 | 439 | 1.8 | 4 | 1 | 1 | 1 | 2 | 3 | 3 | 7 |
| 65+ | 208 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 6 |
| **GRAND TOTAL** | **1362** | **1.8** | **2** | **1** | **1** | **2** | **2** | **3** | **3** | **5** |

## 536: Tympanoplasty (19.4-19.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 566 | 1.7 | | <1 | 1 | 2 | 2 | 2 | 3 | 3 |
| 20-34 | 431 | 1.7 | | <1 | 1 | 2 | 2 | 2 | 3 | 5 |
| 35-49 | 303 | 1.7 | | <1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 50-64 | 270 | 1.7 | | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 65+ | 113 | 1.7 | | <1 | 1 | 2 | 2 | 3 | 3 | 4 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 349 | 1.9 | 2 | <1 | 1 | 2 | 2 | 3 | 3 | 6 |
| 20-34 | 263 | 2.2 | 26 | <1 | 1 | 2 | 2 | 3 | 4 | 7 |
| 35-49 | 215 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 7 |
| 50-64 | 202 | 2.2 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 65+ | 97 | 2.6 | 6 | 1 | 1 | 2 | 3 | 4 | 7 | 16 |
| **TOT. SINGLE DX** | 1683 | 1.7 | 1 | <1 | 1 | 2 | 2 | 2 | 3 | 4 |
| **" MULTIPLE DX** | 1126 | 2.1 | 8 | <1 | 1 | 2 | 2 | 3 | 4 | 7 |
| TOTAL 0-19 YRS | 915 | 1.8 | 1 | <1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 20-34 | 694 | 1.9 | 10 | <1 | 1 | 2 | 2 | 3 | 3 | 6 |
| 35-49 | 518 | 1.8 | 1 | <1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 50-64 | 472 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 65+ | 210 | 2.1 | 4 | <1 | 1 | 2 | 2 | 3 | 4 | 9 |
| **GRAND TOTAL** | **2809** | **1.9** | **4** | **<1** | **1** | **2** | **2** | **3** | **3** | **5** |

## 537: Myringotomy (20.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3208 | 1.1 | | <1 | <1 | <1 | <1 | 1 | 1 | 2 |
| 20-34 | 56 | 1.6 | 3 | <1 | <1 | 1 | 2 | 2 | 4 | 13 |
| 35-49 | 46 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 4 | 7 |
| 50-64 | 35 | 1.7 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 7 |
| 65+ | 19 | 1.1 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1532 | 1.5 | 3 | <1 | <1 | 1 | 1 | 2 | 4 | 10 |
| 20-34 | 54 | 4.1 | 40 | <1 | <1 | 2 | 4 | 13 | 18 | 36 |
| 35-49 | 39 | 3.3 | 41 | <1 | <1 | 1 | 2 | 5 | 17 | 34 |
| 50-64 | 47 | 3.7 | 14 | <1 | 1 | 2 | 4 | 9 | 13 | 18 |
| 65+ | 31 | 6.3 | 52 | <1 | 1 | 2 | 9 | 17 | 26 | 28 |
| **TOT. SINGLE DX** | 3364 | 1.1 | | <1 | <1 | <1 | <1 | 1 | 1 | 3 |
| **" MULTIPLE DX** | 1703 | 1.8 | 7 | <1 | <1 | 1 | 1 | 3 | 5 | 14 |
| TOTAL 0-19 YRS | 4740 | 1.2 | 1 | <1 | <1 | <1 | 1 | 1 | 2 | 6 |
| 20-34 | 110 | 2.8 | 23 | <1 | <1 | 1 | 2 | 5 | 13 | 26 |
| 35-49 | 85 | 2.4 | 20 | <1 | <1 | 1 | 2 | 4 | 6 | 31 |
| 50-64 | 82 | 2.9 | 10 | <1 | <1 | 2 | 3 | 7 | 9 | 16 |
| 65+ | 50 | 4.3 | 39 | <1 | <1 | 2 | 4 | 12 | 18 | 28 |
| **GRAND TOTAL** | **5067** | **1.3** | **3** | **<1** | **<1** | **<1** | **1** | **2** | **2** | **8** |

## 538: Mastoidectomy (20.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 98 | 2.2 | 3 | 1 | 1 | 2 | 2 | 4 | 6 | 10 |
| 20-34 | 102 | 2.0 | 1 | 1 | 1 | 2 | 2 | 3 | 5 | 8 |
| 35-49 | 77 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 7 |
| 50-64 | 67 | 2.5 | 1 | 1 | 1 | 2 | 3 | 3 | 6 | 8 |
| 65+ | 27 | 3.0 | 4 | 1 | 1 | 2 | 3 | 6 | 9 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 185 | 2.5 | 3 | 1 | 1 | 2 | 3 | 4 | 5 | 12 |
| 20-34 | 158 | 2.6 | 6 | 1 | 1 | 2 | 3 | 4 | 6 | 12 |
| 35-49 | 127 | 2.5 | 10 | 1 | 1 | 2 | 3 | 4 | 6 | 45 |
| 50-64 | 148 | 3.7 | 33 | 1 | 1 | 2 | 3 | 6 | 12 | 37 |
| 65+ | 65 | 3.8 | 7 | 1 | 2 | 3 | 4 | 7 | 8 | 17 |
| **TOT. SINGLE DX** | 371 | 2.3 | 2 | 1 | 1 | 2 | 3 | 4 | 6 | 8 |
| **" MULTIPLE DX** | 683 | 2.9 | 13 | 1 | 1 | 2 | 3 | 5 | 7 | 20 |
| TOTAL 0-19 YRS | 283 | 2.4 | 3 | 1 | 1 | 2 | 3 | 4 | 6 | 11 |
| 20-34 | 260 | 2.3 | 5 | 1 | 1 | 2 | 3 | 4 | 6 | 11 |
| 35-49 | 204 | 2.4 | 7 | 1 | 1 | 2 | 3 | 4 | 5 | 25 |
| 50-64 | 215 | 3.3 | 24 | 1 | 1 | 2 | 3 | 5 | 8 | 27 |
| 65+ | 92 | 3.5 | 7 | 1 | 1 | 3 | 4 | 6 | 9 | 15 |
| **GRAND TOTAL** | **1054** | **2.7** | **9** | **1** | **1** | **2** | **3** | **4** | **6** | **15** |

## 539: Miscellaneous operations on middle ear and inner ear (19.6-19.9; 20.1-20.32; 20.5-20.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 253 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 6 |
| 20-34 | 198 | 2.2 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 10 |
| 35-49 | 183 | 2.3 | 4 | <1 | 1 | 2 | 3 | 4 | 5 | 11 |
| 50-64 | 254 | 2.6 | 5 | 1 | 1 | 2 | 3 | 4 | 6 | 9 |
| 65+ | 82 | 2.7 | 4 | 1 | 1 | 2 | 3 | 5 | 7 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 155 | 2.3 | 5 | <1 | <1 | 2 | 2 | 4 | 8 | 14 |
| 20-34 | 84 | 3.2 | 10 | <1 | 1 | 2 | 3 | 7 | 8 | 18 |
| 35-49 | 104 | 3.5 | 11 | 1 | 1 | 3 | 4 | 8 | 11 | 18 |
| 50-64 | 94 | 3.9 | 17 | 1 | 1 | 3 | 4 | 7 | 12 | 25 |
| 65+ | 41 | 6.6 | 122 | 1 | 1 | 3 | 6 | 14 | 16 | 70 |
| **TOT. SINGLE DX** | 970 | 2.2 | 4 | <1 | 1 | 2 | 2 | 4 | 5 | 9 |
| **" MULTIPLE DX** | 478 | 3.4 | 21 | <1 | <1 | 2 | 3 | 7 | 11 | 18 |
| TOTAL 0-19 YRS | 408 | 1.9 | 3 | <1 | <1 | 1 | 2 | 3 | 4 | 11 |
| 20-34 | 282 | 2.5 | 5 | 1 | 1 | 2 | 3 | 4 | 7 | 13 |
| 35-49 | 287 | 2.8 | 7 | 1 | 1 | 2 | 3 | 5 | 8 | 17 |
| 50-64 | 348 | 2.9 | 9 | 1 | 1 | 2 | 3 | 5 | 7 | 15 |
| 65+ | 123 | 4.0 | 46 | 1 | 1 | 2 | 4 | 7 | 13 | 31 |
| **GRAND TOTAL** | **1448** | **2.6** | **10** | **<1** | **<1** | **2** | **3** | **5** | **7** | **14** |

## 540: Other diagnostic and therapeutic procedures on ear (18.01; 18.11; 18.19; 20.39)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 541: Submucous resection of nasal septum (21.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 63 | 1.6 | | ≤1 | ≤1 | 1 | 2 | 3 | 3 | 5 |
| 20-34 | 204 | 1.6 | | ≤1 | ≤1 | 1 | 2 | 3 | 3 | 5 |
| 35-49 | 105 | 1.6 | | ≤1 | ≤1 | 1 | 2 | 3 | 3 | 4 |
| 50-64 | 50 | 2.0 | 1 | ≤1 | ≤1 | 2 | 2 | 4 | 5 | 7 |
| 65+ | 13 | 2.6 | 4 | ≤1 | 1 | 2 | 3 | 4 | 8 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 135 | 2.0 | 1 | ≤1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 20-34 | 386 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 35-49 | 207 | 2.1 | 2 | ≤1 | 1 | 2 | 3 | 3 | 4 | 11 |
| 50-64 | 108 | 2.5 | 4 | 1 | 1 | 2 | 3 | 4 | 6 | 13 |
| 65+ | 37 | 3.2 | 12 | 1 | 1 | 2 | 3 | 7 | 13 | 17 |
| **TOT. SINGLE DX** | 435 | 1.7 | 1 | ≤1 | ≤1 | 1 | 2 | 3 | 3 | 5 |
| **" MULTIPLE DX** | 873 | 2.1 | 2 | 1 | 1 | 2 | 3 | 3 | 4 | 8 |
| TOTAL 0-19 YRS | 198 | 1.9 | 1 | ≤1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 20-34 | 590 | 1.8 | 1 | ≤1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 35-49 | 312 | 2.0 | 2 | ≤1 | 1 | 2 | 2 | 3 | 4 | 9 |
| 50-64 | 158 | 2.4 | 3 | ≤1 | 1 | 2 | 3 | 4 | 5 | 11 |
| 65+ | 50 | 3.1 | 10 | 1 | 1 | 2 | 3 | 6 | 9 | 17 |
| **GRAND TOTAL** | **1308** | **2.0** | **2** | **≤1** | **1** | **2** | **2** | **3** | **4** | **8** |

## 542: Rhinoplasty (21.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 427 | 1.7 | | ≤1 | 1 | 1 | 2 | 3 | 3 | 5 |
| 20-34 | 966 | 1.7 | 4 | ≤1 | 1 | 1 | 2 | 3 | 3 | 4 |
| 35-49 | 473 | 1.7 | | ≤1 | 1 | 1 | 2 | 3 | 4 | 5 |
| 50-64 | 235 | 1.8 | | ≤1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 65+ | 71 | 2.3 | 6 | ≤1 | 1 | 2 | 3 | 4 | 4 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 754 | 2.2 | 12 | ≤1 | 1 | 2 | 2 | 3 | 4 | 8 |
| 20-34 | 1590 | 2.1 | 9 | ≤1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 35-49 | 894 | 2.3 | 6 | ≤1 | 1 | 2 | 3 | 4 | 4 | 11 |
| 50-64 | 538 | 2.8 | 15 | ≤1 | 1 | 2 | 3 | 4 | 6 | 21 |
| 65+ | 174 | 3.1 | 8 | ≤1 | 1 | 2 | 3 | 6 | 8 | 16 |
| **TOT. SINGLE DX** | 2172 | 1.7 | 3 | ≤1 | 1 | 1 | 2 | 3 | 3 | 5 |
| **" MULTIPLE DX** | 3950 | 2.3 | 10 | ≤1 | 1 | 2 | 3 | 4 | 4 | 10 |
| TOTAL 0-19 YRS | 1181 | 2.0 | 8 | ≤1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 20-34 | 2556 | 2.0 | 8 | ≤1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 35-49 | 1367 | 2.1 | 4 | ≤1 | 1 | 2 | 2 | 3 | 4 | 8 |
| 50-64 | 773 | 2.5 | 12 | ≤1 | 1 | 2 | 3 | 4 | 5 | 14 |
| 65+ | 245 | 2.9 | 8 | ≤1 | 1 | 2 | 3 | 5 | 8 | 16 |
| **GRAND TOTAL** | **6122** | **2.1** | **8** | **≤1** | **1** | **2** | **2** | **3** | **4** | **8** |

## 543: Other incision, excision of nose (21.1; 21.22; 21.3-21.4; 21.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 79 | 1.5 | | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 5 |
| 20-34 | 129 | 1.7 | 2 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 9 |
| 35-49 | 123 | 1.7 | 1 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 6 |
| 50-64 | 161 | 1.7 | 1 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 5 |
| 65+ | 190 | 2.3 | 4 | ≤1 | ≤1 | 2 | 2 | 4 | 7 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 121 | 2.6 | 8 | ≤1 | 1 | 2 | 3 | 4 | 6 | 19 |
| 20-34 | 219 | 2.6 | 37 | ≤1 | 1 | 2 | 3 | 4 | 4 | 9 |
| 35-49 | 231 | 2.3 | 2 | ≤1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 50-64 | 212 | 3.3 | 44 | ≤1 | 1 | 2 | 3 | 5 | 6 | 31 |
| 65+ | 226 | 4.5 | 35 | ≤1 | 1 | 2 | 5 | 9 | 16 | 33 |
| **TOT. SINGLE DX** | 682 | 1.8 | 2 | ≤1 | ≤1 | 1 | 2 | 3 | 4 | 8 |
| **" MULTIPLE DX** | 1009 | 3.3 | 28 | ≤1 | 1 | 2 | 3 | 5 | 8 | 21 |
| TOTAL 0-19 YRS | 200 | 2.2 | 6 | ≤1 | ≤1 | 2 | 2 | 4 | 5 | 14 |
| 20-34 | 348 | 2.3 | 25 | ≤1 | ≤1 | 2 | 2 | 3 | 4 | 7 |
| 35-49 | 354 | 2.1 | 2 | ≤1 | ≤1 | 2 | 3 | 3 | 4 | 8 |
| 50-64 | 373 | 2.6 | 27 | ≤1 | ≤1 | 2 | 3 | 4 | 5 | 20 |
| 65+ | 416 | 3.5 | 22 | ≤1 | ≤1 | 2 | 4 | 7 | 11 | 23 |
| **GRAND TOTAL** | **1691** | **2.6** | **18** | **≤1** | **≤1** | **2** | **3** | **4** | **6** | **17** |

## 544: Miscellaneous operations on nose (21.02-21.09; 21.7; 21.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 227 | 1.2 | | ≤1 | ≤1 | 1 | 1 | 2 | 2 | 3 |
| 20-34 | 168 | 2.1 | 3 | ≤1 | ≤1 | 1 | 3 | 5 | 6 | 9 |
| 35-49 | 79 | 2.7 | 3 | ≤1 | 1 | 2 | 4 | 6 | 7 | 9 |
| 50-64 | 92 | 4.0 | 10 | 1 | 1 | 3 | 5 | 8 | 10 | 16 |
| 65+ | 51 | 3.6 | 7 | 1 | 1 | 3 | 5 | 7 | 8 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 323 | 2.3 | 24 | ≤1 | ≤1 | 1 | 2 | 4 | 6 | 11 |
| 20-34 | 403 | 2.7 | 15 | ≤1 | 1 | 2 | 3 | 5 | 7 | 21 |
| 35-49 | 179 | 3.6 | 37 | ≤1 | 1 | 2 | 4 | 7 | 9 | 21 |
| 50-64 | 167 | 5.1 | 41 | 1 | 1 | 4 | 6 | 9 | 11 | 30 |
| 65+ | 172 | 6.3 | 41 | 1 | 2 | 5 | 7 | 12 | 17 | 29 |
| **TOT. SINGLE DX** | 617 | 2.3 | 5 | ≤1 | ≤1 | 1 | 3 | 5 | 7 | 11 |
| **" MULTIPLE DX** | 1244 | 3.5 | 30 | ≤1 | 1 | 2 | 4 | 7 | 10 | 21 |
| TOTAL 0-19 YRS | 550 | 1.9 | 15 | ≤1 | ≤1 | 1 | 2 | 3 | 5 | 10 |
| 20-34 | 571 | 2.5 | 12 | ≤1 | ≤1 | 2 | 3 | 5 | 7 | 18 |
| 35-49 | 258 | 3.3 | 27 | ≤1 | 1 | 2 | 4 | 6 | 8 | 20 |
| 50-64 | 259 | 4.7 | 31 | 1 | 1 | 4 | 6 | 9 | 11 | 19 |
| 65+ | 223 | 5.7 | 35 | 1 | 1 | 4 | 7 | 10 | 16 | 29 |
| **GRAND TOTAL** | **1861** | **3.1** | **22** | **≤1** | **≤1** | **2** | **4** | **6** | **9** | **19** |

## 545: Radical maxillary antrotomy (22.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 32 | 1.9 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 20-34 | 129 | 2.2 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 35-49 | 142 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 7 |
| 50-64 | 113 | 2.2 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 65+ | 61 | 2.6 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 38 | 2.9 | 3 | 1 | 1 | 2 | 4 | 5 | 7 | 10 |
| 20-34 | 173 | 3.0 | 15 | 1 | 1 | 2 | 3 | 5 | 8 | 23 |
| 35-49 | 176 | 3.0 | 11 | 1 | 1 | 2 | 3 | 5 | 6 | 11 |
| 50-64 | 159 | 3.5 | 14 | 1 | 1 | 3 | 4 | 6 | 11 | 21 |
| 65+ | 96 | 5.6 | 82 | 1 | 1 | 3 | 5 | 11 | 17 | 59 |
| **TOT. SINGLE DX** | 477 | 2.3 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| **" MULTIPLE DX** | 642 | 3.5 | 24 | 1 | 1 | 2 | 3 | 6 | 9 | 21 |
| TOTAL 0-19 YRS | 70 | 2.4 | 3 | 1 | 1 | 2 | 3 | 4 | 6 | 10 |
| 20-34 | 302 | 2.6 | 10 | 1 | 1 | 2 | 3 | 4 | 6 | 15 |
| 35-49 | 318 | 2.7 | 7 | 1 | 1 | 2 | 3 | 4 | 6 | 9 |
| 50-64 | 272 | 3.0 | 10 | 1 | 1 | 2 | 3 | 5 | 8 | 18 |
| 65+ | 157 | 4.4 | 53 | 1 | 1 | 3 | 4 | 9 | 12 | 35 |
| **GRAND TOTAL** | **1119** | **3.0** | **15** | **1** | **1** | **2** | **3** | **5** | **7** | **17** |

## 546: Miscellaneous operations on accessory sinus (22.0-22.11; 22.2; 22.4-22.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 57 | 2.6 | 7 | <1 | <1 | 1 | 3 | 5 | 10 | 13 |
| 20-34 | 91 | 2.5 | 3 | <1 | <1 | 2 | 3 | 5 | 7 | 10 |
| 35-49 | 83 | 2.4 | 3 | <1 | 1 | 2 | 3 | 4 | 8 | 10 |
| 50-64 | 68 | 2.5 | 3 | 1 | 1 | 2 | 3 | 5 | 6 | 9 |
| 65+ | 37 | 3.6 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 84 | 5.3 | 54 | <1 | <1 | 2 | 7 | 12 | 17 | 41 |
| 20-34 | 203 | 3.9 | 25 | 1 | 1 | 3 | 4 | 7 | 9 | 32 |
| 35-49 | 194 | 3.5 | 22 | 1 | 1 | 2 | 4 | 7 | 10 | 27 |
| 50-64 | 181 | 4.5 | 25 | 1 | 1 | 3 | 5 | 9 | 15 | 27 |
| 65+ | 115 | 4.8 | 31 | 1 | 1 | 3 | 6 | 11 | 16 | 33 |
| **TOT. SINGLE DX** | 336 | 2.6 | 5 | <1 | <1 | 2 | 3 | 5 | 8 | 12 |
| **" MULTIPLE DX** | 777 | 4.2 | 29 | <1 | 1 | 3 | 5 | 9 | 13 | 30 |
| TOTAL 0-19 YRS | 141 | 4.2 | 37 | <1 | <1 | 2 | 4 | 10 | 14 | 32 |
| 20-34 | 294 | 3.5 | 19 | <1 | 1 | 2 | 4 | 7 | 8 | 29 |
| 35-49 | 277 | 3.2 | 17 | 1 | 1 | 2 | 3 | 7 | 9 | 25 |
| 50-64 | 249 | 3.9 | 20 | 1 | 1 | 3 | 5 | 8 | 11 | 26 |
| 65+ | 152 | 4.5 | 26 | 1 | 1 | 3 | 5 | 9 | 12 | 29 |
| **GRAND TOTAL** | **1113** | **3.7** | **22** | **<1** | **1** | **2** | **4** | **8** | **11** | **28** |

## 547: Tooth extraction (23.0-23.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 203 | 1.3 | | <1 | <1 | 1 | 1 | 2 | 2 | 6 |
| 20-34 | 136 | 1.7 | 2 | <1 | <1 | 1 | 2 | 3 | 3 | 9 |
| 35-49 | 65 | 2.0 | 2 | <1 | 1 | 2 | 2 | 3 | 5 | 10 |
| 50-64 | 48 | 2.0 | 3 | <1 | 1 | 2 | 2 | 3 | 3 | 14 |
| 65+ | 22 | 3.6 | 39 | <1 | <1 | 2 | 3 | 5 | 16 | 31 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 141 | 2.4 | 20 | <1 | <1 | 2 | 3 | 4 | 5 | 13 |
| 20-34 | 167 | 5.5 | 72 | <1 | 1 | 2 | 6 | 15 | 20 | 42 |
| 35-49 | 143 | 5.6 | 83 | <1 | 1 | 2 | 4 | 14 | 26 | 52 |
| 50-64 | 176 | 4.5 | 45 | 1 | 1 | 2 | 4 | 9 | 16 | 44 |
| 65+ | 250 | 5.4 | 52 | 1 | 1 | 2 | 5 | 14 | 22 | 42 |
| **TOT. SINGLE DX** | 474 | 1.7 | 4 | <1 | <1 | 1 | 2 | 3 | 3 | 9 |
| **" MULTIPLE DX** | 877 | 4.8 | 56 | <1 | 1 | 2 | 4 | 11 | 19 | 44 |
| TOTAL 0-19 YRS | 344 | 1.8 | 9 | <1 | <1 | 1 | 2 | 3 | 4 | 7 |
| 20-34 | 303 | 3.8 | 44 | <1 | 1 | 2 | 3 | 8 | 16 | 27 |
| 35-49 | 208 | 4.5 | 61 | <1 | 1 | 2 | 3 | 11 | 19 | 46 |
| 50-64 | 224 | 4.0 | 38 | <1 | 1 | 2 | 3 | 8 | 14 | 34 |
| 65+ | 272 | 5.2 | 51 | 1 | 1 | 2 | 5 | 14 | 22 | 42 |
| **GRAND TOTAL** | **1351** | **3.7** | **40** | **<1** | **<1** | **2** | **3** | **8** | **15** | **31** |

## 548: Tooth restoration and orthodontia (23.2-23.6; 24.6-24.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 188 | 1.1 | | <1 | <1 | <1 | 1 | 1 | 2 | 2 |
| 20-34 | 26 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 3 |
| 35-49 | 13 | 1.9 | | <1 | <1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 5 | 3.2 | 3 | <1 | <1 | 3 | 4 | 6 | 6 | 6 |
| 65+ | 5 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 139 | 1.3 | 2 | <1 | <1 | <1 | 1 | 2 | 2 | 8 |
| 20-34 | 35 | 2.3 | 3 | <1 | <1 | 2 | 2 | 5 | 7 | 10 |
| 35-49 | 22 | 1.8 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 50-64 | 9 | 1.7 | 1 | 1 | 1 | 1 | 2 | 3 | 4 | 4 |
| 65+ | 8 | 9.8 | 331 | <1 | <1 | 2 | 8 | 41 | 54 | 54 |
| **TOT. SINGLE DX** | 237 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 3 |
| **" MULTIPLE DX** | 213 | 1.8 | 16 | <1 | <1 | 1 | 2 | 2 | 4 | 13 |
| TOTAL 0-19 YRS | 327 | 1.1 | 1 | <1 | <1 | <1 | 1 | 1 | 2 | 2 |
| 20-34 | 61 | 1.8 | 3 | <1 | <1 | 1 | 2 | 3 | 5 | 10 |
| 35-49 | 35 | 1.8 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 5 |
| 50-64 | 14 | 2.2 | 2 | <1 | 1 | 2 | 3 | 4 | 6 | 6 |
| 65+ | 13 | 6.6 | 210 | <1 | <1 | 2 | 4 | 20 | 48 | 54 |
| **GRAND TOTAL** | **450** | **1.5** | **8** | **<1** | **<1** | **1** | **1** | **2** | **3** | **7** |

## 549: Operations on teeth and gums except extraction and restoration (23.7; 24.0-24.12; 24.2-24.5; 24.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 41 | 2.0 | 2 | <1 | <1 | 2 | 2 | 4 | 4 | 11 |
| 20-34 | 58 | 1.7 | 2 | 1 | 1 | 1 | 2 | 3 | 4 | 10 |
| 35-49 | 104 | 1.7 | 1 | 1 | 1 | 1 | 2 | 3 | 4 | 5 |
| 50-64 | 90 | 2.6 | 1 | 1 | 1 | 3 | 3 | 4 | 5 | 6 |
| 65+ | 31 | 3.1 | 4 | 1 | 1 | 2 | 4 | 6 | 8 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 39 | 2.6 | 4 | <1 | <1 | 2 | 3 | 4 | 7 | 10 |
| 20-34 | 44 | 2.8 | 6 | 1 | 1 | 2 | 3 | 6 | 7 | 14 |
| 35-49 | 61 | 3.8 | 16 | 1 | 1 | 2 | 4 | 7 | 10 | 26 |
| 50-64 | 79 | 3.4 | 10 | 1 | 1 | 3 | 4 | 5 | 6 | 23 |
| 65+ | 35 | 5.7 | 50 | 1 | 1 | 3 | 6 | 10 | 27 | 31 |
| **TOT. SINGLE DX** | 324 | 2.1 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| **" MULTIPLE DX** | 258 | 3.6 | 16 | 1 | 1 | 3 | 4 | 6 | 9 | 28 |
| TOTAL 0-19 YRS | 80 | 2.3 | 4 | <1 | <1 | 2 | 3 | 4 | 5 | 11 |
| 20-34 | 102 | 2.1 | 4 | 1 | 1 | 2 | 2 | 4 | 6 | 12 |
| 35-49 | 165 | 2.5 | 8 | 1 | 1 | 2 | 3 | 4 | 6 | 15 |
| 50-64 | 169 | 2.9 | 6 | 1 | 1 | 3 | 4 | 4 | 5 | 11 |
| 65+ | 66 | 4.5 | 31 | 1 | 1 | 3 | 5 | 8 | 12 | 31 |
| **GRAND TOTAL** | **582** | **2.7** | **9** | **1** | **1** | **2** | **3** | **5** | **6** | **15** |

## 550: Operations on salivary glands and ducts (26.0-26.11; 26.2-26.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 89 | 2.2 | 1 | <1 | <1 | 2 | 3 | 4 | 4 | 8 |
| 20-34 | 200 | 2.6 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 10 |
| 35-49 | 220 | 2.8 | 1 | 1 | 1 | 3 | 3 | 4 | 5 | 7 |
| 50-64 | 254 | 2.8 | 2 | 1 | 1 | 3 | 4 | 4 | 5 | 8 |
| 65+ | 187 | 3.5 | 4 | 1 | 2 | 3 | 4 | 5 | 7 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 23 | 3.3 | 4 | <1 | 1 | 3 | 5 | 6 | 8 | 9 |
| 20-34 | 87 | 3.2 | 7 | 1 | 1 | 3 | 3 | 6 | 9 | 15 |
| 35-49 | 123 | 4.0 | 13 | 1 | 1 | 3 | 4 | 7 | 11 | 22 |
| 50-64 | 174 | 4.0 | 12 | 1 | 2 | 3 | 4 | 7 | 11 | 24 |
| 65+ | 221 | 5.5 | 30 | 2 | 2 | 4 | 6 | 11 | 16 | 24 |
| **TOT. SINGLE DX** | 950 | 2.8 | 3 | 1 | 1 | 3 | 4 | 5 | 5 | 9 |
| **" MULTIPLE DX** | 628 | 4.4 | 18 | 1 | 1 | 3 | 5 | 8 | 13 | 22 |
| TOTAL 0-19 YRS | 112 | 2.4 | 2 | <1 | <1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 287 | 2.8 | 4 | <1 | 1 | 2 | 3 | 5 | 7 | 11 |
| 35-49 | 343 | 3.2 | 6 | 1 | 1 | 3 | 4 | 5 | 6 | 13 |
| 50-64 | 428 | 3.3 | 7 | 1 | 1 | 3 | 4 | 5 | 7 | 17 |
| 65+ | 408 | 4.6 | 19 | 1 | 2 | 3 | 5 | 8 | 13 | 21 |
| **GRAND TOTAL** | **1578** | **3.4** | **10** | **1** | **1** | **3** | **4** | **5** | **8** | **18** |

## 551: Palatoplasty (27.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 160 | 3.4 | 2 | 1 | 1 | 3 | 4 | 6 | 6 | 7 |
| 20-34 | 7 | 4.1 | 1 | 3 | 3 | 4 | 5 | 6 | 6 | 6 |
| 35-49 | 6 | 3.2 | 1 | 2 | 2 | 3 | 4 | 4 | 4 | 4 |
| 50-64 | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 136 | 3.7 | 4 | 1 | 1 | 3 | 5 | 6 | 7 | 10 |
| 20-34 | 6 | 3.2 | 3 | 1 | 1 | 3 | 4 | 6 | 6 | 6 |
| 35-49 | 3 | 4.0 | 13 | 1 | 1 | 3 | 7 | 8 | 8 | 8 |
| 50-64 | 2 | 3.5 | 13 | <1 | <1 | 3 | 6 | 6 | 6 | 6 |
| 65+ | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| **TOT. SINGLE DX** | 174 | 3.4 | 2 | 1 | 1 | 3 | 4 | 6 | 6 | 7 |
| **" MULTIPLE DX** | 148 | 3.7 | 4 | 1 | 1 | 3 | 5 | 6 | 7 | 9 |
| TOTAL 0-19 YRS | 296 | 3.5 | 3 | 1 | 1 | 3 | 5 | 6 | 6 | 8 |
| 20-34 | 13 | 3.7 | 2 | 1 | 2 | 4 | 5 | 6 | 6 | 6 |
| 35-49 | 9 | 3.4 | 4 | 1 | 1 | 3 | 4 | 7 | 8 | 8 |
| 50-64 | 3 | 3.3 | 6 | <1 | <1 | 3 | 5 | 6 | 6 | 6 |
| 65+ | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| **GRAND TOTAL** | **322** | **3.5** | **3** | **1** | **1** | **3** | **5** | **6** | **6** | **8** |

## 552: Operations on tongue, oral cavity and facial region except palatoplasty and operations on salivary glands (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 299 | 2.9 | 6 | <1 | <1 | 2 | 4 | 6 | 7 | 11 |
| 20-34 | 113 | 3.4 | 56 | <1 | <1 | 2 | 3 | 6 | 8 | 44 |
| 35-49 | 95 | 2.8 | 8 | <1 | <1 | 2 | 3 | 6 | 10 | 15 |
| 50-64 | 175 | 3.8 | 58 | <1 | <1 | 2 | 4 | 8 | 12 | 26 |
| 65+ | 187 | 3.0 | 8 | <1 | <1 | 2 | 3 | 6 | 10 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 237 | 3.7 | 17 | <1 | 1 | 2 | 5 | 7 | 10 | 24 |
| 20-34 | 198 | 4.7 | 34 | <1 | 1 | 3 | 6 | 9 | 13 | 31 |
| 35-49 | 134 | 6.8 | 76 | <1 | 1 | 3 | 7 | 19 | 28 | 40 |
| 50-64 | 230 | 7.5 | 113 | 1 | 1 | 3 | 8 | 21 | 30 | 57 |
| 65+ | 349 | 8.1 | 112 | 1 | 1 | 4 | 10 | 18 | 28 | 68 |
| **TOT. SINGLE DX** | 869 | 3.1 | 24 | <1 | <1 | 2 | 4 | 6 | 9 | 17 |
| **" MULTIPLE DX** | 1148 | 6.3 | 78 | <1 | 1 | 3 | 7 | 16 | 22 | 51 |
| TOTAL 0-19 YRS | 536 | 3.2 | 12 | <1 | <1 | 2 | 4 | 6 | 8 | 17 |
| 20-34 | 311 | 4.2 | 42 | <1 | <1 | 2 | 5 | 9 | 11 | 30 |
| 35-49 | 229 | 5.1 | 52 | <1 | <1 | 3 | 5 | 11 | 22 | 38 |
| 50-64 | 405 | 5.9 | 93 | <1 | 1 | 3 | 6 | 15 | 24 | 54 |
| 65+ | 536 | 6.3 | 82 | <1 | 1 | 3 | 7 | 15 | 21 | 53 |
| **GRAND TOTAL** | **2017** | **5.0** | **57** | **<1** | **<1** | **3** | **5** | **11** | **17** | **42** |

## 553: Tonsillectomy without adenoidectomy (28.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4437 | 1.5 | | <1 | 1 | 1 | 2 | 2 | 3 | 3 |
| 20-34 | 2859 | 1.7 | | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 35-49 | 288 | 1.9 | | 1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 50-64 | 41 | 2.8 | 50 | 1 | 1 | 2 | 2 | 3 | 4 | 47 |
| 65+ | 16 | 2.8 | 24 | <1 | <1 | 2 | 2 | 3 | 16 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 958 | 1.8 | 2 | <1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 20-34 | 653 | 2.0 | 2 | 1 | 1 | 2 | 2 | 3 | 4 | 6 |
| 35-49 | 93 | 2.3 | 3 | 1 | 1 | 2 | 3 | 4 | 6 | 11 |
| 50-64 | 36 | 3.7 | 30 | 1 | 1 | 2 | 2 | 12 | 19 | 25 |
| 65+ | 9 | 8.9 | 356 | <1 | <1 | 3 | 4 | 38 | 59 | 59 |
| **TOT. SINGLE DX** | 7641 | 1.6 | 1 | <1 | 1 | 1 | 2 | 2 | 3 | 4 |
| **" MULTIPLE DX** | 1749 | 2.0 | 5 | <1 | 1 | 2 | 2 | 3 | 4 | 8 |
| TOTAL 0-19 YRS | 5395 | 1.6 | 1 | <1 | 1 | 1 | 2 | 2 | 3 | 4 |
| 20-34 | 3512 | 1.7 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 35-49 | 381 | 2.0 | 2 | 1 | 1 | 2 | 2 | 3 | 4 | 7 |
| 50-64 | 77 | 3.2 | 41 | 1 | 1 | 2 | 2 | 4 | 13 | 41 |
| 65+ | 25 | 5.0 | 143 | <1 | <1 | 2 | 3 | 7 | 30 | 59 |
| **GRAND TOTAL** | **9390** | **1.7** | **2** | **<1** | **1** | **1** | **2** | **3** | **3** | **4** |

## 554: Tonsillectomy with adenoidectomy (28.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 6627 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 2 | 3 |
| 20-34 | 176 | 1.7 | | <1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 35-49 | 9 | 1.6 | | 1 | 1 | 1 | 2 | 3 | 3 | 3 |
| 50-64 | 2 | 1.5 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 65+ | 3 | 1.3 | | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3874 | 1.4 | | <1 | <1 | 1 | 2 | 2 | 2 | 4 |
| 20-34 | 72 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 35-49 | 6 | 2.3 | 2 | 1 | 1 | 2 | 3 | 5 | 5 | 5 |
| 50-64 | 3 | 1.7 | | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| 65+ | 2 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| **TOT. SINGLE DX** | 6817 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 3 |
| **" MULTIPLE DX** | 3957 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 4 |
| TOTAL 0-19 YRS | 10501 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 3 |
| 20-34 | 248 | 1.8 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 35-49 | 15 | 1.9 | 1 | 1 | 1 | 2 | 2 | 3 | 5 | 5 |
| 50-64 | 5 | 1.6 | | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| 65+ | 5 | 1.6 | | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| **GRAND TOTAL** | **10774** | **1.5** | **1** | **<1** | **<1** | **1** | **2** | **2** | **2** | **3** |

## 555: Adenoidectomy without tonsillectomy (28.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 448 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| 20-34 | 4 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 2 | 2 |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 918 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 4 |
| 20-34 | 9 | 1.4 | 1 | 1 | 1 | 1 | 2 | 3 | 4 | 4 |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| **TOT. SINGLE DX** | 452 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| **" MULTIPLE DX** | 928 | 1.2 | 1 | <1 | <1 | 1 | 1 | 2 | 2 | 4 |
| TOTAL 0-19 YRS | 1366 | 1.2 | 1 | <1 | <1 | 1 | 1 | 2 | 2 | 3 |
| 20-34 | 13 | 1.5 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 4 |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| **GRAND TOTAL** | **1380** | **1.2** | **1** | **<1** | **<1** | **1** | **1** | **2** | **2** | **3** |

## 556: Operations on pharynx except tonsillectomy and adenoidectomy (28.0-28.11; 28.4-28.5; 28.7-28.9; 29.0; 29.12; 29.2-29.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 312 | 2.1 | 3 | <1 | <1 | 2 | 3 | 4 | 5 | 10 |
| 20-34 | 224 | 2.3 | 8 | <1 | 1 | 2 | 3 | 4 | 5 | 13 |
| 35-49 | 71 | 2.1 | 1 | <1 | <1 | 2 | 3 | 4 | 4 | 6 |
| 50-64 | 81 | 2.8 | 6 | <1 | 1 | 2 | 3 | 6 | 8 | 15 |
| 65+ | 64 | 3.6 | 19 | <1 | <1 | 2 | 4 | 8 | 13 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 115 | 2.7 | 8 | <1 | <1 | 2 | 3 | 6 | 8 | 17 |
| 20-34 | 92 | 3.8 | 54 | <1 | 1 | 2 | 4 | 6 | 7 | 51 |
| 35-49 | 59 | 5.7 | 76 | <1 | 1 | 3 | 5 | 12 | 24 | 46 |
| 50-64 | 85 | 5.8 | 75 | <1 | 1 | 3 | 6 | 12 | 19 | 53 |
| 65+ | 94 | 9.6 | 182 | 1 | 1 | 5 | 12 | 20 | 40 | 73 |
| **TOT. SINGLE DX** | 752 | 2.4 | 6 | <1 | <1 | 2 | 3 | 4 | 6 | 14 |
| **" MULTIPLE DX** | 445 | 5.4 | 82 | <1 | 1 | 2 | 5 | 12 | 19 | 56 |
| TOTAL 0-19 YRS | 427 | 2.3 | 5 | <1 | <1 | 2 | 3 | 4 | 6 | 13 |
| 20-34 | 316 | 2.7 | 22 | <1 | 1 | 2 | 3 | 5 | 6 | 27 |
| 35-49 | 130 | 3.7 | 38 | <1 | <1 | 2 | 4 | 7 | 12 | 43 |
| 50-64 | 166 | 4.3 | 44 | <1 | 1 | 2 | 5 | 8 | 14 | 38 |
| 65+ | 158 | 7.2 | 124 | <1 | 1 | 3 | 8 | 16 | 23 | 68 |
| **GRAND TOTAL** | **1197** | **3.5** | **37** | **<1** | **1** | **2** | **3** | **6** | **12** | **31** |

## 557: Other diagnostic and therapeutic procedures on nose, mouth, and pharynx (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 41 | 2.1 | 6 | <1 | <1 | <1 | 2 | 5 | 8 | 12 |
| 20-34 | 17 | 2.4 | 1 | <1 | 1 | 3 | 3 | 4 | 4 | 4 |
| 35-49 | 16 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 5 | 5 |
| 50-64 | 32 | 2.7 | 5 | <1 | 1 | 2 | 4 | 6 | 8 | 9 |
| 65+ | 31 | 3.2 | 5 | 1 | 1 | 2 | 4 | 6 | 10 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 52 | 2.7 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 11 | 4.0 | 33 | <1 | <1 | 2 | 4 | 11 | 20 | 21 |
| 35-49 | 13 | 4.7 | 10 | <1 | 1 | 4 | 8 | 9 | 10 | 10 |
| 50-64 | 50 | 5.0 | 25 | <1 | 1 | 4 | 6 | 10 | 15 | 27 |
| 65+ | 100 | 5.3 | 28 | 1 | 1 | 4 | 7 | 11 | 15 | 30 |
| **TOT. SINGLE DX** | 137 | 2.6 | 5 | <1 | <1 | 2 | 4 | 5 | 7 | 10 |
| **" MULTIPLE DX** | 226 | 4.5 | 22 | 1 | 1 | 3 | 5 | 9 | 13 | 25 |
| TOTAL 0-19 YRS | 93 | 2.4 | 4 | <1 | <1 | 2 | 3 | 5 | 6 | 11 |
| 20-34 | 28 | 3.0 | 14 | <1 | <1 | 2 | 3 | 4 | 6 | 21 |
| 35-49 | 29 | 3.8 | 6 | 1 | 1 | 3 | 5 | 8 | 9 | 10 |
| 50-64 | 82 | 4.1 | 18 | <1 | 1 | 3 | 5 | 8 | 11 | 25 |
| 65+ | 131 | 4.8 | 24 | 1 | 1 | 4 | 6 | 10 | 14 | 26 |
| **GRAND TOTAL** | **363** | **3.8** | **16** | **<1** | **1** | **3** | **5** | **8** | **11** | **22** |

## 558: Excision of lesion of larynx (30.0; 31.43)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 87 | 1.4 | 2 | <1 | <1 | 1 | 1 | 2 | 3 | 11 |
| 20-34 | 122 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 3 |
| 35-49 | 180 | 1.7 | 5 | <1 | <1 | 1 | 2 | 2 | 3 | 9 |
| 50-64 | 321 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| 65+ | 240 | 2.0 | 12 | <1 | <1 | 2 | 2 | 2 | 3 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 38 | 2.7 | 34 | <1 | <1 | 1 | 2 | 5 | 8 | 36 |
| 20-34 | 23 | 4.7 | 167 | <1 | <1 | 2 | 3 | 6 | 14 | 70 |
| 35-49 | 96 | 2.8 | 30 | <1 | <1 | 2 | 2 | 5 | 8 | 36 |
| 50-64 | 241 | 3.0 | 18 | <1 | <1 | 2 | 2 | 6 | 11 | 25 |
| 65+ | 230 | 4.1 | 50 | <1 | 1 | 2 | 3 | 9 | 17 | 47 |
| **TOT. SINGLE DX** | 950 | 1.6 | 5 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| **" MULTIPLE DX** | 633 | 3.4 | 40 | <1 | <1 | 2 | 3 | 7 | 12 | 35 |
| TOTAL 0-19 YRS | 125 | 1.8 | 12 | <1 | <1 | 1 | 1 | 2 | 4 | 19 |
| 20-34 | 150 | 1.8 | 32 | <1 | <1 | 1 | 1 | 2 | 3 | 8 |
| 35-49 | 276 | 2.1 | 14 | <1 | <1 | 1 | 2 | 3 | 5 | 16 |
| 50-64 | 562 | 2.1 | 9 | <1 | <1 | 1 | 2 | 3 | 6 | 18 |
| 65+ | 470 | 3.0 | 32 | <1 | <1 | 2 | 2 | 5 | 9 | 30 |
| **GRAND TOTAL** | **1583** | **2.3** | **19** | **<1** | **<1** | **1** | **2** | **3** | **6** | **20** |

## 559: Partial laryngectomy and local excision of lesion of trachea (30.1-30.2; 31.44; 31.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 2.2 | 2 | 1 | 1 | 2 | 3 | 5 | 5 | 5 |
| 20-34 | 5 | 1.4 | | <1 | <1 | 1 | 2 | 2 | 2 | 2 |
| 35-49 | 12 | 4.2 | 46 | <1 | <1 | 1 | 2 | 17 | 20 | 21 |
| 50-64 | 25 | 6.2 | 34 | <1 | <1 | 2 | 13 | 15 | 16 | 16 |
| 65+ | 20 | 6.7 | 21 | <1 | 1 | 7 | 11 | 13 | 14 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 6.9 | 306 | <1 | 1 | 2 | 3 | 9 | 53 | 72 |
| 20-34 | 8 | 2.9 | 6 | 1 | 1 | 2 | 3 | 7 | 9 | 9 |
| 35-49 | 10 | 13.2 | 87 | <1 | 1 | 13 | 21 | 25 | 28 | 28 |
| 50-64 | 31 | 13.5 | 199 | 1 | 1 | 9 | 19 | 27 | 52 | 61 |
| 65+ | 27 | 11.0 | 154 | 1 | 2 | 8 | 14 | 22 | 34 | 61 |
| **TOT. SINGLE DX** | 67 | 5.3 | 30 | <1 | <1 | 2 | 9 | 14 | 16 | 20 |
| **" MULTIPLE DX** | 92 | 10.7 | 181 | 1 | 1 | 6 | 14 | 23 | 30 | 67 |
| TOTAL 0-19 YRS | 21 | 5.8 | 234 | <1 | 1 | 2 | 3 | 7 | 38 | 72 |
| 20-34 | 13 | 2.3 | 5 | <1 | <1 | 2 | 2 | 4 | 8 | 9 |
| 35-49 | 22 | 8.3 | 83 | <1 | <1 | 2 | 16 | 21 | 24 | 28 |
| 50-64 | 56 | 10.2 | 138 | <1 | 1 | 7 | 15 | 22 | 28 | 61 |
| 65+ | 47 | 9.2 | 101 | 1 | 2 | 8 | 13 | 18 | 24 | 61 |
| **GRAND TOTAL** | **159** | **8.4** | **124** | **<1** | **1** | **4** | **12** | **20** | **25** | **61** |

## 560: Complete laryngectomy (30.3-30.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 20-34 | | | | | | | | | | |
| 35-49 | 3 | 13.7 | 9 | 11 | 11 | 13 | 16 | 17 | 17 | 17 |
| 50-64 | 25 | 13.8 | 27 | 7 | 8 | 13 | 17 | 21 | 24 | 25 |
| 65+ | 17 | 16.2 | 109 | 8 | 9 | 13 | 19 | 28 | 44 | 51 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 10.0 | | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 20-34 | | | | | | | | | | |
| 35-49 | 7 | 17.1 | 123 | 6 | 7 | 15 | 16 | 36 | 41 | 41 |
| 50-64 | 65 | 19.7 | 148 | 9 | 10 | 16 | 24 | 37 | 39 | 78 |
| 65+ | 53 | 21.0 | 169 | 8 | 10 | 18 | 27 | 42 | 73 | 99+ |
| **TOT. SINGLE DX** | 46 | 14.4 | 59 | 7 | 8 | 13 | 17 | 21 | 26 | 51 |
| **" MULTIPLE DX** | 126 | 20.0 | 154 | 8 | 10 | 17 | 25 | 38 | 42 | 99+ |
| TOTAL 0-19 YRS | 2 | 6.0 | 32 | 2 | 2 | 6 | 10 | 10 | 10 | 10 |
| 20-34 | | | | | | | | | | |
| 35-49 | 10 | 16.1 | 87 | 6 | 8 | 15 | 16 | 29 | 41 | 41 |
| 50-64 | 90 | 18.0 | 121 | 8 | 9 | 15 | 20 | 32 | 38 | 67 |
| 65+ | 70 | 19.8 | 157 | 8 | 10 | 17 | 24 | 41 | 51 | 99+ |
| **GRAND TOTAL** | **172** | **18.5** | **134** | **8** | **9** | **15** | **21** | **36** | **41** | **96** |

## 561: Permanent tracheostomy (31.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | 3 | 10.3 | 25 | 5 | 5 | 11 | 14 | 15 | 15 | 15 |
| 50-64 | | | | | | | | | | |
| 65+ | 2 | 18.5 | 265 | 7 | 7 | 18 | 30 | 30 | 30 | 30 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 14.8 | 136 | 4 | 4 | 10 | 21 | 34 | 34 | 34 |
| 20-34 | 8 | 34.4 | 174 | 9 | 15 | 33 | 45 | 50 | 51 | 51 |
| 35-49 | 5 | 15.2 | 37 | 8 | 8 | 13 | 21 | 23 | 23 | 23 |
| 50-64 | 20 | 22.7 | 556 | 1 | 2 | 12 | 36 | 63 | 71 | 73 |
| 65+ | 19 | 27.9 | 446 | 2 | 4 | 25 | 47 | 70 | 90 | 99+ |
| **TOT. SINGLE DX** | 5 | 13.6 | 99 | 5 | 5 | 11 | 19 | 30 | 30 | 30 |
| **" MULTIPLE DX** | 57 | 24.7 | 402 | 2 | 4 | 18 | 37 | 56 | 72 | 98 |
| TOTAL 0-19 YRS | 5 | 14.8 | 137 | 4 | 4 | 10 | 21 | 34 | 34 | 34 |
| 20-34 | 8 | 34.4 | 174 | 9 | 15 | 33 | 45 | 50 | 51 | 51 |
| 35-49 | 8 | 13.4 | 35 | 5 | 6 | 12 | 17 | 22 | 23 | 23 |
| 50-64 | 20 | 22.7 | 557 | 1 | 2 | 12 | 36 | 63 | 71 | 73 |
| 65+ | 21 | 27.0 | 423 | 2 | 5 | 25 | 45 | 66 | 88 | 99+ |
| **GRAND TOTAL** | **62** | **23.8** | **385** | **2** | **4** | **17** | **34** | **55** | **71** | **97** |

## 562: Operations on larynx and trachea except excision and permanent tracheostomy (31.0-31.1; 31.3; 31.6-31.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 32 | 5.5 | 65 | <1 | <1 | 2 | 7 | 12 | 20 | 42 |
| 20-34 | 14 | 6.4 | 41 | <1 | <1 | 4 | 10 | 14 | 20 | 22 |
| 35-49 | 16 | 5.1 | 35 | <1 | <1 | 2 | 7 | 10 | 20 | 24 |
| 50-64 | 31 | 8.9 | 108 | 1 | 2 | 6 | 11 | 18 | 31 | 52 |
| 65+ | 22 | 10.0 | 426 | <1 | 1 | 3 | 6 | 28 | 57 | 95 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 65 | 15.9 | 421 | 1 | 1 | 9 | 24 | 67 | 99+ | 99+ |
| 20-34 | 65 | 24.2 | 570 | 1 | 1 | 15 | 46 | 77 | 99+ | 99+ |
| 35-49 | 51 | 27.2 | 647 | 2 | 3 | 23 | 57 | 99+ | 99+ | 99+ |
| 50-64 | 122 | 24.9 | 606 | 1 | 1 | 16 | 45 | 74 | 93 | 99+ |
| 65+ | 156 | 32.5 | 707 | 2 | 2 | 30 | 60 | 93 | 99+ | 99+ |
| **TOT. SINGLE DX** | 115 | 7.3 | 138 | <1 | 1 | 4 | 8 | 18 | 26 | 67 |
| **" MULTIPLE DX** | 459 | 26.4 | 639 | 1 | 2 | 20 | 48 | 86 | 99+ | 99+ |
| TOTAL 0-19 YRS | 97 | 12.5 | 326 | <1 | 1 | 6 | 18 | 44 | 92 | 99+ |
| 20-34 | 79 | 21.0 | 522 | 1 | 1 | 13 | 40 | 69 | 89 | 99+ |
| 35-49 | 67 | 21.9 | 589 | 1 | 2 | 15 | 47 | 95 | 99+ | 99+ |
| 50-64 | 153 | 21.6 | 546 | 1 | 1 | 13 | 40 | 67 | 91 | 99+ |
| 65+ | 178 | 29.7 | 725 | 1 | 2 | 27 | 53 | 92 | 99+ | 99+ |
| **GRAND TOTAL** | **574** | **22.6** | **596** | **1** | **1** | **14** | **42** | **75** | **99+** | **99+** |

## 563: Incision and local excision or destruction of lesion of bronchus (32.0; 33.0; 33.24-33.25)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 7.2 | 57 | <1 | <1 | 5 | 11 | 19 | 23 | 23 |
| 20-34 | 46 | 7.5 | 171 | <1 | <1 | 3 | 10 | 15 | 19 | 86 |
| 35-49 | 92 | 4.5 | 39 | 1 | 1 | 3 | 6 | 10 | 12 | 38 |
| 50-64 | 231 | 5.1 | 38 | 1 | 1 | 3 | 7 | 11 | 17 | 33 |
| 65+ | 301 | 5.9 | 49 | <1 | 1 | 3 | 8 | 14 | 18 | 33 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 21 | 9.5 | 112 | <1 | 1 | 5 | 14 | 26 | 34 | 41 |
| 20-34 | 94 | 8.9 | 67 | 2 | 2 | 7 | 11 | 18 | 27 | 42 |
| 35-49 | 247 | 9.2 | 89 | 1 | 2 | 7 | 12 | 20 | 29 | 53 |
| 50-64 | 829 | 9.5 | 87 | 1 | 2 | 7 | 12 | 20 | 27 | 45 |
| 65+ | 1423 | 11.6 | 123 | 1 | 2 | 9 | 16 | 24 | 31 | 61 |
| **TOT. SINGLE DX** | 680 | 5.5 | 53 | <1 | 1 | 3 | 7 | 13 | 17 | 34 |
| **" MULTIPLE DX** | 2614 | 10.6 | 108 | 1 | 2 | 8 | 14 | 23 | 30 | 55 |
| TOTAL 0-19 YRS | 31 | 8.8 | 94 | <1 | 1 | 5 | 13 | 23 | 28 | 41 |
| 20-34 | 140 | 8.5 | 101 | 1 | 1 | 6 | 10 | 18 | 26 | 51 |
| 35-49 | 339 | 7.9 | 80 | 1 | 1 | 5 | 10 | 18 | 24 | 53 |
| 50-64 | 1060 | 8.5 | 80 | 1 | 1 | 6 | 11 | 18 | 25 | 45 |
| 65+ | 1724 | 10.6 | 115 | 1 | 2 | 8 | 14 | 23 | 30 | 57 |
| **GRAND TOTAL** | **3294** | **9.5** | **101** | **1** | **1** | **7** | **13** | **21** | **28** | **53** |

## 564: Bronchial resection and miscellaneous operations on bronchus (32.1; 33.41-33.42; 33.48; 33.91-33.92; 33.98)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 50-64 | 1 | 1.0 | | <1 | <1 | <1 | <1 | <1 | <1 | <1 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 20.0 | 242 | 9 | 9 | 20 | 31 | 31 | 31 | 31 |
| 20-34 | 2 | 67.5 | 1625 | 39 | 39 | 67 | 96 | 96 | 96 | 96 |
| 35-49 | | | | | | | | | | |
| 50-64 | 4 | 13.8 | 47 | 6 | 6 | 13 | 19 | 22 | 22 | 22 |
| 65+ | 2 | 13.0 | 242 | 2 | 2 | 13 | 24 | 24 | 24 | 24 |
| **TOT. SINGLE DX** | 2 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 2 | 2 |
| **" MULTIPLE DX** | 10 | 25.6 | 745 | 2 | 4 | 19 | 31 | 67 | 96 | 96 |
| TOTAL 0-19 YRS | 2 | 20.0 | 242 | 9 | 9 | 20 | 31 | 31 | 31 | 31 |
| 20-34 | 2 | 67.5 | 1625 | 39 | 39 | 67 | 96 | 96 | 96 | 96 |
| 35-49 | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 50-64 | 5 | 11.2 | 68 | <1 | <1 | 11 | 17 | 22 | 22 | 22 |
| 65+ | 2 | 13.0 | 242 | 2 | 2 | 13 | 24 | 24 | 24 | 24 |
| **GRAND TOTAL** | **12** | **21.6** | **697** | **<1** | **1** | **13** | **27** | **56** | **90** | **96** |

## 565: Local excision or destruction of lesion or tissue of lung (32.2-32.3; 33.26-33.27)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 10.2 | 25 | 5 | 6 | 8 | 17 | 18 | 18 | 18 |
| 20-34 | 50 | 7.6 | 18 | 2 | 2 | 7 | 9 | 14 | 18 | 18 |
| 35-49 | 91 | 7.3 | 14 | 2 | 3 | 7 | 9 | 11 | 13 | 21 |
| 50-64 | 215 | 7.6 | 33 | 1 | 1 | 7 | 10 | 16 | 19 | 29 |
| 65+ | 197 | 7.9 | 38 | 1 | 1 | 7 | 11 | 14 | 17 | 31 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 31 | 10.4 | 38 | 4 | 5 | 9 | 12 | 19 | 21 | 34 |
| 20-34 | 119 | 12.8 | 120 | 3 | 5 | 10 | 14 | 24 | 35 | 59 |
| 35-49 | 200 | 11.4 | 72 | 2 | 3 | 9 | 14 | 23 | 27 | 36 |
| 50-64 | 585 | 11.5 | 77 | 1 | 3 | 9 | 15 | 24 | 30 | 50 |
| 65+ | 798 | 12.5 | 98 | 2 | 2 | 10 | 17 | 25 | 31 | 48 |
| **TOT. SINGLE DX** | 563 | 7.7 | 31 | 1 | 1 | 7 | 10 | 14 | 18 | 27 |
| **" MULTIPLE DX** | 1733 | 12.0 | 89 | 2 | 3 | 10 | 16 | 24 | 31 | 48 |
| TOTAL 0-19 YRS | 41 | 10.3 | 34 | 4 | 5 | 9 | 12 | 18 | 20 | 34 |
| 20-34 | 169 | 11.2 | 96 | 2 | 4 | 9 | 12 | 21 | 29 | 51 |
| 35-49 | 291 | 10.1 | 58 | 2 | 3 | 8 | 13 | 21 | 25 | 34 |
| 50-64 | 800 | 10.4 | 68 | 1 | 2 | 9 | 13 | 21 | 28 | 44 |
| 65+ | 995 | 11.6 | 90 | 1 | 2 | 9 | 15 | 22 | 30 | 47 |
| **GRAND TOTAL** | **2296** | **10.9** | **78** | **1** | **2** | **9** | **14** | **22** | **28** | **46** |

## 566: Lobectomy and complete pneumonectomy (32.4-32.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 7.7 | 9 | 5 | 5 | 7 | 10 | 11 | 11 | 11 |
| 20-34 | 10 | 11.6 | 57 | 5 | 5 | 9 | 14 | 23 | 31 | 31 |
| 35-49 | 28 | 9.6 | 12 | 6 | 7 | 9 | 10 | 15 | 17 | 22 |
| 50-64 | 131 | 10.9 | 17 | 6 | 7 | 10 | 12 | 17 | 20 | 24 |
| 65+ | 97 | 11.3 | 29 | 6 | 7 | 10 | 13 | 19 | 22 | 34 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 15.1 | 75 | 7 | 7 | 14 | 19 | 27 | 34 | 34 |
| 20-34 | 17 | 16.2 | 120 | 7 | 8 | 13 | 20 | 26 | 43 | 52 |
| 35-49 | 80 | 14.3 | 89 | 7 | 7 | 11 | 18 | 22 | 29 | 62 |
| 50-64 | 311 | 14.6 | 71 | 7 | 7 | 12 | 18 | 25 | 36 | 77 |
| 65+ | 361 | 17.3 | 114 | 8 | 9 | 15 | 20 | 28 | 37 | 67 |
| **TOT. SINGLE DX** | 269 | 10.9 | 23 | 6 | 7 | 10 | 12 | 17 | 21 | 26 |
| **" MULTIPLE DX** | 779 | 15.9 | 96 | 7 | 8 | 13 | 19 | 26 | 35 | 67 |
| TOTAL 0-19 YRS | 13 | 13.4 | 63 | 5 | 6 | 11 | 19 | 24 | 32 | 34 |
| 20-34 | 27 | 14.5 | 99 | 6 | 6 | 11 | 16 | 26 | 34 | 52 |
| 35-49 | 108 | 13.1 | 73 | 6 | 7 | 10 | 17 | 21 | 25 | 54 |
| 50-64 | 442 | 13.5 | 58 | 6 | 7 | 11 | 16 | 23 | 29 | 54 |
| 65+ | 458 | 16.0 | 103 | 7 | 8 | 13 | 19 | 27 | 34 | 62 |
| **GRAND TOTAL** | **1048** | **14.6** | **82** | **7** | **7** | **12** | **18** | **25** | **32** | **57** |

## 567: Bronchoscopy (using natural orifice) (33.22-33.23)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 112 | 2.2 | 6 | <1 | <1 | 1 | 2 | 5 | 7 | 15 |
| 20-34 | 54 | 6.1 | 36 | <1 | 1 | 4 | 8 | 17 | 18 | 24 |
| 35-49 | 91 | 5.4 | 36 | <1 | 1 | 3 | 8 | 13 | 15 | 34 |
| 50-64 | 209 | 5.5 | 43 | <1 | <1 | 3 | 7 | 11 | 19 | 44 |
| 65+ | 230 | 5.5 | 31 | <1 | <1 | 3 | 8 | 13 | 17 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 132 | 6.7 | 143 | <1 | 1 | 3 | 9 | 16 | 19 | 84 |
| 20-34 | 149 | 10.6 | 109 | 1 | 1 | 8 | 14 | 24 | 30 | 50 |
| 35-49 | 245 | 10.5 | 145 | 1 | 2 | 8 | 12 | 23 | 31 | 81 |
| 50-64 | 659 | 10.2 | 113 | 1 | 1 | 7 | 13 | 22 | 30 | 66 |
| 65+ | 1172 | 13.6 | 161 | 1 | 2 | 10 | 18 | 29 | 37 | 67 |
| **TOT. SINGLE DX** | 696 | 5.0 | 34 | <1 | <1 | 3 | 7 | 12 | 16 | 32 |
| **" MULTIPLE DX** | 2357 | 11.8 | 146 | 1 | 2 | 8 | 16 | 26 | 34 | 68 |
| TOTAL 0-19 YRS | 244 | 4.6 | 85 | <1 | 1 | 2 | 4 | 11 | 17 | 34 |
| 20-34 | 203 | 9.4 | 94 | 1 | 1 | 7 | 14 | 22 | 27 | 44 |
| 35-49 | 336 | 9.1 | 121 | 1 | 1 | 6 | 11 | 19 | 28 | 76 |
| 50-64 | 868 | 9.0 | 101 | <1 | 1 | 7 | 11 | 20 | 28 | 64 |
| 65+ | 1402 | 12.3 | 149 | 1 | 2 | 9 | 17 | 27 | 35 | 63 |
| **GRAND TOTAL** | **3053** | **10.2** | **128** | **1** | **1** | **7** | **14** | **24** | **31** | **62** |

## 568: Miscellaneous operations on lung (32.6-32.9; 33.1; 33.3; 33.43; 33.49; 33.5; 33.93; 33.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 9 | 5.6 | 21 | 2 | 2 | 5 | 6 | 13 | 17 | 17 |
| 20-34 | 48 | 4.8 | 9 | 1 | 2 | 4 | 6 | 9 | 11 | 15 |
| 35-49 | 13 | 6.5 | 31 | 1 | 2 | 6 | 7 | 13 | 21 | 23 |
| 50-64 | 11 | 8.0 | 65 | 1 | 1 | 5 | 13 | 21 | 26 | 26 |
| 65+ | 11 | 11.4 | 229 | <1 | 1 | 7 | 10 | 36 | 51 | 52 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 20 | 8.6 | 60 | 1 | 2 | 7 | 11 | 18 | 27 | 32 |
| 20-34 | 40 | 9.5 | 142 | 1 | 2 | 6 | 11 | 17 | 25 | 73 |
| 35-49 | 25 | 6.7 | 27 | 2 | 2 | 5 | 10 | 13 | 41 | 99+ |
| 50-64 | 43 | 13.8 | 262 | 2 | 3 | 9 | 19 | 37 | 65 | 99+ |
| 65+ | 75 | 11.5 | 106 | 1 | 2 | 8 | 15 | 28 | 33 | 86 |
| **TOT. SINGLE DX** | 92 | 6.3 | 48 | 1 | 2 | 4 | 7 | 12 | 18 | 41 |
| **" MULTIPLE DX** | 203 | 10.7 | 135 | 2 | 2 | 7 | 13 | 26 | 34 | 99+ |
| TOTAL 0-19 YRS | 29 | 7.7 | 49 | 1 | 2 | 5 | 11 | 16 | 23 | 32 |
| 20-34 | 88 | 6.9 | 74 | 1 | 2 | 5 | 8 | 14 | 16 | 56 |
| 35-49 | 38 | 6.7 | 28 | 2 | 2 | 6 | 8 | 13 | 25 | 99+ |
| 50-64 | 54 | 12.6 | 226 | 1 | 2 | 7 | 16 | 34 | 59 | 99 |
| 65+ | 86 | 11.5 | 120 | 1 | 2 | 8 | 15 | 27 | 35 | 82 |
| **GRAND TOTAL** | **295** | **9.3** | **112** | **1** | **2** | **6** | **11** | **23** | **31** | **88** |

## 569: Thoracotomy (34.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 75 | 5.4 | 11 | 1 | 2 | 4 | 7 | 10 | 12 | 18 |
| 20-34 | 296 | 4.9 | 10 | 2 | 2 | 4 | 6 | 9 | 10 | 22 |
| 35-49 | 104 | 5.7 | 18 | 2 | 2 | 4 | 7 | 10 | 15 | 24 |
| 50-64 | 102 | 7.8 | 41 | 1 | 2 | 5 | 10 | 17 | 22 | 30 |
| 65+ | 68 | 9.8 | 50 | 2 | 3 | 8 | 13 | 18 | 23 | 38 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 203 | 13.3 | 282 | 2 | 3 | 8 | 15 | 28 | 64 | 85 |
| 20-34 | 383 | 8.4 | 76 | 2 | 3 | 6 | 9 | 17 | 24 | 51 |
| 35-49 | 309 | 10.2 | 78 | 2 | 3 | 8 | 13 | 18 | 26 | 50 |
| 50-64 | 517 | 12.6 | 112 | 3 | 4 | 9 | 16 | 27 | 34 | 48 |
| 65+ | 636 | 14.1 | 161 | 3 | 4 | 10 | 17 | 30 | 41 | 79 |
| **TOT. SINGLE DX** | 645 | 6.1 | 24 | 2 | 2 | 5 | 7 | 11 | 16 | 25 |
| **" MULTIPLE DX** | 2048 | 12.0 | 137 | 3 | 3 | 8 | 15 | 25 | 35 | 74 |
| TOTAL 0-19 YRS | 278 | 11.2 | 222 | 2 | 2 | 7 | 12 | 21 | 43 | 84 |
| 20-34 | 679 | 6.9 | 51 | 2 | 2 | 5 | 8 | 13 | 19 | 38 |
| 35-49 | 413 | 9.1 | 67 | 2 | 3 | 7 | 12 | 17 | 23 | 46 |
| 50-64 | 619 | 11.8 | 104 | 2 | 3 | 9 | 15 | 25 | 32 | 46 |
| 65+ | 704 | 13.7 | 153 | 3 | 4 | 10 | 17 | 29 | 39 | 77 |
| **GRAND TOTAL** | **2693** | **10.6** | **116** | **2** | **3** | **7** | **13** | **22** | **31** | **62** |

## 570: Operations on chest wall, pleura, and mediastinum except thoracotomy (34.1-34.27; 34.3-34.8; 34.93-34.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 75 | 5.7 | 14 | <1 | <1 | 5 | 7 | 11 | 14 | 18 |
| 20-34 | 81 | 6.5 | 23 | 1 | 1 | 6 | 9 | 13 | 17 | 22 |
| 35-49 | 63 | 4.8 | 19 | <1 | <1 | 3 | 7 | 9 | 17 | 19 |
| 50-64 | 116 | 5.0 | 19 | <1 | 1 | 3 | 7 | 11 | 14 | 22 |
| 65+ | 82 | 6.6 | 86 | 1 | 1 | 3 | 8 | 13 | 21 | 56 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 8.8 | 62 | 1 | 2 | 7 | 11 | 14 | 20 | 48 |
| 20-34 | 128 | 11.5 | 135 | 1 | 2 | 8 | 14 | 28 | 43 | 99+ |
| 35-49 | 151 | 8.8 | 88 | 1 | 2 | 6 | 10 | 21 | 25 | 58 |
| 50-64 | 325 | 9.8 | 77 | 1 | 2 | 8 | 14 | 22 | 27 | 51 |
| 65+ | 477 | 11.9 | 104 | 2 | 2 | 9 | 16 | 28 | 34 | 49 |
| **TOT. SINGLE DX** | 417 | 5.7 | 33 | <1 | 1 | 4 | 7 | 11 | 16 | 28 |
| **" MULTIPLE DX** | 1135 | 10.7 | 97 | 1 | 2 | 8 | 14 | 23 | 32 | 55 |
| TOTAL 0-19 YRS | 129 | 7.0 | 36 | <1 | 1 | 6 | 8 | 13 | 16 | 36 |
| 20-34 | 209 | 9.6 | 98 | 1 | 2 | 7 | 12 | 21 | 32 | 86 |
| 35-49 | 214 | 7.6 | 71 | 1 | 1 | 5 | 9 | 18 | 22 | 56 |
| 50-64 | 441 | 8.5 | 67 | 1 | 2 | 7 | 11 | 18 | 24 | 46 |
| 65+ | 559 | 11.2 | 106 | 1 | 2 | 8 | 15 | 26 | 34 | 49 |
| **GRAND TOTAL** | **1552** | **9.4** | **85** | **1** | **2** | **7** | **12** | **21** | **29** | **49** |

## 571: Open heart operations (35.1-35.51; 35.53-35.99; 37.5-37.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 92 | 8.1 | 18 | 6 | 6 | 8 | 9 | 10 | 11 | 31 |
| 20-34 | 42 | 10.3 | 18 | 5 | 7 | 9 | 12 | 14 | 19 | 29 |
| 35-49 | 48 | 10.4 | 14 | 6 | 7 | 10 | 12 | 15 | 16 | 21 |
| 50-64 | 81 | 12.2 | 23 | 7 | 8 | 11 | 13 | 17 | 19 | 36 |
| 65+ | 39 | 14.7 | 34 | 8 | 10 | 13 | 16 | 23 | 29 | 33 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 198 | 10.6 | 63 | 6 | 7 | 8 | 11 | 15 | 19 | 57 |
| 20-34 | 97 | 14.6 | 100 | 7 | 8 | 12 | 16 | 24 | 40 | 58 |
| 35-49 | 166 | 13.8 | 53 | 7 | 8 | 11 | 16 | 25 | 30 | 38 |
| 50-64 | 462 | 16.8 | 98 | 8 | 9 | 14 | 19 | 27 | 35 | 64 |
| 65+ | 502 | 19.0 | 127 | 9 | 10 | 16 | 22 | 32 | 39 | 67 |
| **TOT. SINGLE DX** | 302 | 10.7 | 26 | 6 | 7 | 9 | 12 | 16 | 19 | 31 |
| **" MULTIPLE DX** | 1425 | 16.2 | 106 | 7 | 8 | 13 | 19 | 28 | 35 | 63 |
| TOTAL 0-19 YRS | 290 | 9.8 | 50 | 6 | 7 | 8 | 10 | 14 | 17 | 49 |
| 20-34 | 139 | 13.3 | 80 | 7 | 7 | 10 | 14 | 21 | 29 | 54 |
| 35-49 | 214 | 13.1 | 46 | 7 | 7 | 11 | 15 | 23 | 29 | 36 |
| 50-64 | 543 | 16.1 | 90 | 8 | 9 | 13 | 18 | 27 | 34 | 63 |
| 65+ | 541 | 18.6 | 122 | 9 | 10 | 15 | 22 | 31 | 37 | 66 |
| **GRAND TOTAL** | **1727** | **15.2** | **97** | **7** | **8** | **12** | **18** | **27** | **33** | **62** |

## 572: Direct heart revascularization (36.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 10.0 | | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 20-34 | 4 | 9.8 | 4 | 7 | 7 | 10 | 11 | 12 | 12 | 12 |
| 35-49 | 159 | 9.6 | 7 | 6 | 7 | 9 | 11 | 13 | 14 | 20 |
| 50-64 | 547 | 10.6 | 14 | 7 | 8 | 10 | 12 | 14 | 16 | 24 |
| 65+ | 256 | 12.1 | 21 | 7 | 8 | 11 | 14 | 17 | 21 | 32 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 13.4 | 15 | 9 | 9 | 14 | 16 | 18 | 18 | 18 |
| 20-34 | 35 | 13.8 | 159 | 7 | 8 | 10 | 13 | 19 | 34 | 78 |
| 35-49 | 917 | 11.6 | 24 | 7 | 7 | 10 | 13 | 17 | 21 | 33 |
| 50-64 | 3488 | 12.8 | 42 | 7 | 8 | 11 | 14 | 19 | 24 | 40 |
| 65+ | 2165 | 15.9 | 87 | 8 | 9 | 13 | 18 | 25 | 32 | 65 |
| **TOT. SINGLE DX** | 967 | 10.8 | 16 | 7 | 8 | 10 | 12 | 15 | 17 | 27 |
| **" MULTIPLE DX** | 6610 | 13.6 | 58 | 7 | 8 | 12 | 15 | 21 | 27 | 47 |
| TOTAL 0-19 YRS | 6 | 12.8 | 14 | 9 | 9 | 12 | 16 | 18 | 18 | 18 |
| 20-34 | 39 | 13.4 | 145 | 7 | 8 | 10 | 13 | 18 | 32 | 78 |
| 35-49 | 1076 | 11.3 | 23 | 7 | 7 | 10 | 13 | 17 | 20 | 32 |
| 50-64 | 4035 | 12.5 | 39 | 7 | 8 | 11 | 14 | 19 | 23 | 39 |
| 65+ | 2421 | 15.5 | 82 | 8 | 9 | 13 | 17 | 24 | 31 | 61 |
| **GRAND TOTAL** | **7577** | **13.3** | **53** | **7** | **8** | **11** | **15** | **20** | **26** | **44** |

## 573: Cardiac catheterization (37.21-37.23)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 247 | 1.6 | 1 | <1 | 1 | 1 | 2 | 2 | 3 | 8 |
| 20-34 | 155 | 2.3 | 2 | 1 | 1 | 2 | 2 | 4 | 6 | 8 |
| 35-49 | 902 | 2.6 | 4 | 1 | 1 | 2 | 2 | 5 | 8 | 12 |
| 50-64 | 1988 | 2.6 | 5 | 1 | 1 | 2 | 2 | 5 | 7 | 14 |
| 65+ | 816 | 3.1 | 10 | 1 | 1 | 2 | 3 | 7 | 9 | 19 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 320 | 2.9 | 39 | <1 | 1 | 1 | 2 | 5 | 10 | 27 |
| 20-34 | 315 | 4.0 | 17 | 1 | 1 | 2 | 5 | 10 | 12 | 18 |
| 35-49 | 2642 | 3.7 | 17 | 1 | 1 | 2 | 4 | 8 | 11 | 19 |
| 50-64 | 7320 | 4.0 | 22 | 1 | 1 | 2 | 4 | 9 | 13 | 23 |
| 65+ | 4492 | 5.4 | 43 | 1 | 2 | 2 | 7 | 13 | 17 | 33 |
| **TOT. SINGLE DX** | 4108 | 2.6 | 6 | 1 | 1 | 2 | 2 | 5 | 8 | 14 |
| **" MULTIPLE DX** | 15089 | 4.4 | 29 | 1 | 1 | 2 | 5 | 10 | 14 | 26 |
| TOTAL 0-19 YRS | 567 | 2.3 | 23 | <1 | 1 | 1 | 2 | 3 | 7 | 15 |
| 20-34 | 470 | 3.5 | 13 | 1 | 1 | 2 | 4 | 8 | 11 | 17 |
| 35-49 | 3544 | 3.4 | 15 | 1 | 1 | 2 | 3 | 8 | 10 | 18 |
| 50-64 | 9308 | 3.7 | 19 | 1 | 1 | 2 | 4 | 8 | 12 | 22 |
| 65+ | 5308 | 5.1 | 39 | 1 | 1 | 2 | 6 | 12 | 16 | 31 |
| **GRAND TOTAL** | **19197** | **4.0** | **24** | **1** | **1** | **2** | **4** | **9** | **13** | **24** |

## 574: Pacemaker insertion, replacement, removal, and repair (37.7-37.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 2.8 | 3 | 1 | 1 | 2 | 3 | 6 | 6 | 6 |
| 20-34 | 13 | 4.0 | 7 | 1 | 1 | 3 | 5 | 9 | 10 | 10 |
| 35-49 | 40 | 4.1 | 9 | 1 | 1 | 3 | 5 | 8 | 11 | 14 |
| 50-64 | 146 | 4.9 | 19 | 1 | 1 | 3 | 7 | 10 | 11 | 20 |
| 65+ | 610 | 4.6 | 15 | 1 | 1 | 3 | 6 | 9 | 12 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 3.3 | 6 | 1 | 1 | 3 | 4 | 6 | 10 | 11 |
| 20-34 | 43 | 6.2 | 27 | 2 | 2 | 4 | 8 | 12 | 15 | 30 |
| 35-49 | 151 | 9.2 | 44 | 2 | 2 | 8 | 12 | 18 | 22 | 25 |
| 50-64 | 721 | 9.5 | 57 | 2 | 2 | 8 | 12 | 17 | 24 | 35 |
| 65+ | 4127 | 9.8 | 67 | 2 | 3 | 8 | 13 | 19 | 25 | 42 |
| **TOT. SINGLE DX** | 815 | 4.6 | 15 | 1 | 1 | 3 | 6 | 9 | 12 | 18 |
| **" MULTIPLE DX** | 5056 | 9.7 | 65 | 2 | 3 | 8 | 13 | 19 | 24 | 40 |
| TOTAL 0-19 YRS | 20 | 3.2 | 6 | 1 | 1 | 3 | 4 | 6 | 9 | 11 |
| 20-34 | 56 | 5.7 | 24 | 1 | 2 | 4 | 8 | 12 | 13 | 29 |
| 35-49 | 191 | 8.1 | 42 | 1 | 2 | 6 | 11 | 16 | 21 | 25 |
| 50-64 | 867 | 8.7 | 54 | 1 | 2 | 7 | 12 | 17 | 23 | 35 |
| 65+ | 4737 | 9.1 | 63 | 2 | 2 | 7 | 12 | 18 | 24 | 40 |
| **GRAND TOTAL** | **5871** | **9.0** | **61** | **2** | **2** | **7** | **12** | **18** | **24** | **39** |

## 575: Miscellaneous operations on heart (35.0; 35.52; 36.0; 36.2-36.9; 37.0-37.1; 37.24-37.25; 37.3-37.4; 37.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 6.6 | 1 | 4 | 5 | 7 | 7 | 8 | 8 | 8 |
| 20-34 | 7 | 8.0 | 59 | 1 | 1 | 6 | 9 | 21 | 24 | 24 |
| 35-49 | 33 | 6.4 | 21 | 1 | 2 | 7 | 8 | 11 | 16 | 22 |
| 50-64 | 53 | 4.9 | 22 | 1 | 2 | 3 | 5 | 13 | 16 | 23 |
| 65+ | 20 | 5.2 | 21 | <1 | 1 | 3 | 4 | 14 | 15 | 16 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 22 | 15.6 | 258 | 5 | 6 | 9 | 18 | 32 | 60 | 70 |
| 20-34 | 44 | 11.5 | 103 | 2 | 3 | 8 | 15 | 19 | 31 | 58 |
| 35-49 | 131 | 10.0 | 96 | 2 | 2 | 7 | 13 | 23 | 33 | 47 |
| 50-64 | 288 | 9.1 | 82 | 2 | 2 | 7 | 12 | 19 | 28 | 47 |
| 65+ | 179 | 14.2 | 128 | 2 | 3 | 11 | 20 | 31 | 39 | 88 |
| **TOT. SINGLE DX** | 123 | 5.7 | 22 | 1 | 2 | 4 | 7 | 13 | 16 | 23 |
| **" MULTIPLE DX** | 664 | 11.0 | 109 | 2 | 2 | 8 | 14 | 24 | 35 | 59 |
| TOTAL 0-19 YRS | 32 | 12.8 | 193 | 4 | 5 | 7 | 14 | 21 | 50 | 70 |
| 20-34 | 51 | 11.1 | 98 | 2 | 2 | 8 | 15 | 19 | 28 | 58 |
| 35-49 | 164 | 9.3 | 84 | 1 | 2 | 7 | 12 | 21 | 27 | 46 |
| 50-64 | 341 | 8.5 | 75 | 2 | 2 | 6 | 11 | 17 | 25 | 46 |
| 65+ | 199 | 13.3 | 125 | 2 | 3 | 11 | 19 | 30 | 37 | 81 |
| **GRAND TOTAL** | **787** | **10.2** | **100** | **2** | **2** | **7** | **13** | **22** | **32** | **56** |

## 576: Other diagnostic and therapeutic procedures on larynx, trachea, bronchus, lungs and heart (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 160 | 3.0 | 11 | <1 | <1 | 1 | 4 | 7 | 9 | 19 |
| 20-34 | 94 | 3.5 | 11 | <1 | <1 | 2 | 5 | 8 | 10 | 16 |
| 35-49 | 130 | 3.3 | 27 | <1 | <1 | 1 | 3 | 9 | 10 | 35 |
| 50-64 | 183 | 3.0 | 14 | <1 | <1 | 2 | 3 | 7 | 11 | 18 |
| 65+ | 168 | 4.7 | 27 | <1 | 1 | 2 | 6 | 12 | 17 | 23 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 227 | 6.6 | 78 | <1 | 1 | 4 | 8 | 16 | 21 | 39 |
| 20-34 | 163 | 9.5 | 105 | 1 | 1 | 7 | 11 | 20 | 35 | 49 |
| 35-49 | 256 | 8.9 | 96 | 1 | 1 | 6 | 12 | 19 | 27 | 61 |
| 50-64 | 679 | 9.5 | 85 | 1 | 2 | 7 | 12 | 21 | 28 | 44 |
| 65+ | 1484 | 11.2 | 99 | 1 | 2 | 9 | 15 | 23 | 30 | 52 |
| **TOT. SINGLE DX** | 735 | 3.5 | 19 | <1 | <1 | 2 | 4 | 8 | 12 | 21 |
| **" MULTIPLE DX** | 2809 | 10.1 | 96 | 1 | 2 | 7 | 13 | 22 | 29 | 50 |
| TOTAL 0-19 YRS | 387 | 5.1 | 54 | <1 | <1 | 3 | 6 | 12 | 16 | 35 |
| 20-34 | 257 | 7.3 | 79 | 1 | 1 | 4 | 9 | 17 | 23 | 42 |
| 35-49 | 386 | 7.0 | 80 | <1 | 1 | 4 | 10 | 17 | 22 | 53 |
| 50-64 | 862 | 8.1 | 78 | <1 | 1 | 5 | 11 | 19 | 27 | 42 |
| 65+ | 1652 | 10.5 | 96 | 1 | 2 | 8 | 14 | 22 | 29 | 50 |
| **GRAND TOTAL** | **3544** | **8.7** | **87** | **1** | **1** | **5** | **12** | **20** | **27** | **46** |

## 577: Ligation and stripping of varicose veins (38.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 11 | 2.7 | 1 | 1 | 1 | 3 | 4 | 4 | 4 | 4 |
| 20-34 | 432 | 3.2 | 2 | 1 | 1 | 3 | 4 | 5 | 5 | 10 |
| 35-49 | 562 | 3.4 | 3 | 1 | 2 | 3 | 4 | 5 | 7 | 11 |
| 50-64 | 391 | 3.7 | 4 | 1 | 2 | 3 | 4 | 6 | 7 | 13 |
| 65+ | 138 | 4.5 | 10 | 1 | 2 | 4 | 5 | 8 | 9 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 5.7 | 2 | 4 | 4 | 6 | 7 | 7 | 7 | 7 |
| 20-34 | 98 | 4.4 | 20 | 2 | 2 | 4 | 5 | 7 | 9 | 29 |
| 35-49 | 171 | 4.3 | 8 | 2 | 2 | 4 | 5 | 7 | 10 | 14 |
| 50-64 | 224 | 5.1 | 20 | 2 | 2 | 4 | 6 | 10 | 13 | 24 |
| 65+ | 124 | 8.2 | 102 | 2 | 2 | 5 | 8 | 17 | 25 | 63 |
| **TOT. SINGLE DX** | 1534 | 3.5 | 4 | 1 | 2 | 3 | 4 | 5 | 7 | 12 |
| **" MULTIPLE DX** | 620 | 5.4 | 35 | 2 | 2 | 4 | 6 | 10 | 15 | 34 |
| TOTAL 0-19 YRS | 14 | 3.4 | 3 | 1 | 1 | 3 | 4 | 6 | 7 | 7 |
| 20-34 | 530 | 3.4 | 6 | 1 | 2 | 3 | 4 | 5 | 6 | 12 |
| 35-49 | 733 | 3.6 | 5 | 1 | 2 | 3 | 4 | 6 | 8 | 12 |
| 50-64 | 615 | 4.2 | 11 | 1 | 2 | 3 | 5 | 7 | 10 | 18 |
| 65+ | 262 | 6.2 | 57 | 1 | 2 | 4 | 6 | 12 | 18 | 35 |
| **GRAND TOTAL** | **2154** | **4.0** | **14** | **1** | **2** | **3** | **4** | **6** | **9** | **19** |

## 578: Systemic shunt or graft bypass (39.0-39.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 15 | 12.5 | 470 | 2 | 3 | 7 | 9 | 13 | 71 | 90 |
| 20-34 | 24 | 4.3 | 21 | <1 | 1 | 2 | 7 | 8 | 12 | 22 |
| 35-49 | 91 | 6.5 | 13 | 1 | 2 | 7 | 9 | 11 | 12 | 17 |
| 50-64 | 244 | 7.7 | 15 | 2 | 2 | 8 | 10 | 12 | 14 | 24 |
| 65+ | 280 | 9.4 | 43 | 2 | 3 | 8 | 12 | 16 | 20 | 33 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 45 | 12.5 | 154 | 2 | 3 | 8 | 14 | 31 | 44 | 53 |
| 20-34 | 127 | 11.5 | 124 | 2 | 2 | 8 | 15 | 24 | 39 | 68 |
| 35-49 | 327 | 12.2 | 120 | 1 | 2 | 10 | 15 | 23 | 30 | 74 |
| 50-64 | 1149 | 12.7 | 87 | 3 | 5 | 10 | 15 | 23 | 32 | 61 |
| 65+ | 1619 | 16.0 | 141 | 5 | 6 | 12 | 19 | 32 | 42 | 67 |
| **TOT. SINGLE DX** | 654 | 8.2 | 39 | 1 | 2 | 8 | 10 | 13 | 17 | 25 |
| **" MULTIPLE DX** | 3267 | 14.2 | 123 | 3 | 5 | 11 | 17 | 28 | 38 | 64 |
| TOTAL 0-19 YRS | 60 | 12.5 | 227 | 2 | 3 | 8 | 12 | 30 | 46 | 86 |
| 20-34 | 151 | 10.4 | 114 | 1 | 2 | 7 | 13 | 22 | 37 | 59 |
| 35-49 | 418 | 11.0 | 103 | 1 | 2 | 9 | 13 | 22 | 28 | 67 |
| 50-64 | 1393 | 11.8 | 79 | 3 | 4 | 10 | 14 | 22 | 29 | 54 |
| 65+ | 1899 | 15.0 | 133 | 4 | 6 | 12 | 18 | 30 | 40 | 65 |
| **GRAND TOTAL** | **3921** | **13.2** | **114** | **2** | **4** | **10** | **16** | **26** | **36** | **63** |

## 579: Extracorporeal circulation (39.61)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | 1 | 13.0 | | 13 | 13 | 13 | 13 | 13 | 13 | 13 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 10.0 | | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 20-34 | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 35-49 | 5 | 15.8 | 128 | 7 | 7 | 10 | 21 | 35 | 35 | 35 |
| 50-64 | 3 | 10.0 | 81 | 1 | 1 | 10 | 17 | 19 | 19 | 19 |
| 65+ | 2 | 18.0 | 18 | 15 | 15 | 18 | 21 | 21 | 21 | 21 |
| **TOT. SINGLE DX** | 1 | 13.0 | | 13 | 13 | 13 | 13 | 13 | 13 | 13 |
| **" MULTIPLE DX** | 13 | 12.9 | 77 | 1 | 3 | 10 | 17 | 24 | 33 | 35 |
| TOTAL 0-19 YRS | 2 | 10.0 | | 10 | 10 | 10 | 10 | 10 | 10 | 10 |
| 20-34 | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 35-49 | 5 | 15.8 | 129 | 7 | 7 | 10 | 21 | 35 | 35 | 35 |
| 50-64 | 4 | 10.8 | 56 | 1 | 1 | 11 | 16 | 19 | 19 | 19 |
| 65+ | 2 | 18.0 | 18 | 15 | 15 | 18 | 21 | 21 | 21 | 21 |
| **GRAND TOTAL** | **14** | **12.9** | **71** | **1** | **3** | **10** | **17** | **22** | **32** | **35** |

## 580: Hemodialysis (39.95)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 5 | 5 |
| 20-34 | 8 | 2.8 | 4 | 1 | 1 | 2 | 3 | 6 | 7 | 7 |
| 35-49 | 21 | 2.3 | 4 | <1 | <1 | 1 | 3 | 6 | 7 | 8 |
| 50-64 | 15 | 3.9 | 10 | <1 | <1 | 4 | 5 | 7 | 11 | 13 |
| 65+ | 11 | 5.6 | 5 | <1 | 2 | 6 | 6 | 9 | 10 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 45 | 4.4 | 13 | 1 | 1 | 4 | 5 | 8 | 15 | 16 |
| 20-34 | 350 | 6.2 | 59 | <1 | 1 | 4 | 7 | 12 | 19 | 32 |
| 35-49 | 349 | 5.7 | 34 | <1 | 1 | 4 | 7 | 12 | 16 | 34 |
| 50-64 | 709 | 6.4 | 47 | <1 | 1 | 4 | 8 | 14 | 18 | 39 |
| 65+ | 682 | 8.2 | 73 | 1 | 2 | 5 | 10 | 17 | 26 | 51 |
| **TOT. SINGLE DX** | 60 | 3.4 | 7 | <1 | <1 | 3 | 5 | 7 | 8 | 13 |
| **" MULTIPLE DX** | 2135 | 6.8 | 56 | 1 | 1 | 5 | 8 | 14 | 20 | 43 |
| TOTAL 0-19 YRS | 50 | 4.2 | 12 | 1 | 1 | 3 | 5 | 8 | 15 | 16 |
| 20-34 | 358 | 6.1 | 59 | <1 | 1 | 4 | 7 | 12 | 18 | 32 |
| 35-49 | 370 | 5.5 | 33 | <1 | 1 | 4 | 7 | 11 | 16 | 34 |
| 50-64 | 724 | 6.3 | 47 | <1 | 1 | 4 | 8 | 13 | 18 | 39 |
| 65+ | 693 | 8.1 | 73 | 1 | 2 | 5 | 10 | 17 | 25 | 51 |
| **GRAND TOTAL** | **2195** | **6.7** | **55** | **1** | **1** | **4** | **8** | **14** | **20** | **43** |

## 581: Other operations on blood vessels (38.0-38.21; 38.3-38.4; 38.6-38.92; 38.94; 38.98; 39.3-39.5; 39.62-39.91; 39.93-39.94; 39.96-39.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 181 | 7.7 | 122 | <1 | 1 | 6 | 8 | 12 | 21 | 79 |
| 20-34 | 169 | 5.1 | 31 | 1 | 1 | 3 | 7 | 11 | 18 | 23 |
| 35-49 | 256 | 7.2 | 50 | 1 | 1 | 5 | 9 | 15 | 22 | 32 |
| 50-64 | 688 | 6.3 | 28 | 1 | 2 | 5 | 8 | 11 | 13 | 27 |
| 65+ | 1078 | 7.2 | 28 | 2 | 2 | 6 | 9 | 13 | 16 | 28 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2303 | 15.4 | 267 | 2 | 3 | 9 | 20 | 41 | 58 | 99+ |
| 20-34 | 605 | 8.7 | 117 | 1 | 1 | 5 | 10 | 20 | 32 | 64 |
| 35-49 | 821 | 10.6 | 155 | 1 | 2 | 7 | 12 | 24 | 36 | 64 |
| 50-64 | 2863 | 10.5 | 109 | 2 | 2 | 8 | 13 | 22 | 32 | 57 |
| 65+ | 5292 | 12.4 | 127 | 2 | 3 | 9 | 15 | 24 | 35 | 66 |
| **TOT. SINGLE DX** | 2372 | 6.8 | 39 | 1 | 2 | 5 | 9 | 12 | 16 | 31 |
| **" MULTIPLE DX** | 11884 | 12.2 | 155 | 2 | 3 | 9 | 15 | 26 | 39 | 75 |
| TOTAL 0-19 YRS | 2484 | 14.8 | 261 | 2 | 3 | 9 | 19 | 40 | 56 | 99+ |
| 20-34 | 774 | 7.9 | 101 | 1 | 1 | 5 | 9 | 18 | 27 | 63 |
| 35-49 | 1077 | 9.8 | 133 | 1 | 1 | 6 | 11 | 22 | 32 | 58 |
| 50-64 | 3551 | 9.7 | 97 | 2 | 2 | 7 | 12 | 20 | 29 | 53 |
| 65+ | 6370 | 11.5 | 114 | 2 | 3 | 9 | 14 | 23 | 32 | 62 |
| **GRAND TOTAL** | **14256** | **11.3** | **140** | **2** | **2** | **8** | **14** | **24** | **36** | **71** |

## 582: Incision, excision of lymphatic structures (40.0-40.11; 40.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 230 | 2.6 | 8 | <1 | <1 | 1 | 3 | 6 | 7 | 16 |
| 20-34 | 277 | 2.9 | 6 | <1 | <1 | 2 | 3 | 6 | 9 | 13 |
| 35-49 | 199 | 3.1 | 10 | <1 | <1 | 2 | 4 | 7 | 9 | 17 |
| 50-64 | 251 | 3.8 | 17 | <1 | <1 | 2 | 4 | 9 | 11 | 24 |
| 65+ | 190 | 4.9 | 24 | <1 | <1 | 3 | 6 | 11 | 15 | 22 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 120 | 3.8 | 11 | <1 | 1 | 2 | 5 | 8 | 11 | 16 |
| 20-34 | 169 | 6.6 | 44 | <1 | 1 | 4 | 9 | 16 | 19 | 32 |
| 35-49 | 234 | 6.4 | 49 | <1 | 1 | 4 | 9 | 14 | 19 | 35 |
| 50-64 | 486 | 7.4 | 72 | 1 | 1 | 5 | 9 | 15 | 22 | 42 |
| 65+ | 607 | 9.4 | 112 | 1 | 2 | 6 | 12 | 22 | 29 | 54 |
| **TOT. SINGLE DX** | 1147 | 3.4 | 13 | <1 | <1 | 2 | 4 | 8 | 10 | 19 |
| **" MULTIPLE DX** | 1616 | 7.7 | 79 | 1 | 1 | 5 | 10 | 17 | 24 | 45 |
| TOTAL 0-19 YRS | 350 | 3.0 | 10 | <1 | <1 | 2 | 4 | 7 | 9 | 16 |
| 20-34 | 446 | 4.3 | 24 | <1 | <1 | 2 | 5 | 10 | 14 | 26 |
| 35-49 | 433 | 4.9 | 34 | <1 | <1 | 3 | 7 | 11 | 15 | 33 |
| 50-64 | 737 | 6.2 | 57 | <1 | 1 | 4 | 8 | 13 | 19 | 33 |
| 65+ | 797 | 8.3 | 95 | 1 | 1 | 5 | 10 | 20 | 27 | 51 |
| **GRAND TOTAL** | **2763** | **5.9** | **56** | **<1** | **1** | **3** | **7** | **13** | **20** | **34** |

## 583: Radical excision of lymph nodes (40.3-40.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 9 | 2.8 | 4 | <1 | 1 | 2 | 3 | 6 | 8 | 8 |
| 20-34 | 40 | 6.2 | 13 | 1 | 1 | 6 | 9 | 11 | 12 | 15 |
| 35-49 | 28 | 5.8 | 8 | 1 | 2 | 6 | 8 | 9 | 10 | 15 |
| 50-64 | 48 | 6.7 | 33 | 1 | 2 | 6 | 7 | 10 | 17 | 36 |
| 65+ | 37 | 7.3 | 47 | 2 | 2 | 6 | 8 | 12 | 21 | 40 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 11 | 9.3 | 31 | 1 | 2 | 9 | 11 | 16 | 22 | 23 |
| 20-34 | 84 | 7.8 | 10 | 2 | 3 | 9 | 9 | 11 | 14 | 16 |
| 35-49 | 59 | 8.1 | 35 | 1 | 3 | 6 | 11 | 16 | 22 | 26 |
| 50-64 | 134 | 8.5 | 30 | 3 | 4 | 7 | 10 | 15 | 20 | 29 |
| 65+ | 175 | 9.7 | 89 | 3 | 4 | 7 | 11 | 19 | 23 | 44 |
| **TOT. SINGLE DX** | 162 | 6.3 | 26 | 1 | 2 | 6 | 8 | 10 | 14 | 35 |
| **" MULTIPLE DX** | 463 | 8.8 | 50 | 3 | 4 | 7 | 10 | 15 | 21 | 32 |
| TOTAL 0-19 YRS | 20 | 6.4 | 29 | <1 | 1 | 5 | 9 | 11 | 17 | 23 |
| 20-34 | 124 | 7.3 | 12 | 1 | 2 | 8 | 9 | 11 | 13 | 16 |
| 35-49 | 87 | 7.3 | 28 | 1 | 2 | 6 | 9 | 14 | 20 | 26 |
| 50-64 | 182 | 8.0 | 32 | 2 | 3 | 7 | 9 | 15 | 19 | 33 |
| 65+ | 212 | 9.3 | 83 | 3 | 3 | 7 | 10 | 18 | 23 | 43 |
| **GRAND TOTAL** | **625** | **8.2** | **45** | **2** | **3** | **7** | **9** | **15** | **20** | **33** |

## 584: Miscellaneous operations on lymphatic system (40.6-40.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 1 | 19.0 | | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| 35-49 | | | | | | | | | | |
| 50-64 | 4 | 6.3 | 31 | 1 | 1 | 6 | 11 | 12 | 12 | 12 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | 2 | 8.0 | 18 | 5 | 5 | 8 | 11 | 11 | 11 | 11 |
| 65+ | 1 | 4.0 | | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| **TOT. SINGLE DX** | 5 | 8.8 | 56 | 1 | 1 | 10 | 14 | 19 | 19 | 19 |
| **" MULTIPLE DX** | 3 | 6.7 | 14 | 4 | 4 | 5 | 9 | 11 | 11 | 11 |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | 1 | 19.0 | | 19 | 19 | 19 | 19 | 19 | 19 | 19 |
| 35-49 | | | | | | | | | | |
| 50-64 | 6 | 6.8 | 23 | 1 | 1 | 7 | 11 | 12 | 12 | 12 |
| 65+ | 1 | 4.0 | | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| **GRAND TOTAL** | **8** | **8.0** | **37** | **1** | **1** | **7** | **11** | **17** | **19** | **19** |

## 585: Bone marrow transplant (41.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 1 | 31.0 | | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 41.8 | 42 | 33 | 33 | 42 | 46 | 50 | 51 | 51 |
| 20-34 | 2 | 39.0 | 162 | 30 | 30 | 39 | 48 | 48 | 48 | 48 |
| 35-49 | | | | | | | | | | |
| 50-64 | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 1 | 31.0 | | 31 | 31 | 31 | 31 | 31 | 31 | 31 |
| **" MULTIPLE DX** | 9 | 37.1 | 193 | 5 | 15 | 40 | 46 | 50 | 51 | 51 |
| TOTAL 0-19 YRS | 6 | 41.8 | 42 | 33 | 33 | 42 | 46 | 50 | 51 | 51 |
| 20-34 | 3 | 36.3 | 102 | 30 | 30 | 31 | 44 | 48 | 48 | 48 |
| 35-49 | | | | | | | | | | |
| 50-64 | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **10** | **36.5** | **175** | **5** | **17** | **38** | **46** | **49** | **51** | **51** |

## 586: Miscellaneous operations on spleen and bone marrow (41.1-41.33; 41.4-41.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 209 | 6.5 | 57 | 1 | 1 | 5 | 7 | 13 | 19 | 39 |
| 20-34 | 181 | 6.7 | 36 | 1 | 2 | 6 | 8 | 13 | 19 | 35 |
| 35-49 | 123 | 8.8 | 95 | 1 | 2 | 6 | 9 | 18 | 24 | 59 |
| 50-64 | 124 | 7.6 | 49 | 1 | 2 | 6 | 10 | 15 | 22 | 36 |
| 65+ | 203 | 8.1 | 48 | 1 | 2 | 6 | 10 | 17 | 24 | 34 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 306 | 9.5 | 98 | 2 | 3 | 7 | 10 | 20 | 30 | 50 |
| 20-34 | 485 | 11.7 | 149 | 2 | 3 | 8 | 13 | 23 | 37 | 81 |
| 35-49 | 436 | 11.8 | 126 | 2 | 3 | 8 | 14 | 27 | 33 | 58 |
| 50-64 | 904 | 11.5 | 111 | 2 | 3 | 8 | 14 | 23 | 32 | 50 |
| 65+ | 2215 | 12.3 | 113 | 3 | 4 | 9 | 15 | 25 | 33 | 59 |
| **TOT. SINGLE DX** | 840 | 7.4 | 56 | 1 | 2 | 6 | 9 | 15 | 22 | 36 |
| **" MULTIPLE DX** | 4346 | 11.8 | 118 | 2 | 3 | 9 | 14 | 24 | 32 | 60 |
| TOTAL 0-19 YRS | 515 | 8.3 | 84 | 1 | 2 | 6 | 9 | 16 | 27 | 46 |
| 20-34 | 666 | 10.4 | 124 | 2 | 2 | 7 | 12 | 21 | 31 | 71 |
| 35-49 | 559 | 11.2 | 121 | 2 | 3 | 8 | 13 | 25 | 32 | 58 |
| 50-64 | 1028 | 11.0 | 106 | 2 | 3 | 8 | 13 | 22 | 31 | 49 |
| 65+ | 2418 | 11.9 | 109 | 2 | 3 | 9 | 14 | 24 | 31 | 58 |
| **GRAND TOTAL** | **5186** | **11.1** | **110** | **2** | **3** | **8** | **13** | **23** | **31** | **58** |

## 587: Other diagnostic and therapeutic procedures on vessels, lymphatic structures, and spleen (38.29; 38.93; 38.99; 39.92; 40.19; 41.38-41.39)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 54 | 6.3 | 65 | 1 | 1 | 3 | 7 | 13 | 26 | 44 |
| 20-34 | 38 | 10.7 | 199 | <1 | 1 | 5 | 13 | 30 | 41 | 64 |
| 35-49 | 31 | 8.6 | 84 | 1 | 1 | 4 | 12 | 25 | 30 | 31 |
| 50-64 | 27 | 10.4 | 129 | 1 | 1 | 4 | 14 | 29 | 35 | 44 |
| 65+ | 35 | 9.3 | 63 | 2 | 3 | 7 | 13 | 19 | 30 | 34 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 125 | 12.6 | 301 | 1 | 1 | 6 | 17 | 35 | 55 | 99+ |
| 20-34 | 119 | 13.7 | 174 | 1 | 2 | 9 | 25 | 35 | 44 | 99+ |
| 35-49 | 160 | 14.4 | 251 | 2 | 2 | 8 | 18 | 40 | 55 | 66 |
| 50-64 | 292 | 14.4 | 212 | 1 | 2 | 9 | 19 | 35 | 45 | 86 |
| 65+ | 297 | 18.0 | 275 | 2 | 3 | 13 | 24 | 38 | 49 | 85 |
| **TOT. SINGLE DX** | 185 | 8.8 | 106 | 1 | 1 | 4 | 12 | 25 | 30 | 45 |
| **" MULTIPLE DX** | 993 | 15.1 | 247 | 1 | 2 | 10 | 21 | 36 | 48 | 93 |
| TOTAL 0-19 YRS | 179 | 10.7 | 238 | 1 | 1 | 5 | 12 | 28 | 49 | 98 |
| 20-34 | 157 | 13.0 | 181 | 1 | 2 | 8 | 20 | 35 | 43 | 99+ |
| 35-49 | 191 | 13.4 | 229 | 1 | 2 | 8 | 16 | 35 | 52 | 66 |
| 50-64 | 319 | 14.0 | 206 | 1 | 2 | 9 | 19 | 34 | 43 | 81 |
| 65+ | 332 | 17.1 | 260 | 2 | 3 | 12 | 23 | 36 | 48 | 84 |
| **GRAND TOTAL** | **1178** | **14.1** | **230** | **1** | **2** | **9** | **19** | **35** | **46** | **90** |

## 588: Operations on the esophagus (42.0-42.21; 42.24; 42.3-42.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 27 | 3.7 | 13 | <1 | <1 | 2 | 4 | 11 | 12 | 13 |
| 20-34 | 50 | 3.6 | 12 | <1 | <1 | 2 | 5 | 9 | 11 | 16 |
| 35-49 | 58 | 4.4 | 23 | <1 | <1 | 2 | 5 | 10 | 15 | 23 |
| 50-64 | 99 | 6.0 | 45 | 1 | 1 | 4 | 8 | 14 | 17 | 35 |
| 65+ | 139 | 4.9 | 42 | <1 | <1 | 3 | 6 | 11 | 15 | 29 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 39 | 9.1 | 165 | 1 | 1 | 3 | 8 | 28 | 44 | 50 |
| 20-34 | 92 | 8.6 | 156 | <1 | 1 | 5 | 9 | 18 | 29 | 93 |
| 35-49 | 183 | 9.8 | 119 | 1 | 2 | 6 | 11 | 21 | 29 | 60 |
| 50-64 | 407 | 9.1 | 98 | 1 | 2 | 6 | 12 | 19 | 25 | 55 |
| 65+ | 792 | 10.9 | 134 | 1 | 2 | 7 | 13 | 23 | 31 | 65 |
| **TOT. SINGLE DX** | 373 | 4.9 | 35 | <1 | <1 | 3 | 6 | 11 | 15 | 28 |
| **" MULTIPLE DX** | 1513 | 10.1 | 126 | 1 | 2 | 7 | 12 | 22 | 30 | 64 |
| TOTAL 0-19 YRS | 66 | 6.9 | 109 | <1 | 1 | 3 | 7 | 18 | 33 | 50 |
| 20-34 | 142 | 6.8 | 111 | <1 | 1 | 4 | 7 | 15 | 21 | 84 |
| 35-49 | 241 | 8.5 | 101 | <1 | 1 | 5 | 10 | 20 | 27 | 56 |
| 50-64 | 506 | 8.5 | 90 | 1 | 1 | 6 | 11 | 18 | 23 | 50 |
| 65+ | 931 | 10.0 | 125 | 1 | 1 | 7 | 12 | 22 | 30 | 61 |
| **GRAND TOTAL** | **1886** | **9.1** | **112** | **1** | **1** | **6** | **11** | **20** | **28** | **59** |

## 589: Esophagoscopy and gastroscopy (natural orifice) (42.23; 44.13)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 590: Gastrotomy, pyloromyotomy (43.0; 43.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 289 | 3.2 | 1 | 2 | 2 | 3 | 4 | 5 | 6 | 8 |
| 20-34 | 3 | 4.0 | 7 | 2 | 2 | 3 | 6 | 7 | 7 | 7 |
| 35-49 | | | | | | | | | | |
| 50-64 | 5 | 10.0 | 3 | 8 | 8 | 11 | 11 | 12 | 12 | 12 |
| 65+ | 2 | 9.5 | 85 | 3 | 3 | 9 | 16 | 16 | 16 | 16 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 97 | 6.2 | 63 | 2 | 2 | 4 | 6 | 10 | 12 | 53 |
| 20-34 | 7 | 14.9 | 236 | 5 | 5 | 9 | 14 | 42 | 49 | 49 |
| 35-49 | 4 | 23.3 | 875 | 3 | 3 | 11 | 41 | 67 | 67 | 67 |
| 50-64 | 23 | 17.5 | 120 | 7 | 8 | 14 | 23 | 31 | 41 | 53 |
| 65+ | 35 | 23.3 | 197 | 7 | 8 | 21 | 31 | 40 | 54 | 68 |
| **TOT. SINGLE DX** | 299 | 3.4 | 3 | 2 | 2 | 3 | 4 | 5 | 7 | 11 |
| **" MULTIPLE DX** | 166 | 12.1 | 173 | 2 | 3 | 7 | 16 | 31 | 40 | 67 |
| TOTAL 0-19 YRS | 386 | 4.0 | 19 | 2 | 2 | 3 | 4 | 6 | 8 | 15 |
| 20-34 | 10 | 11.6 | 187 | 2 | 2 | 8 | 11 | 32 | 49 | 49 |
| 35-49 | 4 | 23.3 | 875 | 3 | 3 | 11 | 41 | 67 | 67 | 67 |
| 50-64 | 28 | 16.1 | 107 | 7 | 8 | 12 | 21 | 29 | 36 | 53 |
| 65+ | 37 | 22.6 | 199 | 5 | 8 | 20 | 31 | 39 | 53 | 68 |
| **GRAND TOTAL** | **465** | **6.5** | **81** | **2** | **2** | **4** | **6** | **14** | **24** | **52** |

## 591: Gastrostomy (43.1-43.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 6.5 | 5 | 5 | 5 | 6 | 8 | 8 | 8 | 8 |
| 20-34 | | | | | | | | | | |
| 35-49 | 6 | 24.3 | 1264 | 4 | 4 | 10 | 19 | 88 | 96 | 96 |
| 50-64 | 12 | 13.3 | 102 | 2 | 3 | 10 | 19 | 28 | 35 | 36 |
| 65+ | 12 | 22.9 | 547 | 2 | 4 | 18 | 24 | 57 | 84 | 88 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 35 | 29.9 | 650 | 4 | 6 | 19 | 38 | 84 | 97 | 99+ |
| 20-34 | 14 | 29.7 | 665 | 8 | 8 | 20 | 44 | 78 | 85 | 87 |
| 35-49 | 23 | 21.5 | 426 | 4 | 5 | 13 | 28 | 52 | 70 | 81 |
| 50-64 | 71 | 19.9 | 347 | 4 | 6 | 15 | 23 | 51 | 80 | 99+ |
| 65+ | 230 | 24.6 | 284 | 5 | 8 | 20 | 33 | 50 | 60 | 93 |
| **TOT. SINGLE DX** | 32 | 18.6 | 468 | 2 | 4 | 12 | 21 | 38 | 84 | 96 |
| **" MULTIPLE DX** | 373 | 24.2 | 357 | 5 | 7 | 18 | 32 | 56 | 69 | 99+ |
| TOTAL 0-19 YRS | 37 | 28.7 | 644 | 4 | 5 | 19 | 37 | 80 | 97 | 99+ |
| 20-34 | 14 | 29.7 | 665 | 8 | 8 | 20 | 44 | 78 | 85 | 87 |
| 35-49 | 29 | 22.1 | 562 | 4 | 5 | 13 | 27 | 58 | 82 | 96 |
| 50-64 | 83 | 19.0 | 316 | 4 | 6 | 14 | 22 | 44 | 75 | 99+ |
| 65+ | 242 | 24.5 | 296 | 5 | 7 | 20 | 32 | 50 | 61 | 92 |
| **GRAND TOTAL** | **405** | **23.8** | **367** | **5** | **6** | **18** | **31** | **54** | **71** | **99+** |

## 592: Local excision or destruction of lesion or tissue of stomach (43.4; 44.14-44.15)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 12 | 3.7 | 5 | 1 | 1 | 4 | 5 | 7 | 8 | 8 |
| 20-34 | 55 | 4.4 | 9 | <1 | 1 | 4 | 6 | 9 | 10 | 14 |
| 35-49 | 89 | 5.4 | 17 | <1 | <1 | 5 | 7 | 9 | 12 | 26 |
| 50-64 | 111 | 5.5 | 38 | <1 | <1 | 5 | 7 | 10 | 12 | 36 |
| 65+ | 146 | 5.2 | 20 | <1 | <1 | 4 | 7 | 9 | 13 | 22 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 8.9 | 178 | 1 | 2 | 6 | 9 | 14 | 41 | 62 |
| 20-34 | 161 | 7.3 | 51 | 2 | 2 | 6 | 8 | 12 | 20 | 50 |
| 35-49 | 337 | 7.0 | 47 | 2 | 3 | 6 | 8 | 12 | 15 | 35 |
| 50-64 | 746 | 8.0 | 56 | 2 | 3 | 6 | 10 | 15 | 21 | 41 |
| 65+ | 1490 | 10.4 | 117 | 2 | 3 | 7 | 12 | 19 | 29 | 65 |
| **TOT. SINGLE DX** | 413 | 5.2 | 23 | <1 | <1 | 4 | 7 | 9 | 12 | 22 |
| **" MULTIPLE DX** | 2753 | 9.1 | 91 | 2 | 3 | 7 | 11 | 17 | 25 | 54 |
| TOTAL 0-19 YRS | 31 | 6.9 | 116 | 1 | 1 | 5 | 8 | 10 | 15 | 62 |
| 20-34 | 216 | 6.5 | 42 | 1 | 2 | 5 | 8 | 11 | 15 | 36 |
| 35-49 | 426 | 6.7 | 42 | 1 | 2 | 5 | 8 | 12 | 15 | 33 |
| 50-64 | 857 | 7.6 | 55 | 1 | 2 | 6 | 9 | 14 | 20 | 41 |
| 65+ | 1636 | 9.9 | 111 | 2 | 3 | 7 | 11 | 18 | 28 | 52 |
| **GRAND TOTAL** | **3166** | **8.6** | **84** | **1** | **2** | **6** | **10** | **16** | **23** | **52** |

## 593: Partial gastrectomy (43.5-43.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 8.7 | 4 | 7 | 7 | 8 | 10 | 11 | 11 | 11 |
| 20-34 | 47 | 10.1 | 12 | 6 | 7 | 10 | 12 | 15 | 17 | 19 |
| 35-49 | 105 | 10.3 | 15 | 6 | 7 | 9 | 11 | 16 | 18 | 27 |
| 50-64 | 89 | 11.1 | 17 | 7 | 7 | 10 | 13 | 16 | 20 | 28 |
| 65+ | 71 | 14.5 | 48 | 7 | 8 | 13 | 17 | 24 | 32 | 37 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 14.8 | 85 | 7 | 7 | 11 | 21 | 29 | 29 | 29 |
| 20-34 | 54 | 13.2 | 150 | 6 | 7 | 10 | 13 | 20 | 32 | 73 |
| 35-49 | 154 | 14.6 | 126 | 7 | 8 | 11 | 17 | 23 | 30 | 71 |
| 50-64 | 314 | 16.8 | 147 | 7 | 8 | 13 | 20 | 31 | 44 | 97 |
| 65+ | 442 | 21.2 | 191 | 8 | 10 | 17 | 25 | 39 | 49 | 77 |
| **TOT. SINGLE DX** | 315 | 11.5 | 26 | 6 | 7 | 10 | 13 | 17 | 21 | 33 |
| **" MULTIPLE DX** | 969 | 18.2 | 172 | 7 | 8 | 14 | 21 | 35 | 47 | 82 |
| TOTAL 0-19 YRS | 8 | 12.5 | 60 | 7 | 7 | 9 | 15 | 26 | 29 | 29 |
| 20-34 | 101 | 11.8 | 88 | 6 | 7 | 10 | 12 | 17 | 21 | 68 |
| 35-49 | 259 | 12.8 | 86 | 6 | 7 | 10 | 14 | 21 | 24 | 62 |
| 50-64 | 403 | 15.6 | 124 | 7 | 8 | 12 | 18 | 30 | 39 | 90 |
| 65+ | 513 | 20.3 | 177 | 8 | 9 | 16 | 23 | 37 | 47 | 76 |
| **GRAND TOTAL** | **1284** | **16.6** | **144** | **7** | **8** | **12** | **19** | **31** | **41** | **72** |

## 594: Total gastrectomy (43.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 2 | 12.0 | 8 | 10 | 10 | 12 | 14 | 14 | 14 | 14 |
| 35-49 | 4 | 13.8 | 31 | 10 | 10 | 11 | 17 | 22 | 22 | 22 |
| 50-64 | 12 | 16.5 | 43 | 10 | 11 | 14 | 19 | 29 | 30 | 30 |
| 65+ | 10 | 16.5 | 16 | 9 | 11 | 17 | 19 | 22 | 24 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 3 | 18.0 | 39 | 13 | 13 | 16 | 23 | 25 | 25 | 25 |
| 35-49 | 15 | 23.1 | 114 | 11 | 12 | 20 | 29 | 40 | 44 | 45 |
| 50-64 | 60 | 20.8 | 153 | 10 | 11 | 16 | 25 | 39 | 50 | 64 |
| 65+ | 74 | 20.7 | 71 | 10 | 12 | 20 | 26 | 30 | 36 | 90 |
| **TOT. SINGLE DX** | 28 | 15.8 | 29 | 10 | 10 | 14 | 18 | 23 | 28 | 30 |
| **" MULTIPLE DX** | 152 | 20.9 | 106 | 11 | 12 | 18 | 26 | 35 | 40 | 65 |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | 5 | 15.6 | 32 | 10 | 10 | 14 | 18 | 25 | 25 | 25 |
| 35-49 | 19 | 21.1 | 110 | 10 | 11 | 19 | 23 | 40 | 43 | 45 |
| 50-64 | 72 | 20.1 | 137 | 10 | 11 | 16 | 23 | 38 | 43 | 64 |
| 65+ | 84 | 20.2 | 66 | 10 | 12 | 19 | 25 | 30 | 35 | 86 |
| **GRAND TOTAL** | **180** | **20.1** | **97** | **10** | **11** | **17** | **25** | **32** | **40** | **64** |

## 595: Vagotomy (44.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 39 | 8.9 | 8 | 6 | 6 | 8 | 10 | 12 | 15 | 19 |
| 35-49 | 58 | 10.0 | 17 | 5 | 6 | 9 | 11 | 14 | 18 | 27 |
| 50-64 | 75 | 10.5 | 12 | 6 | 7 | 9 | 13 | 16 | 17 | 20 |
| 65+ | 44 | 14.5 | 51 | 7 | 8 | 12 | 16 | 29 | 32 | 32 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 11.0 | | 11 | 11 | 11 | 11 | 11 | 11 | 11 |
| 20-34 | 31 | 13.0 | 101 | 5 | 6 | 9 | 13 | 31 | 39 | 42 |
| 35-49 | 98 | 12.6 | 48 | 6 | 6 | 11 | 15 | 21 | 26 | 43 |
| 50-64 | 131 | 12.9 | 58 | 7 | 7 | 11 | 15 | 20 | 26 | 74 |
| 65+ | 174 | 18.6 | 136 | 8 | 9 | 14 | 23 | 36 | 46 | 59 |
| **TOT. SINGLE DX** | 217 | 10.9 | 24 | 6 | 6 | 9 | 12 | 17 | 20 | 32 |
| **" MULTIPLE DX** | 435 | 15.1 | 98 | 6 | 7 | 12 | 18 | 27 | 39 | 54 |
| TOTAL 0-19 YRS | 2 | 9.0 | 8 | 7 | 7 | 9 | 11 | 11 | 11 | 11 |
| 20-34 | 70 | 10.7 | 53 | 6 | 6 | 8 | 11 | 17 | 29 | 41 |
| 35-49 | 156 | 11.6 | 38 | 6 | 6 | 10 | 13 | 20 | 25 | 41 |
| 50-64 | 206 | 12.0 | 43 | 6 | 7 | 10 | 15 | 18 | 22 | 55 |
| 65+ | 218 | 17.8 | 122 | 8 | 9 | 14 | 21 | 32 | 44 | 56 |
| **GRAND TOTAL** | **652** | **13.7** | **77** | **6** | **7** | **11** | **16** | **24** | **32** | **49** |

## 596: Miscellaneous operations on stomach (44.11; 44.2-44.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 35 | 6.5 | 8 | 2 | 3 | 6 | 8 | 10 | 12 | 14 |
| 20-34 | 227 | 7.3 | 6 | 5 | 5 | 7 | 8 | 9 | 11 | 17 |
| 35-49 | 227 | 7.7 | 6 | 5 | 5 | 7 | 9 | 10 | 13 | 17 |
| 50-64 | 103 | 8.8 | 14 | 5 | 5 | 8 | 10 | 13 | 16 | 25 |
| 65+ | 91 | 11.5 | 32 | 6 | 6 | 10 | 14 | 17 | 26 | 29 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 48 | 18.7 | 360 | 4 | 5 | 10 | 25 | 50 | 75 | 99+ |
| 20-34 | 411 | 9.1 | 40 | 5 | 5 | 8 | 9 | 14 | 18 | 48 |
| 35-49 | 548 | 10.9 | 71 | 5 | 6 | 8 | 11 | 19 | 31 | 53 |
| 50-64 | 421 | 13.9 | 117 | 6 | 7 | 10 | 15 | 25 | 37 | 62 |
| 65+ | 471 | 18.4 | 153 | 7 | 8 | 15 | 23 | 34 | 46 | 81 |
| **TOT. SINGLE DX** | 683 | 8.2 | 13 | 5 | 5 | 7 | 9 | 12 | 14 | 26 |
| **" MULTIPLE DX** | 1899 | 13.2 | 115 | 5 | 6 | 9 | 15 | 25 | 36 | 62 |
| TOTAL 0-19 YRS | 83 | 13.5 | 247 | 3 | 4 | 7 | 14 | 39 | 55 | 94 |
| 20-34 | 638 | 8.5 | 29 | 5 | 5 | 7 | 9 | 12 | 16 | 32 |
| 35-49 | 775 | 10.0 | 55 | 5 | 6 | 8 | 10 | 16 | 25 | 50 |
| 50-64 | 524 | 12.9 | 101 | 5 | 6 | 10 | 14 | 22 | 35 | 59 |
| 65+ | 562 | 17.3 | 140 | 7 | 8 | 14 | 21 | 30 | 40 | 70 |
| **GRAND TOTAL** | **2582** | **11.9** | **93** | **5** | **6** | **9** | **13** | **22** | **31** | **57** |

## 597: Local excision or destruction of intestinal lesion or tissue (45.14-45.15; 45.25-45.27; 45.3-45.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 67 | 4.7 | 16 | <1 | <1 | 4 | 6 | 8 | 10 | 26 |
| 20-34 | 153 | 5.5 | 15 | 1 | 1 | 5 | 8 | 11 | 13 | 20 |
| 35-49 | 165 | 5.1 | 20 | <1 | <1 | 4 | 7 | 9 | 13 | 19 |
| 50-64 | 209 | 3.8 | 12 | <1 | <1 | 2 | 5 | 9 | 11 | 15 |
| 65+ | 259 | 5.0 | 27 | <1 | 1 | 3 | 7 | 11 | 15 | 23 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 67 | 8.7 | 189 | 1 | 2 | 5 | 8 | 12 | 37 | 74 |
| 20-34 | 252 | 7.2 | 35 | 1 | 2 | 5 | 9 | 14 | 21 | 34 |
| 35-49 | 327 | 7.5 | 46 | 1 | 1 | 6 | 9 | 15 | 21 | 37 |
| 50-64 | 740 | 8.3 | 74 | 1 | 1 | 6 | 10 | 17 | 22 | 46 |
| 65+ | 1499 | 9.5 | 100 | 1 | 2 | 7 | 12 | 19 | 28 | 54 |
| **TOT. SINGLE DX** | 853 | 4.8 | 20 | <1 | 1 | 4 | 7 | 10 | 13 | 19 |
| **" MULTIPLE DX** | 2885 | 8.7 | 85 | 1 | 2 | 6 | 11 | 18 | 25 | 49 |
| TOTAL 0-19 YRS | 134 | 6.7 | 106 | <1 | 1 | 5 | 7 | 10 | 14 | 66 |
| 20-34 | 405 | 6.5 | 28 | 1 | 2 | 5 | 8 | 12 | 17 | 29 |
| 35-49 | 492 | 6.7 | 39 | <1 | 1 | 5 | 9 | 13 | 18 | 36 |
| 50-64 | 949 | 7.3 | 64 | <1 | 1 | 5 | 9 | 15 | 21 | 41 |
| 65+ | 1758 | 8.8 | 92 | 1 | 1 | 6 | 11 | 19 | 26 | 51 |
| **GRAND TOTAL** | **3738** | **7.8** | **73** | **1** | **1** | **6** | **10** | **16** | **22** | **45** |

## 598: Resection of small intestine (45.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 12 | 9.0 | 33 | 3 | 3 | 7 | 12 | 18 | 21 | 21 |
| 20-34 | 29 | 10.0 | 45 | 4 | 4 | 8 | 11 | 16 | 22 | 38 |
| 35-49 | 30 | 11.3 | 56 | 6 | 7 | 9 | 11 | 18 | 22 | 45 |
| 50-64 | 29 | 13.5 | 134 | 7 | 8 | 10 | 12 | 25 | 44 | 60 |
| 65+ | 50 | 12.6 | 12 | 8 | 8 | 12 | 15 | 18 | 19 | 21 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 48 | 14.9 | 115 | 6 | 6 | 10 | 24 | 35 | 51 | 99+ |
| 20-34 | 99 | 13.6 | 116 | 5 | 7 | 10 | 15 | 25 | 37 | 60 |
| 35-49 | 114 | 16.8 | 169 | 6 | 7 | 12 | 20 | 31 | 43 | 77 |
| 50-64 | 199 | 17.7 | 189 | 6 | 7 | 13 | 21 | 37 | 50 | 99+ |
| 65+ | 396 | 20.4 | 207 | 8 | 9 | 16 | 24 | 40 | 54 | 97 |
| **TOT. SINGLE DX** | 150 | 11.7 | 53 | 5 | 6 | 10 | 13 | 18 | 21 | 45 |
| **" MULTIPLE DX** | 856 | 18.2 | 187 | 7 | 8 | 14 | 22 | 37 | 51 | 99+ |
| TOTAL 0-19 YRS | 60 | 13.7 | 104 | 4 | 5 | 10 | 18 | 33 | 43 | 99+ |
| 20-34 | 128 | 12.8 | 102 | 5 | 6 | 10 | 14 | 23 | 37 | 56 |
| 35-49 | 144 | 15.7 | 151 | 6 | 7 | 11 | 19 | 29 | 42 | 73 |
| 50-64 | 228 | 17.1 | 184 | 6 | 7 | 13 | 20 | 35 | 48 | 99+ |
| 65+ | 446 | 19.5 | 191 | 8 | 9 | 15 | 23 | 38 | 53 | 95 |
| **GRAND TOTAL** | **1006** | **17.2** | **173** | **6** | **7** | **13** | **20** | **35** | **47** | **97** |

## 599: Colectomy (45.7-45.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 26 | 7.6 | 10 | 4 | 4 | 7 | 10 | 11 | 14 | 17 |
| 20-34 | 83 | 10.0 | 19 | 6 | 6 | 9 | 12 | 16 | 18 | 29 |
| 35-49 | 156 | 10.7 | 19 | 6 | 7 | 10 | 12 | 16 | 19 | 31 |
| 50-64 | 433 | 10.6 | 20 | 7 | 7 | 10 | 12 | 15 | 18 | 24 |
| 65+ | 590 | 12.0 | 20 | 7 | 8 | 11 | 14 | 17 | 21 | 30 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 47 | 14.4 | 179 | 5 | 6 | 10 | 13 | 29 | 49 | 67 |
| 20-34 | 174 | 15.2 | 91 | 6 | 7 | 12 | 18 | 31 | 36 | 48 |
| 35-49 | 444 | 14.3 | 99 | 7 | 7 | 11 | 16 | 24 | 35 | 69 |
| 50-64 | 1233 | 14.8 | 96 | 7 | 8 | 12 | 17 | 25 | 35 | 62 |
| 65+ | 2816 | 17.1 | 105 | 8 | 9 | 14 | 20 | 29 | 37 | 60 |
| **TOT. SINGLE DX** | 1288 | 11.1 | 21 | 6 | 7 | 10 | 13 | 17 | 19 | 28 |
| **" MULTIPLE DX** | 4714 | 16.1 | 104 | 7 | 8 | 13 | 19 | 28 | 36 | 61 |
| TOTAL 0-19 YRS | 73 | 12.0 | 129 | 4 | 5 | 9 | 11 | 23 | 33 | 66 |
| 20-34 | 257 | 13.5 | 74 | 6 | 7 | 10 | 14 | 27 | 33 | 40 |
| 35-49 | 600 | 13.3 | 81 | 6 | 7 | 11 | 15 | 22 | 29 | 63 |
| 50-64 | 1666 | 13.8 | 80 | 7 | 8 | 11 | 15 | 22 | 31 | 56 |
| 65+ | 3406 | 16.2 | 95 | 8 | 9 | 13 | 19 | 27 | 35 | 59 |
| **GRAND TOTAL** | **6002** | **15.1** | **90** | **7** | **8** | **12** | **17** | **25** | **33** | **58** |

## 600: Other incision, resection of intestine (45.0; 45.5; 46.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 6 | 8.0 | 10 | 4 | 4 | 8 | 11 | 12 | 12 | 12 |
| 20-34 | 5 | 7.6 | 16 | 3 | 3 | 7 | 9 | 14 | 14 | 14 |
| 35-49 | 18 | 9.7 | 13 | 4 | 6 | 9 | 11 | 14 | 18 | 20 |
| 50-64 | 18 | 11.0 | 24 | 3 | 5 | 11 | 13 | 15 | 21 | 25 |
| 65+ | 30 | 11.0 | 28 | 6 | 7 | 9 | 11 | 19 | 21 | 28 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 10 | 7.8 | 13 | 2 | 3 | 7 | 11 | 13 | 13 | 13 |
| 20-34 | 19 | 13.2 | 201 | 1 | 3 | 10 | 12 | 33 | 49 | 59 |
| 35-49 | 29 | 16.0 | 72 | 6 | 7 | 14 | 21 | 27 | 31 | 44 |
| 50-64 | 67 | 18.3 | 228 | 7 | 8 | 14 | 18 | 38 | 56 | 79 |
| 65+ | 148 | 20.0 | 176 | 8 | 8 | 16 | 23 | 36 | 44 | 74 |
| **TOT. SINGLE DX** | 77 | 10.2 | 22 | 4 | 6 | 10 | 11 | 16 | 20 | 27 |
| **" MULTIPLE DX** | 273 | 18.2 | 180 | 6 | 7 | 14 | 22 | 34 | 44 | 73 |
| TOTAL 0-19 YRS | 16 | 7.9 | 11 | 3 | 4 | 8 | 11 | 13 | 13 | 13 |
| 20-34 | 24 | 12.0 | 166 | 1 | 3 | 8 | 11 | 27 | 44 | 59 |
| 35-49 | 47 | 13.6 | 59 | 6 | 7 | 11 | 18 | 22 | 27 | 44 |
| 50-64 | 85 | 16.7 | 193 | 7 | 7 | 12 | 17 | 32 | 47 | 76 |
| 65+ | 178 | 18.5 | 163 | 7 | 8 | 15 | 22 | 33 | 42 | 73 |
| **GRAND TOTAL** | **350** | **16.5** | **156** | **6** | **7** | **13** | **19** | **31** | **41** | **71** |

## 601: Intestinal anastomosis or repair (45.9; 46.4-46.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 93 | 6.3 | 9 | 2 | 3 | 6 | 8 | 10 | 12 | 16 |
| 20-34 | 97 | 8.4 | 17 | 3 | 4 | 8 | 10 | 13 | 17 | 25 |
| 35-49 | 96 | 9.3 | 25 | 4 | 5 | 9 | 11 | 13 | 16 | 36 |
| 50-64 | 122 | 8.6 | 16 | 3 | 5 | 8 | 10 | 12 | 14 | 30 |
| 65+ | 168 | 10.3 | 20 | 5 | 5 | 10 | 12 | 16 | 18 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 112 | 10.1 | 96 | 3 | 4 | 7 | 11 | 19 | 27 | 61 |
| 20-34 | 221 | 11.9 | 106 | 4 | 5 | 9 | 13 | 22 | 35 | 68 |
| 35-49 | 238 | 12.2 | 79 | 5 | 6 | 9 | 14 | 21 | 37 | 56 |
| 50-64 | 370 | 13.2 | 87 | 5 | 6 | 10 | 15 | 25 | 34 | 72 |
| 65+ | 587 | 15.3 | 123 | 6 | 7 | 12 | 17 | 29 | 41 | 60 |
| **TOT. SINGLE DX** | 576 | 8.8 | 20 | 3 | 4 | 8 | 11 | 13 | 16 | 27 |
| **" MULTIPLE DX** | 1528 | 13.5 | 106 | 5 | 6 | 10 | 15 | 25 | 37 | 62 |
| TOTAL 0-19 YRS | 205 | 8.4 | 60 | 3 | 3 | 6 | 9 | 14 | 21 | 48 |
| 20-34 | 318 | 10.9 | 82 | 4 | 5 | 9 | 12 | 19 | 28 | 59 |
| 35-49 | 334 | 11.4 | 65 | 5 | 5 | 9 | 12 | 19 | 29 | 50 |
| 50-64 | 492 | 12.1 | 74 | 5 | 6 | 10 | 13 | 23 | 30 | 65 |
| 65+ | 755 | 14.2 | 105 | 5 | 7 | 11 | 16 | 25 | 37 | 59 |
| **GRAND TOTAL** | **2104** | **12.2** | **86** | **4** | **5** | **10** | **13** | **22** | **32** | **58** |

## 602: Ileostomy, colostomy, and other enterostomy (46.1-46.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 10.1 | 70 | 3 | 3 | 6 | 13 | 25 | 27 | 27 |
| 20-34 | 4 | 11.5 | 9 | 9 | 9 | 10 | 13 | 16 | 16 | 16 |
| 35-49 | 12 | 13.5 | 87 | 7 | 8 | 9 | 15 | 27 | 38 | 40 |
| 50-64 | 25 | 14.3 | 85 | 6 | 7 | 11 | 16 | 27 | 38 | 43 |
| 65+ | 40 | 17.6 | 185 | 6 | 7 | 13 | 20 | 34 | 47 | 73 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 15 | 18.3 | 374 | 6 | 8 | 12 | 17 | 30 | 70 | 84 |
| 20-34 | 29 | 17.7 | 201 | 6 | 7 | 13 | 20 | 31 | 59 | 67 |
| 35-49 | 58 | 18.7 | 260 | 4 | 6 | 13 | 24 | 48 | 64 | 98 |
| 50-64 | 128 | 19.8 | 182 | 6 | 9 | 16 | 27 | 41 | 67 | 99+ |
| 65+ | 332 | 22.3 | 205 | 8 | 9 | 19 | 28 | 41 | 51 | 96 |
| **TOT. SINGLE DX** | 88 | 15.2 | 129 | 5 | 7 | 11 | 18 | 27 | 39 | 66 |
| **" MULTIPLE DX** | 562 | 21.1 | 211 | 6 | 9 | 17 | 27 | 41 | 55 | 99+ |
| TOTAL 0-19 YRS | 22 | 15.7 | 285 | 4 | 5 | 11 | 15 | 28 | 52 | 84 |
| 20-34 | 33 | 16.9 | 181 | 6 | 8 | 13 | 18 | 28 | 55 | 67 |
| 35-49 | 70 | 17.8 | 233 | 5 | 7 | 13 | 21 | 44 | 56 | 95 |
| 50-64 | 153 | 18.9 | 171 | 6 | 8 | 15 | 25 | 39 | 59 | 99+ |
| 65+ | 372 | 21.8 | 205 | 7 | 9 | 18 | 28 | 40 | 51 | 94 |
| **GRAND TOTAL** | **650** | **20.3** | **204** | **6** | **8** | **16** | **26** | **40** | **52** | **99+** |

## 603: Appendectomy (47.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 5116 | 3.7 | 4 | 2 | 2 | 3 | 4 | 6 | 8 | 11 |
| 20-34 | 3933 | 3.8 | 3 | 2 | 2 | 3 | 4 | 6 | 7 | 11 |
| 35-49 | 1228 | 4.5 | 5 | 2 | 2 | 4 | 5 | 8 | 9 | 12 |
| 50-64 | 530 | 5.6 | 8 | 2 | 3 | 5 | 7 | 9 | 11 | 15 |
| 65+ | 217 | 6.6 | 11 | 3 | 3 | 6 | 8 | 11 | 13 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1287 | 5.7 | 16 | 2 | 2 | 4 | 7 | 10 | 13 | 23 |
| 20-34 | 1283 | 5.6 | 16 | 2 | 3 | 4 | 7 | 10 | 13 | 21 |
| 35-49 | 496 | 7.3 | 26 | 3 | 3 | 6 | 9 | 12 | 18 | 27 |
| 50-64 | 331 | 8.8 | 36 | 3 | 3 | 7 | 10 | 15 | 20 | 34 |
| 65+ | 342 | 11.6 | 67 | 4 | 5 | 9 | 14 | 22 | 27 | 45 |
| **TOT. SINGLE DX** | 11024 | 4.0 | 5 | 2 | 2 | 3 | 5 | 7 | 8 | 12 |
| **" MULTIPLE DX** | 3789 | 6.7 | 28 | 2 | 3 | 5 | 8 | 12 | 16 | 27 |
| TOTAL 0-19 YRS | 6403 | 4.1 | 7 | 2 | 2 | 3 | 5 | 7 | 9 | 15 |
| 20-34 | 5216 | 4.3 | 7 | 2 | 2 | 4 | 5 | 7 | 9 | 15 |
| 35-49 | 1724 | 5.3 | 13 | 2 | 2 | 4 | 6 | 9 | 11 | 20 |
| 50-64 | 911 | 6.9 | 23 | 3 | 3 | 6 | 8 | 12 | 15 | 26 |
| 65+ | 559 | 9.6 | 52 | 3 | 4 | 8 | 11 | 17 | 23 | 43 |
| **GRAND TOTAL** | **14813** | **4.7** | **12** | **2** | **2** | **4** | **5** | **8** | **11** | **19** |

## 604: Incidental appendectomy (47.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 69 | 3.7 | 3 | 1 | 2 | 3 | 4 | 6 | 8 | 9 |
| 20-34 | 51 | 3.8 | 2 | 2 | 2 | 4 | 5 | 5 | 6 | 9 |
| 35-49 | 13 | 3.7 | 3 | 1 | 1 | 4 | 5 | 6 | 7 | 7 |
| 50-64 | 5 | 4.4 | 1 | 3 | 3 | 4 | 5 | 6 | 6 | 6 |
| 65+ | 2 | 8.0 | 8 | 6 | 6 | 8 | 10 | 10 | 10 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 37 | 4.9 | 10 | 2 | 3 | 4 | 6 | 8 | 9 | 20 |
| 20-34 | 45 | 5.3 | 6 | 2 | 3 | 5 | 6 | 9 | 12 | 13 |
| 35-49 | 15 | 7.1 | 27 | 2 | 3 | 6 | 8 | 11 | 21 | 24 |
| 50-64 | 8 | 17.6 | 228 | 7 | 7 | 10 | 25 | 43 | 49 | 49 |
| 65+ | 9 | 14.4 | 84 | 7 | 7 | 9 | 21 | 29 | 33 | 33 |
| **TOT. SINGLE DX** | 140 | 3.8 | 3 | 2 | 2 | 4 | 4 | 5 | 7 | 9 |
| **" MULTIPLE DX** | 114 | 7.0 | 44 | 2 | 3 | 5 | 7 | 13 | 21 | 39 |
| TOTAL 0-19 YRS | 106 | 4.1 | 6 | 2 | 2 | 4 | 5 | 7 | 8 | 14 |
| 20-34 | 96 | 4.5 | 5 | 2 | 3 | 4 | 5 | 7 | 9 | 13 |
| 35-49 | 28 | 5.5 | 19 | 1 | 2 | 5 | 7 | 9 | 12 | 24 |
| 50-64 | 13 | 12.5 | 178 | 3 | 4 | 7 | 15 | 34 | 46 | 49 |
| 65+ | 11 | 13.3 | 75 | 6 | 6 | 9 | 19 | 27 | 32 | 33 |
| **GRAND TOTAL** | **254** | **5.3** | **24** | **2** | **2** | **4** | **6** | **8** | **12** | **30** |

## 605: Operations on appendix, except appendectomy (47.2-47.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 23 | 8.0 | 23 | 2 | 2 | 7 | 11 | 16 | 18 | 19 |
| 20-34 | 20 | 6.7 | 13 | 2 | 3 | 6 | 9 | 11 | 14 | 16 |
| 35-49 | 6 | 11.2 | 15 | 4 | 5 | 12 | 14 | 15 | 15 | 15 |
| 50-64 | 3 | 11.7 | 22 | 8 | 8 | 10 | 15 | 17 | 17 | 17 |
| 65+ | 5 | 15.8 | 23 | 8 | 8 | 17 | 19 | 21 | 21 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 7.9 | 5 | 4 | 4 | 8 | 9 | 11 | 12 | 12 |
| 20-34 | 8 | 9.3 | 28 | 1 | 2 | 8 | 14 | 16 | 17 | 17 |
| 35-49 | 5 | 10.2 | 10 | 6 | 6 | 10 | 11 | 15 | 15 | 15 |
| 50-64 | 9 | 14.1 | 46 | 6 | 6 | 13 | 16 | 24 | 29 | 29 |
| 65+ | 5 | 16.4 | 128 | 7 | 7 | 11 | 27 | 31 | 31 | 31 |
| **TOT. SINGLE DX** | 57 | 8.7 | 25 | 2 | 3 | 8 | 12 | 17 | 18 | 21 |
| **" MULTIPLE DX** | 41 | 10.8 | 40 | 4 | 6 | 9 | 13 | 17 | 27 | 31 |
| TOTAL 0-19 YRS | 37 | 7.9 | 16 | 2 | 3 | 8 | 10 | 12 | 17 | 19 |
| 20-34 | 28 | 7.4 | 18 | 2 | 3 | 7 | 9 | 14 | 16 | 17 |
| 35-49 | 11 | 10.7 | 12 | 4 | 5 | 10 | 14 | 15 | 15 | 15 |
| 50-64 | 12 | 13.5 | 39 | 6 | 6 | 13 | 16 | 21 | 28 | 29 |
| 65+ | 10 | 16.1 | 67 | 7 | 7 | 16 | 21 | 28 | 31 | 31 |
| **GRAND TOTAL** | **98** | **9.6** | **32** | **2** | **4** | **8** | **12** | **17** | **19** | **30** |

## 606: Local excision or destruction of lesion or tissue of rectum (48.24-48.26; 48.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 59 | 4.7 | 74 | <1 | <1 | 2 | 4 | 11 | 21 | 53 |
| 20-34 | 77 | 3.1 | 5 | <1 | <1 | 2 | 4 | 6 | 7 | 13 |
| 35-49 | 71 | 3.0 | 6 | 1 | 1 | 2 | 3 | 6 | 9 | 13 |
| 50-64 | 102 | 3.5 | 12 | <1 | 1 | 2 | 4 | 7 | 9 | 22 |
| 65+ | 171 | 4.2 | 12 | <1 | 1 | 3 | 5 | 8 | 11 | 20 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 28 | 4.4 | 61 | <1 | 1 | 2 | 4 | 6 | 29 | 35 |
| 20-34 | 73 | 6.7 | 71 | 1 | 1 | 4 | 8 | 12 | 14 | 55 |
| 35-49 | 86 | 6.7 | 78 | 1 | 2 | 4 | 6 | 15 | 28 | 48 |
| 50-64 | 173 | 7.4 | 52 | 1 | 2 | 5 | 10 | 15 | 23 | 38 |
| 65+ | 438 | 8.3 | 74 | 1 | 2 | 5 | 10 | 18 | 25 | 45 |
| **TOT. SINGLE DX** | 480 | 3.7 | 18 | <1 | 1 | 3 | 4 | 7 | 11 | 22 |
| **" MULTIPLE DX** | 798 | 7.7 | 70 | 1 | 2 | 5 | 9 | 16 | 24 | 44 |
| TOTAL 0-19 YRS | 87 | 4.6 | 70 | <1 | <1 | 2 | 4 | 11 | 27 | 48 |
| 20-34 | 150 | 4.8 | 40 | <1 | 1 | 3 | 6 | 9 | 12 | 45 |
| 35-49 | 157 | 5.1 | 49 | 1 | 1 | 3 | 5 | 11 | 16 | 39 |
| 50-64 | 275 | 6.0 | 41 | <1 | 1 | 4 | 7 | 14 | 19 | 31 |
| 65+ | 609 | 7.1 | 60 | 1 | 2 | 5 | 9 | 15 | 21 | 43 |
| **GRAND TOTAL** | **1278** | **6.2** | **54** | **1** | **1** | **4** | **7** | **13** | **20** | **42** |

## 607: Other incision of rectum and incision or excision of perirectal tissue (48.0; 48.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 30 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 7 |
| 20-34 | 133 | 2.7 | 2 | 1 | 1 | 2 | 3 | 4 | 6 | 8 |
| 35-49 | 135 | 3.2 | 5 | 1 | 1 | 2 | 4 | 7 | 8 | 11 |
| 50-64 | 67 | 2.6 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 65+ | 28 | 5.2 | 49 | 1 | 1 | 4 | 5 | 8 | 13 | 39 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 8.8 | 82 | 1 | 1 | 5 | 14 | 23 | 30 | 30 |
| 20-34 | 59 | 3.9 | 12 | 1 | 1 | 3 | 4 | 10 | 13 | 15 |
| 35-49 | 80 | 5.4 | 37 | 1 | 1 | 3 | 6 | 13 | 17 | 34 |
| 50-64 | 70 | 5.7 | 27 | 1 | 1 | 4 | 8 | 13 | 14 | 26 |
| 65+ | 45 | 6.8 | 27 | 2 | 3 | 5 | 8 | 17 | 21 | 21 |
| **TOT. SINGLE DX** | 393 | 3.0 | 7 | 1 | 1 | 2 | 4 | 5 | 8 | 10 |
| **" MULTIPLE DX** | 264 | 5.5 | 30 | 1 | 1 | 4 | 6 | 13 | 17 | 26 |
| TOTAL 0-19 YRS | 40 | 4.0 | 28 | 1 | 1 | 3 | 4 | 7 | 15 | 30 |
| 20-34 | 192 | 3.1 | 6 | 1 | 1 | 2 | 4 | 5 | 8 | 14 |
| 35-49 | 215 | 4.0 | 19 | 1 | 1 | 3 | 5 | 8 | 12 | 21 |
| 50-64 | 137 | 4.2 | 17 | 1 | 1 | 3 | 5 | 8 | 13 | 26 |
| 65+ | 73 | 6.2 | 36 | 1 | 2 | 5 | 7 | 11 | 21 | 35 |
| **GRAND TOTAL** | **657** | **4.0** | **18** | **1** | **1** | **3** | **4** | **8** | **12** | **21** |

## 608: Proctectomy (48.4-48.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 7.8 | 2 | 5 | 5 | 8 | 9 | 10 | 10 | 10 |
| 20-34 | 11 | 12.4 | 49 | 6 | 6 | 11 | 14 | 23 | 30 | 31 |
| 35-49 | 36 | 11.4 | 28 | 1 | 5 | 12 | 13 | 16 | 22 | 28 |
| 50-64 | 110 | 12.3 | 29 | 5 | 7 | 12 | 15 | 18 | 20 | 32 |
| 65+ | 136 | 14.2 | 28 | 8 | 9 | 13 | 16 | 20 | 24 | 35 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 12.2 | 44 | 6 | 7 | 10 | 14 | 17 | 28 | 36 |
| 20-34 | 33 | 17.0 | 155 | 7 | 8 | 11 | 19 | 37 | 45 | 56 |
| 35-49 | 71 | 16.3 | 114 | 7 | 8 | 12 | 18 | 30 | 51 | 90 |
| 50-64 | 285 | 16.8 | 119 | 8 | 9 | 14 | 19 | 27 | 35 | 69 |
| 65+ | 522 | 18.5 | 113 | 9 | 10 | 15 | 23 | 33 | 41 | 95 |
| **TOT. SINGLE DX** | 301 | 12.9 | 30 | 5 | 8 | 12 | 15 | 19 | 22 | 33 |
| **" MULTIPLE DX** | 930 | 17.6 | 117 | 8 | 9 | 15 | 21 | 32 | 40 | 75 |
| TOTAL 0-19 YRS | 27 | 10.9 | 36 | 6 | 6 | 9 | 12 | 16 | 21 | 36 |
| 20-34 | 44 | 15.9 | 131 | 6 | 7 | 11 | 18 | 34 | 41 | 56 |
| 35-49 | 107 | 14.6 | 90 | 5 | 8 | 12 | 15 | 28 | 40 | 73 |
| 50-64 | 395 | 15.5 | 99 | 7 | 8 | 13 | 17 | 23 | 33 | 60 |
| 65+ | 658 | 17.6 | 99 | 8 | 10 | 15 | 21 | 31 | 38 | 61 |
| **GRAND TOTAL** | **1231** | **16.5** | **100** | **7** | **9** | **14** | **19** | **28** | **37** | **58** |

## 609: Miscellaneous operations on intestine (45.11; 45.13; 45.21; 45.23; 48.1; 48.21; 48.7; 48.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 91 | 3.4 | 6 | <1 | <1 | 3 | 5 | 6 | 7 | 15 |
| 20-34 | 383 | 4.5 | 18 | <1 | 1 | 4 | 6 | 8 | 10 | 19 |
| 35-49 | 409 | 4.8 | 15 | <1 | 1 | 4 | 6 | 9 | 12 | 23 |
| 50-64 | 408 | 4.7 | 13 | <1 | <1 | 4 | 6 | 9 | 11 | 18 |
| 65+ | 408 | 5.7 | 29 | <1 | 1 | 4 | 7 | 10 | 14 | 31 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 152 | 6.3 | 58 | 1 | 1 | 4 | 7 | 13 | 19 | 50 |
| 20-34 | 1010 | 6.1 | 33 | 1 | 2 | 5 | 7 | 12 | 16 | 32 |
| 35-49 | 1427 | 6.5 | 30 | 2 | 2 | 5 | 8 | 12 | 16 | 30 |
| 50-64 | 2240 | 7.3 | 49 | 1 | 2 | 5 | 9 | 14 | 19 | 33 |
| 65+ | 3453 | 9.1 | 65 | 2 | 3 | 7 | 11 | 18 | 25 | 43 |
| **TOT. SINGLE DX** | 1699 | 4.8 | 19 | <1 | 1 | 4 | 6 | 9 | 11 | 21 |
| **" MULTIPLE DX** | 8282 | 7.8 | 52 | 2 | 2 | 6 | 9 | 15 | 21 | 38 |
| TOTAL 0-19 YRS | 243 | 5.2 | 41 | <1 | 1 | 4 | 6 | 9 | 15 | 43 |
| 20-34 | 1393 | 5.7 | 30 | 1 | 2 | 4 | 7 | 11 | 15 | 29 |
| 35-49 | 1836 | 6.1 | 27 | 1 | 2 | 5 | 7 | 11 | 15 | 30 |
| 50-64 | 2648 | 6.9 | 45 | 1 | 2 | 5 | 8 | 13 | 18 | 32 |
| 65+ | 3861 | 8.8 | 63 | 2 | 3 | 6 | 11 | 17 | 24 | 42 |
| **GRAND TOTAL** | **9981** | **7.3** | **48** | **1** | **2** | **5** | **9** | **14** | **20** | **37** |

## 610: Incision or excision of anal fistula (49.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 28 | 1.9 | 2 | <1 | <1 | 2 | 2 | 3 | 4 | 8 |
| 20-34 | 264 | 2.8 | 2 | 1 | 1 | 3 | 4 | 4 | 5 | 9 |
| 35-49 | 254 | 2.8 | 2 | 1 | 1 | 3 | 3 | 4 | 5 | 7 |
| 50-64 | 171 | 3.0 | 2 | 1 | 1 | 3 | 3 | 5 | 5 | 10 |
| 65+ | 62 | 3.2 | 1 | 1 | 2 | 3 | 4 | 5 | 5 | 8 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 3.9 | 12 | 2 | 2 | 2 | 4 | 10 | 13 | 13 |
| 20-34 | 169 | 3.4 | 8 | 1 | 1 | 3 | 4 | 6 | 8 | 16 |
| 35-49 | 219 | 4.1 | 20 | 1 | 2 | 3 | 5 | 6 | 8 | 22 |
| 50-64 | 179 | 3.9 | 7 | 1 | 2 | 3 | 4 | 6 | 9 | 19 |
| 65+ | 79 | 4.9 | 10 | 1 | 2 | 4 | 6 | 9 | 13 | 14 |
| **TOT. SINGLE DX** | 779 | 2.8 | 3 | 1 | 1 | 3 | 3 | 4 | 5 | 9 |
| **" MULTIPLE DX** | 656 | 4.0 | 13 | 1 | 1 | 3 | 4 | 7 | 9 | 17 |
| TOTAL 0-19 YRS | 38 | 2.4 | 5 | <1 | 1 | 2 | 2 | 4 | 8 | 13 |
| 20-34 | 433 | 3.0 | 5 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 35-49 | 473 | 3.4 | 11 | 1 | 1 | 3 | 4 | 5 | 7 | 14 |
| 50-64 | 350 | 3.5 | 6 | 1 | 2 | 3 | 4 | 5 | 7 | 14 |
| 65+ | 141 | 4.2 | 7 | 1 | 2 | 3 | 5 | 8 | 9 | 14 |
| **GRAND TOTAL** | **1435** | **3.3** | **7** | **1** | **1** | **3** | **4** | **5** | **7** | **14** |

## 611: Local excision or destruction of lesion or tissue of anus (49.23; 49.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 32 | 1.8 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 4 |
| 20-34 | 229 | 2.3 | 2 | <1 | <1 | 2 | 3 | 4 | 5 | 6 |
| 35-49 | 128 | 2.7 | 2 | <1 | 1 | 2 | 3 | 4 | 6 | 9 |
| 50-64 | 67 | 3.2 | 5 | 1 | 1 | 3 | 4 | 5 | 6 | 14 |
| 65+ | 43 | 3.4 | 2 | 1 | 1 | 4 | 4 | 5 | 6 | 8 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 20-34 | 156 | 3.5 | 5 | 1 | 1 | 3 | 4 | 6 | 8 | 13 |
| 35-49 | 148 | 4.7 | 14 | 1 | 2 | 4 | 6 | 9 | 11 | 15 |
| 50-64 | 141 | 4.0 | 5 | 1 | 1 | 4 | 5 | 7 | 9 | 10 |
| 65+ | 92 | 6.2 | 26 | 2 | 2 | 5 | 7 | 13 | 15 | 30 |
| **TOT. SINGLE DX** | 499 | 2.6 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 8 |
| **" MULTIPLE DX** | 547 | 4.4 | 12 | 1 | 1 | 4 | 5 | 8 | 11 | 15 |
| TOTAL 0-19 YRS | 42 | 1.9 | 1 | <1 | <1 | 2 | 3 | 3 | 4 | 4 |
| 20-34 | 385 | 2.8 | 4 | <1 | 1 | 2 | 4 | 5 | 6 | 10 |
| 35-49 | 276 | 3.8 | 10 | 1 | 1 | 3 | 4 | 6 | 9 | 14 |
| 50-64 | 208 | 3.7 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 13 |
| 65+ | 135 | 5.3 | 21 | 1 | 2 | 4 | 6 | 10 | 14 | 27 |
| **GRAND TOTAL** | **1046** | **3.5** | **9** | **<1** | **1** | **3** | **4** | **6** | **8** | **15** |

## 612: Hemorrhoidectomy (49.43-49.46)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 3.8 | 5 | 2 | 2 | 3 | 5 | 7 | 7 | 7 |
| 20-34 | 215 | 4.0 | 3 | 2 | 2 | 4 | 5 | 6 | 8 | 9 |
| 35-49 | 341 | 4.2 | 3 | 2 | 2 | 4 | 5 | 7 | 8 | 9 |
| 50-64 | 249 | 4.4 | 4 | 2 | 2 | 4 | 5 | 7 | 8 | 12 |
| 65+ | 115 | 5.2 | 8 | 2 | 3 | 4 | 6 | 8 | 10 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 4.1 | 4 | <1 | 1 | 4 | 6 | 7 | 8 | 8 |
| 20-34 | 745 | 4.5 | 6 | 2 | 2 | 4 | 5 | 7 | 8 | 14 |
| 35-49 | 1239 | 4.7 | 6 | 2 | 3 | 4 | 5 | 7 | 9 | 14 |
| 50-64 | 1040 | 5.1 | 7 | 2 | 3 | 4 | 6 | 8 | 10 | 16 |
| 65+ | 513 | 6.8 | 33 | 3 | 3 | 5 | 7 | 12 | 17 | 31 |
| **TOT. SINGLE DX** | 924 | 4.3 | 4 | 2 | 2 | 4 | 5 | 7 | 8 | 10 |
| **" MULTIPLE DX** | 3556 | 5.1 | 11 | 2 | 3 | 4 | 6 | 8 | 10 | 18 |
| TOTAL 0-19 YRS | 23 | 4.0 | 5 | <1 | 1 | 3 | 6 | 7 | 8 | 8 |
| 20-34 | 960 | 4.4 | 6 | 2 | 2 | 4 | 5 | 7 | 8 | 12 |
| 35-49 | 1580 | 4.6 | 5 | 2 | 2 | 4 | 5 | 7 | 8 | 12 |
| 50-64 | 1289 | 4.9 | 7 | 2 | 3 | 4 | 6 | 8 | 10 | 16 |
| 65+ | 628 | 6.5 | 29 | 3 | 3 | 5 | 7 | 11 | 15 | 28 |
| **GRAND TOTAL** | **4480** | **4.9** | **10** | **2** | **3** | **4** | **6** | **8** | **10** | **17** |

## 613: Miscellaneous operations on anus (49.0; 49.22; 49.47-49.49; 49.5-49.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 31 | 2.1 | 1 | <1 | 1 | 2 | 3 | 4 | 4 | 7 |
| 20-34 | 204 | 2.7 | 3 | <1 | 1 | 2 | 3 | 5 | 6 | 10 |
| 35-49 | 192 | 2.8 | 3 | 1 | 1 | 2 | 4 | 5 | 7 | 10 |
| 50-64 | 121 | 3.2 | 5 | 1 | 1 | 3 | 4 | 6 | 8 | 13 |
| 65+ | 76 | 4.0 | 9 | 1 | 1 | 3 | 5 | 7 | 8 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 26 | 5.8 | 34 | 1 | 1 | 4 | 7 | 13 | 18 | 29 |
| 20-34 | 148 | 4.4 | 37 | 1 | 1 | 3 | 5 | 8 | 11 | 29 |
| 35-49 | 176 | 4.1 | 10 | 1 | 1 | 3 | 5 | 7 | 9 | 18 |
| 50-64 | 157 | 5.4 | 32 | 1 | 1 | 4 | 6 | 11 | 14 | 24 |
| 65+ | 157 | 8.7 | 86 | 1 | 2 | 6 | 11 | 19 | 25 | 45 |
| **TOT. SINGLE DX** | 624 | 2.9 | 5 | 1 | 1 | 2 | 4 | 6 | 7 | 12 |
| **" MULTIPLE DX** | 664 | 5.6 | 44 | 1 | 1 | 4 | 6 | 11 | 17 | 30 |
| TOTAL 0-19 YRS | 57 | 3.8 | 20 | 1 | 1 | 3 | 4 | 7 | 12 | 28 |
| 20-34 | 352 | 3.4 | 19 | 1 | 1 | 2 | 4 | 6 | 8 | 19 |
| 35-49 | 368 | 3.4 | 7 | 1 | 1 | 3 | 4 | 7 | 8 | 13 |
| 50-64 | 278 | 4.4 | 22 | 1 | 1 | 3 | 5 | 8 | 12 | 22 |
| 65+ | 233 | 7.2 | 66 | 1 | 2 | 5 | 8 | 16 | 21 | 43 |
| **GRAND TOTAL** | **1288** | **4.3** | **27** | **1** | **1** | **3** | **5** | **8** | **12** | **25** |

## 614: Incision of liver (50.0; 50.91)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 9.0 | | 9 | 9 | 9 | 9 | 9 | 9 | 9 |
| 20-34 | 13 | 11.4 | 69 | 2 | 4 | 9 | 15 | 24 | 30 | 32 |
| 35-49 | 6 | 3.5 | 10 | 1 | 1 | 2 | 6 | 9 | 9 | 9 |
| 50-64 | 7 | 7.7 | 61 | <1 | <1 | 6 | 11 | 21 | 23 | 23 |
| 65+ | 4 | 8.3 | 16 | 5 | 5 | 7 | 11 | 14 | 14 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 11.8 | 210 | 2 | 2 | 6 | 21 | 33 | 33 | 33 |
| 20-34 | 15 | 12.2 | 87 | 2 | 4 | 10 | 16 | 23 | 33 | 37 |
| 35-49 | 12 | 16.8 | 185 | 2 | 5 | 14 | 22 | 32 | 51 | 54 |
| 50-64 | 25 | 18.0 | 144 | 6 | 6 | 16 | 26 | 39 | 41 | 43 |
| 65+ | 23 | 16.4 | 151 | 2 | 4 | 15 | 19 | 35 | 43 | 54 |
| **TOT. SINGLE DX** | 31 | 8.5 | 52 | 1 | 1 | 6 | 11 | 19 | 23 | 32 |
| **" MULTIPLE DX** | 79 | 15.9 | 142 | 2 | 4 | 13 | 22 | 35 | 40 | 54 |
| TOTAL 0-19 YRS | 5 | 11.2 | 159 | 2 | 2 | 9 | 15 | 33 | 33 | 33 |
| 20-34 | 28 | 11.8 | 77 | 2 | 4 | 9 | 15 | 23 | 32 | 37 |
| 35-49 | 18 | 12.3 | 165 | 1 | 1 | 8 | 18 | 23 | 42 | 54 |
| 50-64 | 32 | 15.8 | 143 | 2 | 5 | 11 | 23 | 36 | 41 | 43 |
| 65+ | 27 | 15.2 | 138 | 2 | 4 | 13 | 17 | 33 | 40 | 54 |
| **GRAND TOTAL** | **110** | **13.9** | **127** | **2** | **2** | **10** | **18** | **31** | **37** | **54** |

## 615: Miscellaneous operations on liver (50.11-50.12; 50.2-50.6; 50.92-50.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 36 | 4.3 | 16 | 1 | 1 | 2 | 8 | 11 | 12 | 15 |
| 20-34 | 179 | 3.1 | 28 | <1 | 1 | 1 | 3 | 7 | 11 | 28 |
| 35-49 | 206 | 3.7 | 24 | <1 | 1 | 2 | 5 | 9 | 12 | 19 |
| 50-64 | 199 | 3.5 | 19 | 1 | 1 | 2 | 4 | 9 | 13 | 19 |
| 65+ | 128 | 4.8 | 32 | 1 | 1 | 2 | 6 | 14 | 19 | 23 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 89 | 12.2 | 154 | 1 | 2 | 8 | 16 | 29 | 46 | 86 |
| 20-34 | 282 | 9.0 | 89 | 1 | 1 | 7 | 11 | 17 | 32 | 54 |
| 35-49 | 392 | 8.6 | 87 | 1 | 1 | 6 | 11 | 18 | 26 | 46 |
| 50-64 | 723 | 9.2 | 90 | 1 | 1 | 7 | 12 | 21 | 26 | 43 |
| 65+ | 774 | 10.9 | 95 | 1 | 1 | 9 | 15 | 23 | 29 | 49 |
| **TOT. SINGLE DX** | 748 | 3.7 | 25 | <1 | 1 | 2 | 4 | 10 | 13 | 23 |
| **" MULTIPLE DX** | 2260 | 9.8 | 95 | 1 | 1 | 7 | 13 | 21 | 28 | 49 |
| TOTAL 0-19 YRS | 125 | 9.9 | 127 | 1 | 1 | 7 | 11 | 26 | 36 | 73 |
| 20-34 | 461 | 6.7 | 74 | 1 | 1 | 4 | 9 | 14 | 25 | 46 |
| 35-49 | 598 | 6.9 | 71 | 1 | 1 | 4 | 9 | 15 | 22 | 45 |
| 50-64 | 922 | 8.0 | 81 | 1 | 1 | 5 | 11 | 19 | 25 | 40 |
| 65+ | 902 | 10.1 | 91 | 1 | 1 | 8 | 14 | 22 | 28 | 49 |
| **GRAND TOTAL** | **3008** | **8.3** | **85** | **1** | **1** | **6** | **11** | **19** | **26** | **47** |

## 616: Cholecystectomy (51.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 214 | 6.1 | 8 | 3 | 4 | 6 | 7 | 8 | 10 | 17 |
| 20-34 | 2388 | 6.0 | 4 | 4 | 4 | 6 | 7 | 8 | 10 | 13 |
| 35-49 | 2186 | 6.4 | 6 | 4 | 4 | 6 | 7 | 9 | 10 | 15 |
| 50-64 | 2072 | 6.8 | 6 | 4 | 5 | 6 | 8 | 10 | 12 | 16 |
| 65+ | 1282 | 8.4 | 14 | 5 | 5 | 8 | 9 | 12 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 153 | 7.5 | 10 | 4 | 4 | 7 | 9 | 12 | 14 | 19 |
| 20-34 | 1789 | 7.6 | 14 | 4 | 5 | 7 | 8 | 12 | 14 | 22 |
| 35-49 | 2324 | 8.3 | 33 | 4 | 5 | 7 | 9 | 13 | 17 | 28 |
| 50-64 | 3172 | 9.3 | 35 | 5 | 5 | 8 | 10 | 15 | 19 | 34 |
| 65+ | 3860 | 12.7 | 71 | 6 | 6 | 10 | 15 | 21 | 29 | 49 |
| **TOT. SINGLE DX** | 8142 | 6.7 | 8 | 4 | 4 | 6 | 8 | 10 | 11 | 17 |
| **" MULTIPLE DX** | 11298 | 10.0 | 48 | 5 | 5 | 8 | 11 | 17 | 21 | 40 |
| TOTAL 0-19 YRS | 367 | 6.7 | 10 | 4 | 4 | 6 | 8 | 10 | 12 | 17 |
| 20-34 | 4177 | 6.7 | 9 | 4 | 4 | 6 | 7 | 10 | 12 | 18 |
| 35-49 | 4510 | 7.4 | 22 | 4 | 4 | 6 | 8 | 11 | 14 | 24 |
| 50-64 | 5244 | 8.3 | 25 | 4 | 5 | 7 | 9 | 13 | 16 | 28 |
| 65+ | 5142 | 11.6 | 60 | 5 | 6 | 9 | 13 | 20 | 25 | 46 |
| **GRAND TOTAL** | **19440** | **8.6** | **34** | **4** | **5** | **7** | **9** | **14** | **18** | **33** |

## 617: Operations on biliary tract and gallbladder except cholecystectomy (51.01-51.13; 51.3-51.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 8.4 | 57 | 1 | 1 | 7 | 13 | 20 | 20 | 20 |
| 20-34 | 33 | 8.7 | 35 | 2 | 3 | 8 | 12 | 15 | 23 | 28 |
| 35-49 | 40 | 8.5 | 34 | 1 | 2 | 8 | 11 | 17 | 19 | 26 |
| 50-64 | 53 | 9.2 | 34 | 2 | 2 | 9 | 13 | 17 | 18 | 26 |
| 65+ | 98 | 12.4 | 62 | 1 | 3 | 11 | 16 | 23 | 29 | 36 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 9 | 11.1 | 24 | 4 | 5 | 10 | 15 | 18 | 20 | 20 |
| 20-34 | 34 | 12.2 | 94 | 2 | 3 | 10 | 17 | 21 | 29 | 51 |
| 35-49 | 65 | 14.6 | 130 | 2 | 5 | 11 | 19 | 31 | 35 | 66 |
| 50-64 | 184 | 16.9 | 167 | 4 | 6 | 13 | 21 | 32 | 42 | 70 |
| 65+ | 472 | 18.1 | 162 | 4 | 7 | 15 | 22 | 31 | 43 | 76 |
| **TOT. SINGLE DX** | 229 | 10.4 | 50 | 1 | 2 | 9 | 14 | 20 | 24 | 33 |
| **" MULTIPLE DX** | 764 | 17.2 | 158 | 4 | 6 | 14 | 21 | 31 | 41 | 69 |
| TOTAL 0-19 YRS | 14 | 10.1 | 34 | 1 | 3 | 9 | 15 | 20 | 20 | 20 |
| 20-34 | 67 | 10.5 | 67 | 2 | 3 | 8 | 14 | 20 | 25 | 48 |
| 35-49 | 105 | 12.3 | 102 | 2 | 2 | 10 | 16 | 25 | 32 | 55 |
| 50-64 | 237 | 15.2 | 148 | 3 | 4 | 11 | 19 | 29 | 40 | 61 |
| 65+ | 570 | 17.1 | 149 | 3 | 6 | 14 | 22 | 30 | 38 | 65 |
| **GRAND TOTAL** | **993** | **15.6** | **141** | **3** | **4** | **13** | **20** | **29** | **37** | **60** |

## 618: Operations on pancreas (52.01-52.12; 52.2-52.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 6.9 | 7 | 3 | 3 | 7 | 9 | 10 | 10 | 10 |
| 20-34 | 44 | 6.2 | 29 | 1 | 2 | 4 | 9 | 14 | 18 | 24 |
| 35-49 | 78 | 6.2 | 35 | <1 | 1 | 4 | 10 | 14 | 17 | 31 |
| 50-64 | 87 | 6.2 | 35 | <1 | 1 | 4 | 9 | 16 | 19 | 25 |
| 65+ | 66 | 10.6 | 98 | 1 | 1 | 8 | 14 | 20 | 35 | 45 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 17.2 | 292 | 2 | 2 | 13 | 26 | 36 | 57 | 62 |
| 20-34 | 68 | 13.0 | 150 | 2 | 2 | 11 | 15 | 25 | 35 | 65 |
| 35-49 | 149 | 15.5 | 255 | 1 | 2 | 10 | 19 | 33 | 45 | 78 |
| 50-64 | 213 | 14.6 | 189 | 2 | 3 | 11 | 17 | 30 | 42 | 74 |
| 65+ | 289 | 15.0 | 190 | 1 | 2 | 11 | 19 | 34 | 45 | 68 |
| **TOT. SINGLE DX** | 282 | 7.3 | 51 | 1 | 1 | 5 | 10 | 16 | 19 | 36 |
| **" MULTIPLE DX** | 732 | 14.8 | 201 | 1 | 2 | 11 | 19 | 32 | 45 | 70 |
| TOTAL 0-19 YRS | 20 | 13.6 | 212 | 2 | 2 | 8 | 21 | 28 | 46 | 62 |
| 20-34 | 112 | 10.3 | 113 | 1 | 2 | 8 | 13 | 22 | 27 | 60 |
| 35-49 | 227 | 12.3 | 199 | 1 | 1 | 8 | 15 | 28 | 35 | 76 |
| 50-64 | 300 | 12.2 | 159 | 1 | 1 | 9 | 16 | 25 | 37 | 69 |
| 65+ | 355 | 14.2 | 176 | 1 | 2 | 10 | 18 | 32 | 44 | 67 |
| **GRAND TOTAL** | **1014** | **12.7** | **171** | **1** | **2** | **9** | **16** | **28** | **40** | **68** |

## 619: Miscellaneous diagnostic and therapeutic procedures on digestive tract, abdominal cavity and abdominal wall (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 125 | 4.7 | 27 | <1 | 1 | 3 | 5 | 10 | 12 | 34 |
| 20-34 | 394 | 4.2 | 9 | 1 | 1 | 3 | 5 | 8 | 10 | 18 |
| 35-49 | 324 | 4.8 | 22 | <1 | 1 | 4 | 6 | 9 | 14 | 28 |
| 50-64 | 373 | 4.1 | 12 | <1 | <1 | 3 | 5 | 8 | 10 | 22 |
| 65+ | 461 | 5.0 | 18 | 1 | 1 | 4 | 6 | 10 | 13 | 20 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 115 | 5.4 | 37 | 1 | 1 | 4 | 7 | 10 | 14 | 39 |
| 20-34 | 599 | 6.0 | 25 | 1 | 2 | 5 | 7 | 12 | 16 | 26 |
| 35-49 | 803 | 6.5 | 37 | 1 | 2 | 5 | 8 | 12 | 18 | 34 |
| 50-64 | 1459 | 7.1 | 47 | 1 | 2 | 5 | 8 | 14 | 19 | 37 |
| 65+ | 3469 | 8.2 | 56 | 2 | 2 | 6 | 10 | 16 | 21 | 39 |
| **TOT. SINGLE DX** | 1677 | 4.6 | 17 | <1 | 1 | 4 | 6 | 9 | 11 | 22 |
| **" MULTIPLE DX** | 6445 | 7.5 | 50 | 1 | 2 | 6 | 9 | 15 | 20 | 36 |
| TOTAL 0-19 YRS | 240 | 5.1 | 32 | <1 | 1 | 4 | 6 | 10 | 13 | 34 |
| 20-34 | 993 | 5.3 | 20 | 1 | 2 | 4 | 6 | 10 | 14 | 23 |
| 35-49 | 1127 | 6.0 | 34 | 1 | 2 | 5 | 7 | 11 | 16 | 32 |
| 50-64 | 1832 | 6.4 | 42 | 1 | 1 | 5 | 8 | 13 | 18 | 34 |
| 65+ | 3930 | 7.9 | 53 | 1 | 2 | 6 | 10 | 15 | 20 | 37 |
| **GRAND TOTAL** | **8122** | **6.9** | **44** | **1** | **2** | **5** | **8** | **14** | **18** | **34** |

## 620: Unilateral repair of inguinal hernia (53.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1913 | 1.7 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| 20-34 | 2213 | 2.6 | 1 | 1 | 1 | 3 | 3 | 4 | 5 | 6 |
| 35-49 | 1950 | 2.8 | 1 | 1 | 1 | 3 | 4 | 4 | 5 | 6 |
| 50-64 | 2773 | 3.0 | 1 | 1 | 1 | 3 | 4 | 5 | 5 | 8 |
| 65+ | 2624 | 3.4 | 2 | 1 | 2 | 3 | 4 | 5 | 6 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 869 | 2.1 | 21 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| 20-34 | 666 | 3.3 | 4 | 1 | 2 | 3 | 4 | 5 | 6 | 12 |
| 35-49 | 844 | 3.5 | 6 | 1 | 2 | 3 | 4 | 5 | 7 | 12 |
| 50-64 | 1824 | 4.0 | 8 | 1 | 2 | 3 | 5 | 6 | 8 | 15 |
| 65+ | 2705 | 5.3 | 19 | 2 | 2 | 4 | 6 | 10 | 13 | 24 |
| **TOT. SINGLE DX** | 11473 | 2.8 | 2 | <1 | 1 | 3 | 4 | 4 | 5 | 7 |
| **" MULTIPLE DX** | 6908 | 4.1 | 15 | 1 | 1 | 3 | 5 | 7 | 10 | 19 |
| TOTAL 0-19 YRS | 2782 | 1.8 | 8 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 20-34 | 2879 | 2.8 | 2 | 1 | 1 | 3 | 3 | 4 | 5 | 7 |
| 35-49 | 2794 | 3.0 | 3 | 1 | 1 | 3 | 4 | 5 | 5 | 8 |
| 50-64 | 4597 | 3.4 | 5 | 1 | 1 | 3 | 4 | 5 | 7 | 11 |
| 65+ | 5329 | 4.4 | 12 | 1 | 2 | 4 | 5 | 7 | 10 | 19 |
| **GRAND TOTAL** | **18381** | **3.3** | **7** | **1** | **1** | **3** | **4** | **5** | **7** | **13** |

## 621: Bilateral repair of inguinal hernia (53.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 358 | 1.5 | | <1 | <1 | 1 | 2 | 3 | 3 | 6 |
| 20-34 | 206 | 3.2 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 7 |
| 35-49 | 300 | 3.3 | 1 | 1 | 2 | 3 | 4 | 5 | 5 | 7 |
| 50-64 | 468 | 3.6 | 1 | 1 | 2 | 4 | 4 | 5 | 6 | 8 |
| 65+ | 373 | 4.0 | 12 | 2 | 2 | 4 | 5 | 6 | 7 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 220 | 2.9 | 66 | <1 | <1 | 1 | 2 | 3 | 6 | 69 |
| 20-34 | 89 | 3.7 | 2 | 1 | 2 | 4 | 5 | 6 | 7 | 8 |
| 35-49 | 169 | 4.2 | 9 | 2 | 2 | 4 | 5 | 6 | 7 | 19 |
| 50-64 | 354 | 4.4 | 8 | 2 | 2 | 4 | 5 | 7 | 9 | 14 |
| 65+ | 455 | 6.1 | 25 | 2 | 3 | 5 | 7 | 10 | 15 | 32 |
| **TOT. SINGLE DX** | 1705 | 3.2 | 5 | <1 | 1 | 3 | 4 | 5 | 6 | 8 |
| **" MULTIPLE DX** | 1287 | 4.7 | 25 | <1 | 1 | 4 | 5 | 8 | 10 | 28 |
| TOTAL 0-19 YRS | 578 | 2.0 | 26 | <1 | <1 | 1 | 2 | 3 | 4 | 21 |
| 20-34 | 295 | 3.4 | 2 | 1 | 2 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 469 | 3.6 | 4 | 1 | 2 | 3 | 4 | 5 | 6 | 9 |
| 50-64 | 822 | 4.0 | 5 | 1 | 2 | 4 | 5 | 6 | 7 | 11 |
| 65+ | 828 | 5.2 | 21 | 2 | 2 | 4 | 6 | 8 | 10 | 25 |
| **GRAND TOTAL** | **2992** | **3.8** | **14** | **<1** | **1** | **3** | **5** | **6** | **8** | **17** |

## 622: Unilateral or bilateral repair of femoral hernia (53.2-53.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 2.4 | 1 | <1 | 1 | 2 | 3 | 4 | 5 | 5 |
| 20-34 | 71 | 2.8 | 1 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 89 | 3.0 | 1 | 1 | 1 | 3 | 4 | 5 | 5 | 8 |
| 50-64 | 121 | 3.5 | 2 | 1 | 1 | 3 | 4 | 6 | 7 | 8 |
| 65+ | 139 | 4.2 | 4 | 1 | 2 | 4 | 5 | 7 | 9 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 3.4 | | 2 | 2 | 3 | 4 | 5 | 5 | 5 |
| 20-34 | 21 | 3.8 | 3 | 1 | 1 | 4 | 5 | 6 | 7 | 8 |
| 35-49 | 48 | 5.0 | 9 | 2 | 2 | 4 | 6 | 10 | 11 | 14 |
| 50-64 | 88 | 4.8 | 14 | 1 | 2 | 4 | 6 | 8 | 14 | 20 |
| 65+ | 214 | 8.1 | 40 | 3 | 3 | 6 | 10 | 14 | 20 | 36 |
| **TOT. SINGLE DX** | 439 | 3.5 | 3 | 1 | 1 | 3 | 4 | 6 | 7 | 9 |
| **" MULTIPLE DX** | 379 | 6.6 | 30 | 2 | 2 | 5 | 8 | 13 | 18 | 31 |
| TOTAL 0-19 YRS | 27 | 2.7 | 1 | 1 | 1 | 3 | 3 | 4 | 5 | 5 |
| 20-34 | 92 | 3.1 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 137 | 3.7 | 5 | 1 | 2 | 3 | 4 | 6 | 9 | 12 |
| 50-64 | 209 | 4.0 | 8 | 1 | 2 | 3 | 5 | 7 | 8 | 18 |
| 65+ | 353 | 6.6 | 30 | 2 | 3 | 5 | 8 | 12 | 17 | 31 |
| **GRAND TOTAL** | **818** | **4.9** | **18** | **1** | **2** | **4** | **6** | **9** | **13** | **21** |

## 623: Repair of umbilical hernia (53.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 236 | 1.3 | | <1 | <1 | 1 | 2 | 2 | 2 | 3 |
| 20-34 | 174 | 2.2 | 1 | <1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 35-49 | 287 | 2.6 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 7 |
| 50-64 | 205 | 2.9 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 65+ | 91 | 2.9 | 2 | <1 | 1 | 3 | 4 | 5 | 6 | 8 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 60 | 3.7 | 50 | <1 | <1 | 2 | 3 | 6 | 17 | 41 |
| 20-34 | 73 | 3.4 | 2 | 1 | 1 | 3 | 4 | 6 | 6 | 7 |
| 35-49 | 177 | 3.8 | 9 | 1 | 1 | 3 | 5 | 6 | 8 | 15 |
| 50-64 | 219 | 4.7 | 9 | 1 | 2 | 4 | 6 | 9 | 11 | 17 |
| 65+ | 182 | 6.9 | 37 | 2 | 2 | 5 | 8 | 13 | 17 | 32 |
| **TOT. SINGLE DX** | 993 | 2.3 | 2 | <1 | <1 | 2 | 3 | 4 | 5 | 8 |
| **" MULTIPLE DX** | 711 | 4.8 | 21 | 1 | 1 | 4 | 6 | 9 | 12 | 27 |
| TOTAL 0-19 YRS | 296 | 1.8 | 11 | <1 | <1 | 1 | 2 | 2 | 3 | 18 |
| 20-34 | 247 | 2.6 | 2 | <1 | 1 | 2 | 3 | 5 | 6 | 7 |
| 35-49 | 464 | 3.1 | 5 | 1 | 1 | 3 | 4 | 5 | 6 | 11 |
| 50-64 | 424 | 3.8 | 7 | 1 | 1 | 3 | 5 | 7 | 9 | 14 |
| 65+ | 273 | 5.5 | 29 | 1 | 2 | 4 | 7 | 11 | 14 | 29 |
| **GRAND TOTAL** | **1704** | **3.4** | **12** | **<1** | **1** | **2** | **4** | **6** | **8** | **16** |

## 624: Repair of other hernia of anterior abdominal wall (53.5-53.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 83 | 1.9 | 1 | <1 | <1 | 1 | 3 | 3 | 4 | 8 |
| 20-34 | 282 | 3.3 | 3 | 1 | 1 | 3 | 4 | 6 | 7 | 10 |
| 35-49 | 495 | 4.0 | 4 | 1 | 2 | 4 | 5 | 7 | 8 | 10 |
| 50-64 | 569 | 4.6 | 5 | 2 | 2 | 4 | 6 | 8 | 9 | 13 |
| 65+ | 401 | 5.1 | 5 | 2 | 2 | 5 | 6 | 8 | 10 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 24 | 3.2 | 6 | <1 | <1 | 2 | 5 | 8 | 8 | 8 |
| 20-34 | 153 | 5.6 | 22 | 1 | 2 | 5 | 6 | 10 | 15 | 26 |
| 35-49 | 372 | 6.9 | 54 | 2 | 3 | 5 | 7 | 11 | 15 | 39 |
| 50-64 | 600 | 7.1 | 30 | 2 | 3 | 6 | 9 | 12 | 16 | 26 |
| 65+ | 660 | 8.4 | 51 | 2 | 3 | 7 | 10 | 15 | 21 | 40 |
| **TOT. SINGLE DX** | 1830 | 4.2 | 5 | 1 | 2 | 4 | 6 | 7 | 8 | 11 |
| **" MULTIPLE DX** | 1809 | 7.4 | 43 | 2 | 3 | 6 | 9 | 13 | 17 | 31 |
| TOTAL 0-19 YRS | 107 | 2.2 | 3 | <1 | <1 | 2 | 3 | 4 | 7 | 8 |
| 20-34 | 435 | 4.1 | 12 | 1 | 1 | 3 | 5 | 7 | 9 | 20 |
| 35-49 | 867 | 5.2 | 28 | 1 | 2 | 4 | 6 | 9 | 11 | 24 |
| 50-64 | 1169 | 5.9 | 20 | 2 | 2 | 5 | 7 | 10 | 13 | 21 |
| 65+ | 1061 | 7.2 | 37 | 2 | 3 | 6 | 8 | 12 | 17 | 28 |
| **GRAND TOTAL** | **3639** | **5.8** | **27** | **1** | **2** | **5** | **7** | **10** | **13** | **25** |

## 625: Repair of diaphragmatic and other hernia (53.7-53.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 4.7 | 75 | <1 | <1 | 2 | 4 | 11 | 29 | 34 |
| 20-34 | 19 | 4.8 | 5 | 2 | 2 | 4 | 7 | 8 | 9 | 9 |
| 35-49 | 38 | 5.7 | 6 | 2 | 2 | 5 | 8 | 9 | 10 | 12 |
| 50-64 | 47 | 7.2 | 21 | 2 | 3 | 7 | 8 | 11 | 14 | 32 |
| 65+ | 25 | 9.8 | 108 | 2 | 3 | 8 | 10 | 14 | 25 | 57 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 18 | 13.3 | 155 | 1 | 1 | 8 | 20 | 36 | 41 | 43 |
| 20-34 | 48 | 8.8 | 19 | 4 | 5 | 8 | 10 | 14 | 16 | 25 |
| 35-49 | 101 | 9.7 | 58 | 4 | 5 | 8 | 10 | 16 | 20 | 50 |
| 50-64 | 148 | 10.3 | 43 | 5 | 6 | 9 | 11 | 16 | 20 | 39 |
| 65+ | 176 | 12.2 | 65 | 5 | 6 | 10 | 14 | 20 | 25 | 56 |
| **TOT. SINGLE DX** | 143 | 6.7 | 38 | 1 | 2 | 6 | 8 | 10 | 14 | 36 |
| **" MULTIPLE DX** | 491 | 10.8 | 57 | 4 | 5 | 9 | 12 | 18 | 23 | 47 |
| TOTAL 0-19 YRS | 32 | 9.5 | 135 | <1 | 1 | 6 | 10 | 32 | 38 | 43 |
| 20-34 | 67 | 7.7 | 18 | 2 | 3 | 7 | 9 | 13 | 15 | 25 |
| 35-49 | 139 | 8.6 | 47 | 2 | 4 | 8 | 9 | 13 | 19 | 37 |
| 50-64 | 195 | 9.6 | 40 | 3 | 5 | 8 | 11 | 15 | 19 | 39 |
| 65+ | 201 | 11.9 | 71 | 4 | 6 | 10 | 14 | 19 | 25 | 58 |
| **GRAND TOTAL** | **634** | **9.9** | **56** | **2** | **4** | **8** | **11** | **17** | **22** | **46** |

## 626: Incision, excision of abdominal wall and peritoneum except laparotomy (54.0; 54.22-54.23; 54.3-54.4; 54.91)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 112 | 3.3 | 13 | <1 | <1 | 2 | 4 | 7 | 10 | 21 |
| 20-34 | 209 | 4.0 | 10 | <1 | 1 | 3 | 6 | 8 | 9 | 14 |
| 35-49 | 144 | 4.1 | 16 | <1 | 1 | 3 | 5 | 8 | 10 | 26 |
| 50-64 | 159 | 6.6 | 55 | 1 | 1 | 6 | 8 | 12 | 14 | 51 |
| 65+ | 142 | 8.0 | 42 | 1 | 2 | 6 | 11 | 17 | 21 | 27 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 92 | 6.1 | 43 | <1 | 1 | 4 | 8 | 13 | 15 | 38 |
| 20-34 | 311 | 7.7 | 55 | 1 | 1 | 6 | 9 | 17 | 26 | 42 |
| 35-49 | 427 | 9.8 | 106 | <1 | 1 | 6 | 13 | 23 | 28 | 54 |
| 50-64 | 733 | 11.0 | 111 | 1 | 2 | 8 | 13 | 21 | 32 | 58 |
| 65+ | 842 | 13.2 | 123 | 2 | 3 | 10 | 18 | 26 | 34 | 61 |
| **TOT. SINGLE DX** | 766 | 5.2 | 30 | <1 | 1 | 4 | 7 | 11 | 14 | 26 |
| **" MULTIPLE DX** | 2405 | 10.9 | 109 | 1 | 2 | 8 | 14 | 24 | 30 | 54 |
| TOTAL 0-19 YRS | 204 | 4.5 | 29 | <1 | <1 | 3 | 6 | 10 | 13 | 29 |
| 20-34 | 520 | 6.3 | 40 | 1 | 1 | 5 | 8 | 12 | 19 | 33 |
| 35-49 | 571 | 8.4 | 90 | <1 | 1 | 5 | 10 | 20 | 27 | 51 |
| 50-64 | 892 | 10.2 | 104 | 1 | 2 | 7 | 12 | 21 | 27 | 56 |
| 65+ | 984 | 12.4 | 115 | 1 | 2 | 10 | 17 | 25 | 32 | 55 |
| **GRAND TOTAL** | **3171** | **9.5** | **96** | **1** | **1** | **7** | **12** | **21** | **28** | **50** |

## 627: Laparotomy (54.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 174 | 5.4 | 17 | 1 | 2 | 4 | 7 | 10 | 13 | 18 |
| 20-34 | 426 | 5.5 | 14 | 2 | 3 | 5 | 6 | 9 | 11 | 22 |
| 35-49 | 159 | 7.8 | 36 | 2 | 3 | 6 | 9 | 14 | 18 | 36 |
| 50-64 | 134 | 8.8 | 25 | 3 | 4 | 8 | 11 | 14 | 20 | 29 |
| 65+ | 106 | 11.0 | 47 | 4 | 5 | 10 | 13 | 18 | 21 | 42 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 251 | 9.9 | 150 | 2 | 3 | 6 | 10 | 18 | 34 | 69 |
| 20-34 | 756 | 8.3 | 85 | 2 | 3 | 6 | 9 | 14 | 22 | 54 |
| 35-49 | 399 | 10.7 | 88 | 3 | 4 | 8 | 12 | 21 | 29 | 64 |
| 50-64 | 422 | 15.3 | 197 | 4 | 5 | 10 | 18 | 32 | 40 | 85 |
| 65+ | 562 | 17.7 | 191 | 5 | 6 | 14 | 21 | 34 | 44 | 85 |
| **TOT. SINGLE DX** | 999 | 6.9 | 27 | 2 | 3 | 5 | 8 | 12 | 16 | 29 |
| **" MULTIPLE DX** | 2390 | 12.3 | 152 | 3 | 4 | 8 | 14 | 25 | 37 | 71 |
| TOTAL 0-19 YRS | 425 | 8.1 | 101 | 2 | 3 | 5 | 8 | 14 | 20 | 61 |
| 20-34 | 1182 | 7.3 | 62 | 2 | 3 | 5 | 8 | 12 | 18 | 49 |
| 35-49 | 558 | 9.8 | 75 | 3 | 4 | 7 | 12 | 18 | 27 | 54 |
| 50-64 | 556 | 13.7 | 163 | 4 | 5 | 10 | 16 | 27 | 37 | 81 |
| 65+ | 668 | 16.7 | 175 | 5 | 6 | 13 | 20 | 31 | 41 | 74 |
| **GRAND TOTAL** | **3389** | **10.7** | **121** | **3** | **3** | **7** | **12** | **21** | **31** | **63** |

## 628: Laparoscopy (excluding that for destruction of fallopian tubes) (54.21)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 310 | 2.3 | 3 | <1 | <1 | 1 | 3 | 5 | 6 | 10 |
| 20-34 | 1391 | 2.0 | 4 | <1 | <1 | 1 | 2 | 4 | 6 | 9 |
| 35-49 | 226 | 1.9 | 2 | <1 | <1 | 1 | 2 | 4 | 5 | 8 |
| 50-64 | 42 | 2.6 | 6 | <1 | <1 | 2 | 3 | 6 | 9 | 11 |
| 65+ | 11 | 4.3 | 26 | 1 | 1 | 2 | 4 | 13 | 17 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 230 | 3.3 | 8 | <1 | 1 | 2 | 4 | 7 | 9 | 12 |
| 20-34 | 1502 | 2.4 | 5 | <1 | <1 | 1 | 3 | 5 | 7 | 12 |
| 35-49 | 274 | 2.6 | 8 | <1 | <1 | 1 | 3 | 7 | 9 | 14 |
| 50-64 | 58 | 4.4 | 18 | <1 | 1 | 2 | 6 | 10 | 16 | 19 |
| 65+ | 44 | 10.5 | 144 | 1 | 1 | 7 | 13 | 21 | 42 | 55 |
| **TOT. SINGLE DX** | 1980 | 2.1 | 4 | <1 | <1 | 1 | 2 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 2108 | 2.7 | 11 | <1 | <1 | 1 | 3 | 6 | 8 | 15 |
| TOTAL 0-19 YRS | 540 | 2.7 | 6 | <1 | <1 | 2 | 4 | 6 | 7 | 11 |
| 20-34 | 2893 | 2.2 | 5 | <1 | <1 | 1 | 2 | 5 | 7 | 11 |
| 35-49 | 500 | 2.3 | 6 | <1 | <1 | 1 | 3 | 5 | 7 | 12 |
| 50-64 | 100 | 3.7 | 14 | <1 | <1 | 2 | 5 | 9 | 11 | 18 |
| 65+ | 55 | 9.2 | 126 | 1 | 1 | 6 | 11 | 19 | 39 | 54 |
| **GRAND TOTAL** | **4088** | **2.4** | **8** | **<1** | **<1** | **1** | **3** | **5** | **7** | **13** |

## 629: Peritoneal dialysis (54.98)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| TOT. SINGLE DX | | | | | | | | | | |
| " MULTIPLE DX | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| GRAND TOTAL | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 630: Miscellaneous operations on abdominal wall and peritoneum (54.5-54.7; 54.92-54.97; 54.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 81 | 6.4 | 17 | <1 | 1 | 6 | 8 | 11 | 15 | 21 |
| 20-34 | 287 | 5.5 | 17 | <1 | <1 | 5 | 7 | 10 | 13 | 20 |
| 35-49 | 222 | 7.5 | 20 | 1 | 3 | 7 | 9 | 12 | 14 | 26 |
| 50-64 | 201 | 9.1 | 18 | 4 | 5 | 8 | 10 | 14 | 17 | 25 |
| 65+ | 170 | 11.2 | 39 | 4 | 5 | 10 | 13 | 17 | 22 | 38 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 134 | 8.9 | 109 | 1 | 2 | 6 | 11 | 20 | 22 | 51 |
| 20-34 | 836 | 6.1 | 28 | 1 | 1 | 5 | 7 | 10 | 15 | 32 |
| 35-49 | 426 | 10.9 | 126 | 1 | 4 | 7 | 12 | 21 | 31 | 67 |
| 50-64 | 479 | 14.2 | 138 | 4 | 5 | 11 | 17 | 27 | 35 | 63 |
| 65+ | 636 | 17.2 | 165 | 5 | 7 | 14 | 20 | 33 | 45 | 77 |
| TOT. SINGLE DX | 961 | 7.8 | 27 | 1 | 2 | 7 | 10 | 13 | 17 | 26 |
| " MULTIPLE DX | 2511 | 11.4 | 125 | 1 | 3 | 8 | 14 | 24 | 32 | 63 |
| TOTAL 0-19 YRS | 215 | 8.0 | 76 | 1 | 2 | 6 | 9 | 17 | 21 | 36 |
| 20-34 | 1123 | 5.9 | 26 | <1 | 1 | 5 | 7 | 10 | 14 | 30 |
| 35-49 | 648 | 9.7 | 92 | 1 | 3 | 7 | 11 | 18 | 25 | 65 |
| 50-64 | 680 | 12.7 | 108 | 4 | 5 | 10 | 15 | 24 | 31 | 55 |
| 65+ | 806 | 15.9 | 145 | 5 | 6 | 12 | 18 | 30 | 41 | 70 |
| GRAND TOTAL | 3472 | 10.4 | 100 | 1 | 3 | 8 | 12 | 20 | 28 | 57 |

## 631: Complete nephrectomy (55.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 22 | 7.3 | 12 | 3 | 3 | 7 | 9 | 11 | 16 | 17 |
| 20-34 | 50 | 7.1 | 2 | 5 | 5 | 7 | 8 | 9 | 9 | 11 |
| 35-49 | 53 | 8.9 | 8 | 5 | 6 | 8 | 10 | 13 | 15 | 19 |
| 50-64 | 94 | 9.4 | 10 | 6 | 6 | 9 | 10 | 14 | 16 | 22 |
| 65+ | 89 | 11.9 | 32 | 6 | 7 | 10 | 14 | 19 | 23 | 34 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 66 | 10.6 | 97 | 3 | 4 | 8 | 11 | 17 | 35 | 60 |
| 20-34 | 107 | 12.7 | 135 | 5 | 6 | 9 | 13 | 22 | 36 | 69 |
| 35-49 | 111 | 13.4 | 128 | 6 | 6 | 9 | 14 | 24 | 38 | 70 |
| 50-64 | 239 | 13.5 | 82 | 6 | 7 | 11 | 16 | 22 | 31 | 50 |
| 65+ | 346 | 15.9 | 104 | 7 | 8 | 13 | 19 | 25 | 38 | 55 |
| TOT. SINGLE DX | 308 | 9.5 | 18 | 5 | 6 | 8 | 11 | 15 | 17 | 27 |
| " MULTIPLE DX | 869 | 14.1 | 107 | 6 | 6 | 11 | 16 | 24 | 35 | 55 |
| TOTAL 0-19 YRS | 88 | 9.8 | 78 | 3 | 4 | 8 | 10 | 16 | 27 | 54 |
| 20-34 | 157 | 10.9 | 99 | 5 | 5 | 8 | 11 | 18 | 27 | 64 |
| 35-49 | 164 | 11.9 | 94 | 6 | 6 | 9 | 12 | 20 | 31 | 62 |
| 50-64 | 333 | 12.3 | 66 | 6 | 7 | 10 | 14 | 20 | 27 | 48 |
| 65+ | 435 | 15.1 | 92 | 7 | 8 | 12 | 18 | 24 | 33 | 55 |
| GRAND TOTAL | 1177 | 12.9 | 88 | 6 | 6 | 10 | 15 | 22 | 29 | 55 |

## 632: Miscellaneous operations on kidney (55.0-55.1; 55.23-55.24; 55.3-55.4; 55.6-55.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 104 | 5.4 | 19 | 1 | 1 | 5 | 8 | 10 | 13 | 25 |
| 20-34 | 263 | 7.8 | 25 | 2 | 2 | 7 | 9 | 13 | 18 | 28 |
| 35-49 | 204 | 7.5 | 22 | 2 | 2 | 7 | 9 | 13 | 15 | 28 |
| 50-64 | 207 | 7.8 | 23 | 1 | 2 | 8 | 9 | 13 | 15 | 27 |
| 65+ | 92 | 10.1 | 56 | 1 | 3 | 9 | 11 | 19 | 22 | 47 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 157 | 10.3 | 63 | 2 | 3 | 8 | 12 | 20 | 26 | 48 |
| 20-34 | 415 | 12.5 | 136 | 2 | 2 | 9 | 15 | 29 | 39 | 59 |
| 35-49 | 369 | 13.0 | 119 | 2 | 2 | 10 | 16 | 26 | 35 | 59 |
| 50-64 | 398 | 11.8 | 108 | 2 | 2 | 9 | 14 | 23 | 32 | 54 |
| 65+ | 327 | 13.5 | 131 | 2 | 3 | 11 | 16 | 26 | 32 | 55 |
| TOT. SINGLE DX | 870 | 7.7 | 28 | 1 | 2 | 7 | 9 | 13 | 17 | 28 |
| " MULTIPLE DX | 1666 | 12.4 | 119 | 2 | 2 | 9 | 15 | 25 | 35 | 55 |
| TOTAL 0-19 YRS | 261 | 8.4 | 52 | 1 | 2 | 7 | 10 | 16 | 21 | 35 |
| 20-34 | 678 | 10.7 | 99 | 2 | 2 | 8 | 12 | 22 | 31 | 52 |
| 35-49 | 573 | 11.1 | 92 | 2 | 2 | 9 | 13 | 21 | 31 | 52 |
| 50-64 | 605 | 10.4 | 83 | 1 | 2 | 8 | 12 | 20 | 27 | 50 |
| 65+ | 419 | 12.7 | 116 | 2 | 3 | 10 | 16 | 24 | 30 | 51 |
| GRAND TOTAL | 2536 | 10.8 | 93 | 2 | 2 | 8 | 13 | 21 | 29 | 50 |

## 633: Ureterotomy (56.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 17 | 5.8 | 9 | 2 | 2 | 6 | 7 | 11 | 12 | 12 |
| 20-34 | 222 | 6.3 | 9 | 2 | 3 | 6 | 8 | 10 | 11 | 15 |
| 35-49 | 257 | 6.8 | 7 | 2 | 3 | 7 | 8 | 10 | 12 | 15 |
| 50-64 | 234 | 7.6 | 11 | 2 | 3 | 7 | 9 | 12 | 14 | 17 |
| 65+ | 93 | 8.2 | 8 | 3 | 5 | 8 | 10 | 12 | 14 | 16 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 8.9 | 179 | 1 | 2 | 5 | 11 | 48 | 85 | 99+ |
| 20-34 | 132 | 8.3 | 21 | 2 | 4 | 8 | 10 | 12 | 16 | 29 |
| 35-49 | 198 | 8.2 | 22 | 3 | 3 | 8 | 10 | 12 | 15 | 31 |
| 50-64 | 261 | 9.8 | 38 | 3 | 4 | 8 | 12 | 17 | 21 | 35 |
| 65+ | 236 | 12.3 | 90 | 3 | 5 | 10 | 14 | 22 | 29 | 58 |
| **TOT. SINGLE DX** | 823 | 7.0 | 10 | 2 | 3 | 7 | 9 | 11 | 13 | 16 |
| **" MULTIPLE DX** | 843 | 9.9 | 52 | 3 | 4 | 8 | 11 | 17 | 23 | 43 |
| TOTAL 0-19 YRS | 33 | 7.3 | 91 | 2 | 2 | 5 | 9 | 13 | 48 | 99+ |
| 20-34 | 354 | 7.0 | 15 | 2 | 3 | 7 | 9 | 11 | 13 | 23 |
| 35-49 | 455 | 7.4 | 15 | 2 | 3 | 7 | 9 | 11 | 13 | 24 |
| 50-64 | 495 | 8.7 | 27 | 2 | 4 | 8 | 10 | 14 | 18 | 27 |
| 65+ | 329 | 11.1 | 70 | 3 | 5 | 9 | 12 | 20 | 27 | 50 |
| **GRAND TOTAL** | **1666** | **8.5** | **33** | **2** | **3** | **8** | **10** | **13** | **18** | **31** |

## 634: Ureteral diversion (56.5-56.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 131 | 7.5 | 6 | 4 | 5 | 7 | 9 | 11 | 12 | 15 |
| 20-34 | 21 | 10.3 | 16 | 6 | 6 | 9 | 11 | 18 | 19 | 20 |
| 35-49 | 11 | 9.5 | 11 | 6 | 6 | 9 | 11 | 15 | 16 | 16 |
| 50-64 | 10 | 11.9 | 41 | 3 | 3 | 12 | 16 | 21 | 22 | 22 |
| 65+ | 9 | 14.4 | 9 | 9 | 10 | 14 | 17 | 18 | 18 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 160 | 9.0 | 13 | 4 | 6 | 8 | 10 | 13 | 15 | 24 |
| 20-34 | 49 | 11.8 | 90 | 6 | 7 | 10 | 12 | 17 | 19 | 71 |
| 35-49 | 28 | 13.0 | 71 | 5 | 6 | 11 | 13 | 25 | 29 | 40 |
| 50-64 | 49 | 15.3 | 87 | 5 | 6 | 13 | 18 | 29 | 37 | 99+ |
| 65+ | 74 | 17.8 | 86 | 5 | 6 | 15 | 24 | 29 | 38 | 46 |
| **TOT. SINGLE DX** | 182 | 8.5 | 13 | 4 | 5 | 8 | 10 | 14 | 16 | 20 |
| **" MULTIPLE DX** | 360 | 12.3 | 65 | 5 | 6 | 10 | 15 | 23 | 28 | 46 |
| TOTAL 0-19 YRS | 291 | 8.3 | 11 | 4 | 5 | 8 | 9 | 12 | 14 | 23 |
| 20-34 | 70 | 11.3 | 69 | 6 | 7 | 10 | 12 | 17 | 19 | 62 |
| 35-49 | 39 | 12.1 | 56 | 6 | 6 | 10 | [illegible] | 24 | 27 | 40 |
| 50-64 | 59 | 14.7 | 80 | 4 | 5 | 13 | 18 | 27 | 36 | 95 |
| 65+ | 83 | 17.4 | 79 | 6 | 8 | 15 | 22 | 29 | 37 | 45 |
| **GRAND TOTAL** | **542** | **11.1** | **51** | **4** | **6** | **9** | **13** | **19** | **25** | **40** |

## 635: Miscellaneous operations on ureter (56.0-56.1; 56.32-56.33; 56.4; 56.8-56.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 53 | 4.0 | 8 | <1 | <1 | 4 | 6 | 8 | 11 | 13 |
| 20-34 | 238 | 3.9 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 11 |
| 35-49 | 281 | 3.7 | 5 | 1 | 1 | 3 | 5 | 7 | 8 | 13 |
| 50-64 | 214 | 4.1 | 7 | 1 | 1 | 3 | 6 | 8 | 10 | 13 |
| 65+ | 101 | 5.1 | 17 | 1 | 2 | 4 | 7 | 9 | 13 | 22 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 42 | 7.0 | 27 | <1 | 1 | 6 | 8 | 15 | 18 | 23 |
| 20-34 | 136 | 5.9 | 28 | 1 | 2 | 5 | 8 | 10 | 12 | 40 |
| 35-49 | 165 | 5.5 | 20 | 1 | 1 | 4 | 7 | 12 | 14 | 21 |
| 50-64 | 178 | 5.7 | 16 | 2 | 2 | 5 | 8 | 11 | 13 | 22 |
| 65+ | 189 | 9.1 | 65 | 2 | 2 | 7 | 12 | 18 | 26 | 43 |
| **TOT. SINGLE DX** | 887 | 4.0 | 8 | 1 | 1 | 3 | 5 | 7 | 9 | 14 |
| **" MULTIPLE DX** | 710 | 6.7 | 36 | 1 | 2 | 5 | 8 | 13 | 17 | 34 |
| TOTAL 0-19 YRS | 95 | 5.3 | 19 | <1 | <1 | 4 | 7 | 11 | 14 | 21 |
| 20-34 | 374 | 4.6 | 15 | 1 | 1 | 4 | 6 | 8 | 10 | 14 |
| 35-49 | 446 | 4.4 | 12 | 1 | 1 | 3 | 6 | 9 | 11 | 16 |
| 50-64 | 392 | 4.8 | 13 | 1 | 1 | 4 | 6 | 9 | 11 | 18 |
| 65+ | 290 | 7.7 | 52 | 2 | 2 | 5 | 9 | 16 | 22 | 39 |
| **GRAND TOTAL** | **1597** | **5.2** | **22** | **1** | **1** | **4** | **7** | **10** | **13** | **23** |

## 636: Cystoscopy and urethroscopy (55.21-55.22; 56.31; 57.32; 58.22)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | [illegible]th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 323 | 2.0 | 5 | <1 | <1 | 1 | 2 | 4 | 6 | 11 |
| 20-34 | 358 | 3.4 | 11 | <1 | <1 | 2 | 4 | 8 | 10 | 15 |
| 35-49 | 362 | 3.3 | 11 | <1 | <1 | 2 | 5 | 7 | 9 | 16 |
| 50-64 | 514 | 3.2 | 8 | <1 | <1 | 2 | 4 | 7 | 9 | 15 |
| 65+ | 594 | 3.6 | 12 | <1 | 1 | 2 | 4 | 8 | 11 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 359 | 2.1 | 5 | <1 | <1 | 1 | 2 | 5 | 7 | 11 |
| 20-34 | 594 | 5.7 | 58 | <1 | 1 | 4 | 7 | 12 | 17 | 50 |
| 35-49 | 588 | 5.6 | 33 | <1 | 1 | 4 | 7 | 12 | 16 | 29 |
| 50-64 | 1104 | 6.3 | 55 | <1 | 1 | 4 | 8 | 15 | 19 | 37 |
| 65+ | 2318 | 8.9 | 81 | 1 | 2 | 6 | 11 | 20 | 26 | 47 |
| **TOT. SINGLE DX** | 2151 | 3.1 | 10 | <1 | <1 | 2 | 4 | 7 | 9 | 16 |
| **" MULTIPLE DX** | 4963 | 7.0 | 66 | <1 | 1 | 5 | 9 | 16 | 22 | 42 |
| TOTAL 0-19 YRS | 682 | 2.0 | 5 | <1 | <1 | 1 | 2 | 5 | 7 | 11 |
| 20-34 | 952 | 4.8 | 42 | <1 | <1 | 3 | 6 | 10 | 14 | 41 |
| 35-49 | 950 | 4.7 | 26 | <1 | <1 | 3 | 6 | 10 | 14 | 27 |
| 50-64 | 1618 | 5.3 | 43 | <1 | 1 | 3 | 7 | 11 | 17 | 31 |
| 65+ | 2912 | 7.8 | 72 | 1 | 1 | 5 | 10 | 17 | 25 | 43 |
| **GRAND TOTAL** | **7114** | **5.9** | **52** | **<1** | **1** | **3** | **7** | **13** | **19** | **37** |

### 637: Transurethral excision or destruction of bladder lesion (57.33; 57.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 3.8 | 10 | <1 | 1 | 3 | 4 | 9 | 11 | 12 |
| 20-34 | 57 | 2.6 | 3 | <1 | 1 | 2 | 3 | 5 | 6 | 12 |
| 35-49 | 136 | 2.6 | 5 | <1 | 1 | 2 | 3 | 5 | 7 | 12 |
| 50-64 | 510 | 2.9 | 4 | <1 | 1 | 2 | 4 | 5 | 7 | 11 |
| 65+ | 1163 | 3.4 | 6 | 1 | 1 | 3 | 4 | 6 | 8 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 12 | 3.0 | 7 | <1 | <1 | 2 | 4 | 8 | 9 | 9 |
| 20-34 | 76 | 4.8 | 78 | 1 | 1 | 3 | 5 | 7 | 8 | 62 |
| 35-49 | 178 | 4.2 | 19 | <1 | 1 | 3 | 5 | 10 | 12 | 21 |
| 50-64 | 660 | 4.5 | 26 | 1 | 1 | 3 | 5 | 9 | 13 | 32 |
| 65+ | 2011 | 6.4 | 49 | 1 | 2 | 4 | 8 | 14 | 19 | 37 |
| **TOT. SINGLE DX** | 1882 | 3.2 | 6 | <1 | 1 | 3 | 4 | 6 | 7 | 12 |
| **" MULTIPLE DX** | 2937 | 5.8 | 44 | 1 | 1 | 4 | 7 | 12 | 17 | 35 |
| TOTAL 0-19 YRS | 28 | 3.5 | 9 | <1 | <1 | 2 | 4 | 9 | 10 | 12 |
| 20-34 | 133 | 3.9 | 48 | 1 | 1 | 3 | 4 | 5 | 7 | 45 |
| 35-49 | 314 | 3.5 | 14 | <1 | 1 | 2 | 4 | 7 | 11 | 19 |
| 50-64 | 1170 | 3.8 | 18 | 1 | 1 | 3 | 4 | 7 | 10 | 20 |
| 65+ | 3174 | 5.3 | 36 | 1 | 1 | 4 | 6 | 11 | 15 | 31 |
| **GRAND TOTAL** | **4819** | **4.8** | **31** | **1** | **1** | **3** | **5** | **9** | **14** | **29** |

### 638: Miscellaneous operations on bladder (57.0-57.2; 57.34; 57.51-57.93; 57.96-57.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 36 | 4.6 | 16 | <1 | 1 | 3 | 7 | 11 | 12 | 17 |
| 20-34 | 65 | 5.5 | 19 | 1 | 1 | 5 | 7 | 10 | 13 | 25 |
| 35-49 | 97 | 4.7 | 10 | <1 | 1 | 4 | 7 | 10 | 11 | 14 |
| 50-64 | 178 | 5.1 | 17 | 1 | 1 | 4 | 7 | 11 | 14 | 19 |
| 65+ | 255 | 5.5 | 23 | 1 | 1 | 3 | 8 | 13 | 15 | 22 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 109 | 10.3 | 184 | 2 | 2 | 6 | 11 | 21 | 37 | 98 |
| 20-34 | 124 | 8.2 | 91 | 1 | 1 | 6 | 10 | 16 | 26 | 51 |
| 35-49 | 174 | 9.8 | 147 | 1 | 2 | 6 | 12 | 19 | 30 | 78 |
| 50-64 | 351 | 9.1 | 86 | 1 | 2 | 7 | 12 | 18 | 25 | 51 |
| 65+ | 869 | 10.9 | 110 | 2 | 2 | 8 | 15 | 22 | 31 | 65 |
| **TOT. SINGLE DX** | 631 | 5.2 | 19 | 1 | 1 | 4 | 7 | 11 | 14 | 20 |
| **" MULTIPLE DX** | 1627 | 10.2 | 113 | 1 | 2 | 7 | 13 | 21 | 30 | 67 |
| TOTAL 0-19 YRS | 145 | 8.9 | 149 | 1 | 1 | 6 | 10 | 17 | 32 | 96 |
| 20-34 | 189 | 7.3 | 68 | 1 | 1 | 5 | 9 | 14 | 19 | 43 |
| 35-49 | 271 | 8.0 | 104 | 1 | 1 | 5 | 10 | 15 | 23 | 48 |
| 50-64 | 529 | 7.8 | 67 | 1 | 1 | 6 | 10 | 16 | 21 | 41 |
| 65+ | 1124 | 9.7 | 96 | 1 | 2 | 7 | 13 | 20 | 28 | 58 |
| **GRAND TOTAL** | **2258** | **8.8** | **92** | **1** | **2** | **6** | **11** | **18** | **25** | **52** |

### 639: Urethral meatotomy (58.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 88 | 1.1 | | <1 | <1 | <1 | 1 | 1 | 2 | 2 |
| 20-34 | 6 | 1.7 | | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| 35-49 | 6 | 2.8 | 5 | <1 | <1 | 2 | 4 | 7 | 7 | 7 |
| 50-64 | 3 | 1.3 | | <1 | <1 | <1 | 2 | 2 | 2 | 2 |
| 65+ | 5 | 2.6 | 4 | 1 | 1 | 2 | 4 | 6 | 6 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 79 | 1.4 | | <1 | <1 | <1 | 2 | 3 | 4 | 6 |
| 20-34 | 6 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 3 | 3 |
| 35-49 | 7 | 6.1 | 103 | <1 | <1 | 2 | 4 | 24 | 29 | 29 |
| 50-64 | 12 | 2.6 | 4 | <1 | <1 | 2 | 3 | 7 | 7 | 7 |
| 65+ | 17 | 7.8 | 105 | 2 | 2 | 5 | 9 | 13 | 34 | 45 |
| **TOT. SINGLE DX** | 108 | 1.3 | 1 | <1 | <1 | <1 | 1 | 2 | 2 | 6 |
| **" MULTIPLE DX** | 121 | 2.7 | 26 | <1 | <1 | 1 | 2 | 5 | 8 | 34 |
| TOTAL 0-19 YRS | 167 | 1.2 | | <1 | <1 | <1 | 1 | 2 | 2 | 4 |
| 20-34 | 12 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 3 | 3 |
| 35-49 | 13 | 4.6 | 57 | <1 | <1 | 2 | 4 | 11 | 26 | 29 |
| 50-64 | 15 | 2.3 | 4 | <1 | <1 | 2 | 2 | 7 | 7 | 7 |
| 65+ | 22 | 6.6 | 86 | 1 | 1 | 3 | 8 | 12 | 26 | 45 |
| **GRAND TOTAL** | **229** | **2.0** | **14** | **<1** | **<1** | **1** | **2** | **3** | **6** | **16** |

### 640: Excision of urethral lesion, repair of urethra (58.23-58.24; 58.3-58.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 212 | 5.2 | 20 | <1 | 1 | 5 | 7 | 10 | 12 | 17 |
| 20-34 | 111 | 2.8 | 8 | <1 | <1 | 2 | 3 | 7 | 9 | 13 |
| 35-49 | 80 | 4.1 | 11 | <1 | <1 | 3 | 6 | 8 | 9 | 18 |
| 50-64 | 83 | 3.8 | 8 | <1 | 1 | 3 | 5 | 8 | 9 | 15 |
| 65+ | 89 | 4.5 | 20 | 1 | 1 | 3 | 5 | 8 | 13 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 116 | 6.3 | 29 | <1 | 1 | 5 | 9 | 13 | 16 | 27 |
| 20-34 | 86 | 4.7 | 23 | <1 | 1 | 3 | 6 | 10 | 15 | 26 |
| 35-49 | 79 | 4.7 | 16 | <1 | <1 | 4 | 6 | 10 | 13 | 77 |
| 50-64 | 113 | 5.0 | 21 | 1 | 1 | 3 | 6 | 10 | 15 | 25 |
| 65+ | 171 | 8.1 | 93 | 1 | 2 | 5 | 9 | 20 | 26 | 49 |
| **TOT. SINGLE DX** | 575 | 4.3 | 16 | <1 | <1 | 3 | 6 | 8 | 10 | 18 |
| **" MULTIPLE DX** | 565 | 6.1 | 46 | <1 | 1 | 4 | 8 | 13 | 18 | 35 |
| TOTAL 0-19 YRS | 328 | 5.6 | 24 | <1 | 1 | 5 | 8 | 11 | 14 | 21 |
| 20-34 | 197 | 3.6 | 16 | <1 | <1 | 2 | 4 | 9 | 11 | 19 |
| 35-49 | 159 | 4.4 | 14 | <1 | <1 | 3 | 6 | 9 | 11 | 24 |
| 50-64 | 196 | 4.5 | 16 | <1 | 1 | 3 | 6 | 9 | 12 | 22 |
| 65+ | 260 | 6.9 | 71 | 1 | 2 | 4 | 8 | 15 | 22 | 43 |
| **GRAND TOTAL** | **1140** | **5.2** | **32** | **<1** | **1** | **4** | **7** | **10** | **15** | **28** |

## 641: Urethral dilation (58.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 121 | 1.2 | | <1 | <1 | <1 | 1 | 2 | 2 | 6 |
| 20-34 | 67 | 2.3 | 3 | <1 | <1 | 1 | 3 | 5 | 6 | 8 |
| 35-49 | 51 | 1.5 | 1 | <1 | <1 | <1 | 2 | 3 | 4 | 7 |
| 50-64 | 68 | 2.8 | 9 | <1 | <1 | 2 | 3 | 8 | 10 | 14 |
| 65+ | 63 | 2.7 | 5 | <1 | 1 | 2 | 4 | 6 | 7 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 164 | 1.9 | 7 | <1 | <1 | 1 | 1 | 3 | 6 | 14 |
| 20-34 | 130 | 4.7 | 45 | <1 | <1 | 2 | 6 | 12 | 16 | 38 |
| 35-49 | 150 | 5.8 | 49 | 1 | 1 | 4 | 7 | 13 | 18 | 55 |
| 50-64 | 189 | 5.3 | 23 | <1 | 1 | 3 | 7 | 12 | 16 | 22 |
| 65+ | 401 | 8.9 | 79 | 1 | 2 | 7 | 12 | 18 | 27 | 48 |
| **TOT. SINGLE DX** | 370 | 2.0 | 4 | <1 | <1 | 1 | 2 | 4 | 6 | 12 |
| **" MULTIPLE DX** | 1034 | 6.1 | 55 | <1 | <1 | 3 | 8 | 14 | 19 | 39 |
| TOTAL 0-19 YRS | 285 | 1.6 | 5 | <1 | <1 | <1 | 1 | 2 | 5 | 12 |
| 20-34 | 197 | 3.9 | 32 | <1 | <1 | 2 | 5 | 8 | 14 | 34 |
| 35-49 | 201 | 4.7 | 41 | <1 | <1 | 2 | 6 | 11 | 16 | 46 |
| 50-64 | 257 | 4.7 | 21 | <1 | <1 | 3 | 7 | 11 | 15 | 21 |
| 65+ | 464 | 8.0 | 74 | 1 | 1 | 5 | 11 | 16 | 24 | 44 |
| **GRAND TOTAL** | **1404** | **5.1** | **45** | **<1** | **<1** | **2** | **7** | **12** | **17** | **37** |

## 642: Miscellaneous operations on urethra (58.0; 58.21; 58.5; 58.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 84 | 1.8 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| 20-34 | 112 | 2.8 | 6 | <1 | 1 | 2 | 3 | 7 | 9 | 12 |
| 35-49 | 93 | 2.8 | 8 | 1 | 1 | 2 | 3 | 6 | 8 | 17 |
| 50-64 | 86 | 3.2 | 5 | 1 | 1 | 2 | 4 | 7 | 8 | 10 |
| 65+ | 95 | 3.4 | 4 | 1 | 1 | 3 | 4 | 6 | 8 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 123 | 1.9 | 3 | <1 | <1 | 1 | 2 | 3 | 5 | 12 |
| 20-34 | 141 | 3.5 | 25 | 1 | 1 | 2 | 3 | 9 | 10 | 19 |
| 35-49 | 110 | 4.8 | 61 | 1 | 1 | 3 | 6 | 9 | 11 | 41 |
| 50-64 | 158 | 5.4 | 41 | 1 | 1 | 3 | 6 | 11 | 17 | 32 |
| 65+ | 283 | 6.5 | 46 | 1 | 2 | 4 | 8 | 15 | 19 | 31 |
| **TOT. SINGLE DX** | 470 | 2.8 | 6 | <1 | 1 | 2 | 3 | 6 | 8 | 12 |
| **" MULTIPLE DX** | 815 | 4.8 | 40 | 1 | 1 | 3 | 6 | 11 | 15 | 30 |
| TOTAL 0-19 YRS | 207 | 1.9 | 3 | <1 | <1 | 1 | 2 | 3 | 5 | 11 |
| 20-34 | 253 | 3.2 | 17 | <1 | 1 | 2 | 3 | 8 | 9 | 13 |
| 35-49 | 203 | 3.9 | 38 | 1 | 1 | 2 | 4 | 8 | 10 | 17 |
| 50-64 | 244 | 4.6 | 30 | 1 | 1 | 3 | 5 | 9 | 11 | 30 |
| 65+ | 378 | 5.7 | 38 | 1 | 2 | 4 | 7 | 12 | 18 | 28 |
| **GRAND TOTAL** | **1285** | **4.1** | **28** | **<1** | **1** | **2** | **5** | **9** | **12** | **26** |

## 643: Urethral suspension (59.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 33 | 6.0 | 3 | 2 | 4 | 6 | 7 | 8 | 8 | 10 |
| 35-49 | 162 | 6.6 | 4 | 4 | 4 | 7 | 8 | 9 | 9 | 12 |
| 50-64 | 201 | 6.8 | 4 | 4 | 4 | 7 | 8 | 10 | 10 | 13 |
| 65+ | 104 | 7.7 | 5 | 5 | 5 | 7 | 9 | 10 | 12 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 59 | 7.5 | 10 | 4 | 5 | 7 | 8 | 10 | 12 | 24 |
| 35-49 | 293 | 7.6 | 7 | 4 | 5 | 7 | 9 | 11 | 12 | 18 |
| 50-64 | 293 | 7.9 | 10 | 5 | 5 | 7 | 9 | 11 | 13 | 22 |
| 65+ | 204 | 9.3 | 21 | 5 | 5 | 8 | 10 | 14 | 17 | 29 |
| **TOT. SINGLE DX** | 500 | 6.9 | 5 | 4 | 4 | 7 | 8 | 9 | 10 | 13 |
| **" MULTIPLE DX** | 850 | 8.1 | 13 | 5 | 5 | 7 | 9 | 12 | 14 | 23 |
| TOTAL 0-19 YRS | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 92 | 7.0 | 9 | 4 | 5 | 7 | 8 | 9 | 11 | 22 |
| 35-49 | 455 | 7.3 | 7 | 4 | 5 | 7 | 8 | 10 | 11 | 17 |
| 50-64 | 494 | 7.4 | 8 | 4 | 5 | 7 | 8 | 10 | 12 | 21 |
| 65+ | 308 | 8.8 | 17 | 5 | 5 | 8 | 10 | 13 | 16 | 26 |
| **GRAND TOTAL** | **1350** | **7.6** | **10** | **4** | **5** | **7** | **9** | **11** | **13** | **21** |

## 644: Other repair of stress incontinence (59.3-59.4; 59.6-59.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 12 | 5.4 | 8 | <1 | 1 | 5 | 7 | 10 | 10 | 10 |
| 35-49 | 66 | 6.0 | 3 | 4 | 4 | 6 | 7 | 9 | 9 | 10 |
| 50-64 | 90 | 6.4 | 4 | 3 | 4 | 6 | 8 | 9 | 10 | 12 |
| 65+ | 56 | 6.6 | 4 | 4 | 4 | 6 | 8 | 10 | 10 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 8.0 | 8 | 6 | 6 | 8 | 10 | 10 | 10 | 10 |
| 20-34 | 26 | 5.3 | 3 | 2 | 3 | 5 | 6 | 8 | 9 | 10 |
| 35-49 | 106 | 6.9 | 11 | 3 | 4 | 7 | 8 | 9 | 13 | 22 |
| 50-64 | 139 | 6.9 | 20 | 3 | 4 | 6 | 8 | 10 | 12 | 26 |
| 65+ | 119 | 8.6 | 40 | 3 | 4 | 7 | 9 | 13 | 17 | 41 |
| **TOT. SINGLE DX** | 224 | 6.3 | 4 | 3 | 4 | 6 | 8 | 9 | 10 | 11 |
| **" MULTIPLE DX** | 392 | 7.3 | 24 | 3 | 4 | 7 | 8 | 11 | 14 | 27 |
| TOTAL 0-19 YRS | 2 | 8.0 | 8 | 6 | 6 | 8 | 10 | 10 | 10 | 10 |
| 20-34 | 38 | 5.3 | 5 | 2 | 3 | 5 | 6 | 8 | 10 | 10 |
| 35-49 | 172 | 6.6 | 8 | 3 | 4 | 6 | 8 | 9 | 10 | 17 |
| 50-64 | 229 | 6.7 | 14 | 3 | 4 | 6 | 8 | 9 | 11 | 23 |
| 65+ | 175 | 8.0 | 30 | 3 | 4 | 7 | 9 | 12 | 14 | 30 |
| **GRAND TOTAL** | **616** | **7.0** | **17** | **3** | **4** | **6** | **8** | **10** | **12** | **23** |

### 645: Ureteral catheterization (59.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 2.4 | 4 | <1 | <1 | <1 | 4 | 5 | 5 | 5 |
| 20-34 | 9 | 3.9 | 6 | 1 | 1 | 4 | 5 | 8 | 9 | 9 |
| 35-49 | 5 | 7.8 | 28 | 4 | 4 | 5 | 10 | 17 | 17 | 17 |
| 50-64 | 7 | 4.0 | 33 | 1 | 1 | 2 | 3 | 14 | 17 | 17 |
| 65+ | 8 | 2.9 | 3 | 1 | 1 | 2 | 3 | 6 | 7 | 7 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 2.8 | 11 | <1 | <1 | 1 | 3 | 7 | 12 | 13 |
| 20-34 | 20 | 5.9 | 22 | 1 | 1 | 4 | 9 | 13 | 15 | 16 |
| 35-49 | 22 | 9.5 | 133 | <1 | <1 | 7 | 10 | 18 | 35 | 55 |
| 50-64 | 19 | 8.2 | 32 | 1 | 1 | 7 | 13 | 15 | 18 | 20 |
| 65+ | 37 | 10.1 | 127 | 1 | 2 | 8 | 12 | 21 | 30 | 61 |
| **TOT. SINGLE DX** | 34 | 4.0 | 15 | <1 | 1 | 3 | 6 | 8 | 15 | 17 |
| **" MULTIPLE DX** | 111 | 8.0 | 84 | 1 | 1 | 5 | 11 | 16 | 21 | 57 |
| TOTAL 0-19 YRS | 18 | 2.7 | 9 | <1 | <1 | 1 | 4 | 5 | 10 | 13 |
| 20-34 | 29 | 5.2 | 18 | 1 | 1 | 4 | 7 | 12 | 14 | 16 |
| 35-49 | 27 | 9.2 | 113 | <1 | <1 | 7 | 10 | 17 | 27 | 55 |
| 50-64 | 26 | 7.0 | 35 | 1 | 1 | 5 | 12 | 15 | 18 | 20 |
| 65+ | 45 | 8.8 | 112 | 1 | 2 | 5 | 10 | 20 | 28 | 61 |
| **GRAND TOTAL** | **145** | **7.1** | **70** | **<1** | **1** | **4** | **10** | **15** | **20** | **55** |

### 646: Miscellaneous operations on urinary tract (59.0-59.21, 59.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 4.5 | 5 | 3 | 3 | 4 | 6 | 6 | 6 | 6 |
| 20-34 | 7 | 5.4 | 10 | 1 | 1 | 5 | 8 | 10 | 10 | 10 |
| 35-49 | 5 | 10.2 | 43 | 5 | 5 | 9 | 13 | 21 | 21 | 21 |
| 50-64 | 8 | 5.5 | 6 | 2 | 2 | 6 | 7 | 8 | 8 | 8 |
| 65+ | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 15.5 | 665 | 2 | 2 | 5 | 8 | 62 | 68 | 68 |
| 20-34 | 19 | 8.5 | 23 | 3 | 4 | 8 | 9 | 17 | 20 | 22 |
| 35-49 | 5 | 9.0 | 6 | 6 | 6 | 9 | 10 | 13 | 13 | 13 |
| 50-64 | 21 | 12.1 | 37 | 4 | 7 | 9 | 16 | 22 | 25 | 25 |
| 65+ | 29 | 15.0 | 281 | 2 | 2 | 9 | 17 | 34 | 50 | 80 |
| **TOT. SINGLE DX** | 23 | 6.3 | 17 | 1 | 2 | 5 | 8 | 10 | 14 | 21 |
| **" MULTIPLE DX** | 80 | 12.4 | 165 | 2 | 3 | 9 | 14 | 24 | 33 | 76 |
| TOTAL 0-19 YRS | 8 | 12.8 | 502 | 2 | 2 | 5 | 7 | 50 | 68 | 68 |
| 20-34 | 26 | 7.7 | 21 | 2 | 3 | 7 | 8 | 15 | 18 | 22 |
| 35-49 | 10 | 9.6 | 22 | 5 | 5 | 9 | 11 | 17 | 21 | 21 |
| 50-64 | 29 | 10.3 | 37 | 2 | 3 | 9 | 14 | 19 | 25 | 25 |
| 65+ | 30 | 14.6 | 277 | 2 | 2 | 9 | 15 | 33 | 48 | 80 |
| **GRAND TOTAL** | **103** | **11.0** | **138** | **2** | **2** | **8** | **12** | **22** | **28** | **74** |

### 647: Miscellaneous diagnostic and therapeutic procedures on urinary tract (55.29; 56.39; 57.31; 57.39; 57.94-57.95; 58.29; 59.29)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

### 648: Transurethral prostatectomy (60.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 5.5 | 2 | 4 | 4 | 5 | 7 | 7 | 7 | 7 |
| 20-34 | 5 | 5.6 | 22 | 3 | 3 | 4 | 7 | 14 | 14 | 14 |
| 35-49 | 47 | 4.7 | 2 | 3 | 3 | 5 | 5 | 7 | 8 | 8 |
| 50-64 | 1143 | 5.0 | 4 | 3 | 3 | 5 | 6 | 7 | 8 | 11 |
| 65+ | 2628 | 5.4 | 4 | 3 | 4 | 5 | 6 | 8 | 9 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 6.5 | 9 | 4 | 4 | 5 | 9 | 11 | 12 | 12 |
| 20-34 | 15 | 7.4 | 14 | 3 | 4 | 7 | 8 | 14 | 16 | 17 |
| 35-49 | 114 | 6.9 | 21 | 2 | 3 | 5 | 9 | 13 | 17 | 26 |
| 50-64 | 2064 | 6.5 | 21 | 3 | 4 | 5 | 7 | 11 | 14 | 26 |
| 65+ | 7159 | 8.3 | 46 | 3 | 4 | 6 | 9 | 15 | 21 | 38 |
| **TOT. SINGLE DX** | 3831 | 5.3 | 4 | 3 | 3 | 5 | 6 | 7 | 9 | 13 |
| **" MULTIPLE DX** | 9360 | 7.9 | 41 | 3 | 4 | 6 | 9 | 14 | 19 | 36 |
| TOTAL 0-19 YRS | 16 | 6.0 | 6 | 4 | 4 | 5 | 7 | 10 | 11 | 12 |
| 20-34 | 20 | 7.0 | 16 | 3 | 3 | 6 | 8 | 14 | 16 | 17 |
| 35-49 | 161 | 6.3 | 17 | 2 | 3 | 5 | 7 | 11 | 14 | 24 |
| 50-64 | 3207 | 6.0 | 16 | 3 | 3 | 5 | 6 | 9 | 12 | 22 |
| 65+ | 9787 | 7.6 | 37 | 3 | 4 | 6 | 8 | 13 | 18 | 34 |
| **GRAND TOTAL** | **13191** | **7.2** | **32** | **3** | **4** | **6** | **8** | **12** | **17** | **31** |

## 649: Other prostatectomy (60.3-60.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 1.5 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 20-34 | 2 | 6.5 | 1 | 6 | 6 | 6 | 7 | 7 | 7 | 7 |
| 35-49 | 7 | 11.9 | 79 | <1 | 1 | 12 | 14 | 26 | 29 | 29 |
| 50-64 | 136 | 9.5 | 14 | 4 | 6 | 9 | 12 | 15 | 17 | 21 |
| 65+ | 202 | 10.5 | 18 | 5 | 6 | 10 | 12 | 16 | 19 | 26 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 2.6 | 1 | 1 | 1 | 3 | 3 | 4 | 4 | 4 |
| 20-34 | 2 | 11.5 | 221 | <1 | <1 | 11 | 22 | 22 | 22 | 22 |
| 35-49 | 6 | 8.7 | 29 | <1 | <1 | 9 | 12 | 16 | 16 | 16 |
| 50-64 | 175 | 11.3 | 28 | 5 | 6 | 10 | 13 | 19 | 23 | 27 |
| 65+ | 692 | 12.8 | 56 | 6 | 7 | 11 | 15 | 21 | 26 | 41 |
| **TOT. SINGLE DX** | 349 | 10.1 | 19 | 4 | 6 | 9 | 12 | 15 | 18 | 24 |
| **" MULTIPLE DX** | 882 | 12.4 | 52 | 5 | 7 | 11 | 15 | 21 | 25 | 40 |
| TOTAL 0-19 YRS | 9 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 20-34 | 4 | 9.0 | 82 | <1 | <1 | 6 | 14 | 22 | 22 | 22 |
| 35-49 | 13 | 10.4 | 55 | <1 | <1 | 11 | 13 | 19 | 27 | 29 |
| 50-64 | 311 | 10.5 | 23 | 5 | 6 | 9 | 13 | 17 | 20 | 26 |
| 65+ | 894 | 12.3 | 49 | 6 | 7 | 11 | 14 | 20 | 25 | 40 |
| **GRAND TOTAL** | **1231** | **11.8** | **43** | **5** | **6** | **10** | **14** | **19** | **24** | **39** |

## 650: Miscellaneous operations on prostate and seminal vesicles (60.0; 60.11-60.15; 60.7-60.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 1.5 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 20-34 | 4 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 35-49 | 42 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 3 | 5 |
| 50-64 | 306 | 2.0 | 3 | <1 | <1 | 2 | 2 | 3 | 4 | 7 |
| 65+ | 507 | 2.4 | 4 | <1 | 1 | 2 | 2 | 4 | 6 | 14 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 5.5 | 16 | 2 | 2 | 4 | 8 | 11 | 11 | 11 |
| 20-34 | 3 | 4.7 | 30 | <1 | <1 | 2 | 9 | 11 | 11 | 11 |
| 35-49 | 45 | 3.6 | 9 | <1 | 1 | 2 | 4 | 8 | 10 | 15 |
| 50-64 | 291 | 4.0 | 35 | <1 | 1 | 2 | 4 | 9 | 13 | 37 |
| 65+ | 767 | 6.0 | 49 | 1 | 1 | 3 | 7 | 14 | 20 | 38 |
| **TOT. SINGLE DX** | 861 | 2.2 | 4 | <1 | 1 | 2 | 2 | 3 | 5 | 10 |
| **" MULTIPLE DX** | 1110 | 5.4 | 45 | 1 | 1 | 3 | 6 | 13 | 18 | 38 |
| TOTAL 0-19 YRS | 6 | 4.2 | 14 | 1 | 1 | 3 | 6 | 10 | 11 | 11 |
| 20-34 | 7 | 3.3 | 13 | <1 | <1 | 2 | 4 | 10 | 11 | 11 |
| 35-49 | 87 | 2.6 | 6 | <1 | 1 | 2 | 3 | 5 | 8 | 14 |
| 50-64 | 597 | 2.9 | 20 | <1 | <1 | 2 | 3 | 6 | 9 | 23 |
| 65+ | 1274 | 4.6 | 35 | 1 | 1 | 2 | 5 | 11 | 16 | 32 |
| **GRAND TOTAL** | **1971** | **4.0** | **29** | **<1** | **1** | **2** | **4** | **9** | **14** | **30** |

## 651: Operations on scrotum and tunica vaginalis (61.0-61.11; 61.2-61.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 76 | 2.0 | 2 | <1 | <1 | 1 | 2 | 4 | 6 | 8 |
| 20-34 | 64 | 2.7 | 6 | <1 | 1 | 2 | 3 | 5 | 6 | 17 |
| 35-49 | 49 | 3.1 | 37 | <1 | <1 | 2 | 3 | 4 | 6 | 44 |
| 50-64 | 58 | 2.4 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 65+ | 38 | 2.9 | 3 | 1 | 1 | 2 | 4 | 6 | 7 | 8 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 51 | 2.5 | 5 | <1 | 1 | 2 | 3 | 4 | 5 | 15 |
| 20-34 | 50 | 4.6 | 20 | <1 | 1 | 3 | 6 | 12 | 14 | 20 |
| 35-49 | 44 | 4.9 | 31 | 1 | 1 | 3 | 6 | 10 | 19 | 28 |
| 50-64 | 55 | 6.4 | 69 | 1 | 1 | 3 | 8 | 13 | 17 | 53 |
| 65+ | 73 | 7.2 | 43 | 2 | 2 | 5 | 10 | 14 | 18 | 39 |
| **TOT. SINGLE DX** | 285 | 2.6 | 9 | <1 | 1 | 2 | 3 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 273 | 5.3 | 38 | 1 | 1 | 3 | 7 | 12 | 16 | 28 |
| TOTAL 0-19 YRS | 127 | 2.2 | 4 | <1 | <1 | 2 | 3 | 4 | 6 | 10 |
| 20-34 | 114 | 3.6 | 13 | <1 | 1 | 2 | 4 | 7 | 12 | 19 |
| 35-49 | 93 | 4.0 | 35 | <1 | 1 | 2 | 4 | 7 | 12 | 37 |
| 50-64 | 113 | 4.3 | 38 | 1 | 1 | 3 | 4 | 10 | 13 | 38 |
| 65+ | 111 | 5.8 | 34 | 1 | 2 | 4 | 7 | 12 | 16 | 32 |
| **GRAND TOTAL** | **558** | **3.9** | **25** | **<1** | **1** | **2** | **4** | **8** | **12** | **25** |

## 652: Orchiectomy (62.3-62.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 101 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 20-34 | 197 | 2.8 | 5 | 1 | 1 | 2 | 3 | 5 | 7 | 14 |
| 35-49 | 85 | 2.7 | 3 | 1 | 1 | 2 | 3 | 5 | 6 | 11 |
| 50-64 | 66 | 3.5 | 7 | 1 | 1 | 3 | 4 | 6 | 11 | 15 |
| 65+ | 155 | 3.6 | 7 | 1 | 1 | 3 | 4 | 7 | 9 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 88 | 2.2 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 112 | 4.5 | 28 | 1 | 1 | 3 | 5 | 10 | 15 | 32 |
| 35-49 | 65 | 5.9 | 79 | 1 | 1 | 3 | 5 | 11 | 22 | 50 |
| 50-64 | 120 | 5.9 | 40 | 1 | 1 | 3 | 7 | 13 | 19 | 34 |
| 65+ | 446 | 7.7 | 60 | 2 | 2 | 5 | 10 | 17 | 22 | 39 |
| **TOT. SINGLE DX** | 604 | 3.0 | 6 | 1 | 1 | 2 | 3 | 6 | 8 | 13 |
| **" MULTIPLE DX** | 831 | 6.3 | 52 | 1 | 1 | 3 | 8 | 14 | 20 | 37 |
| TOTAL 0-19 YRS | 189 | 2.3 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 20-34 | 309 | 3.5 | 14 | 1 | 1 | 2 | 4 | 7 | 10 | 17 |
| 35-49 | 150 | 4.1 | 39 | 1 | 1 | 3 | 4 | 7 | 11 | 43 |
| 50-64 | 186 | 5.1 | 30 | 1 | 1 | 3 | 5 | 11 | 15 | 31 |
| 65+ | 601 | 6.6 | 50 | 1 | 2 | 4 | 8 | 15 | 20 | 35 |
| **GRAND TOTAL** | **1435** | **4.9** | **35** | **1** | **1** | **3** | **5** | **11** | **15** | **31** |

## 653: Orchiopexy (62.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 636 | 1.8 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 6 |
| 20-34 | 83 | 2.1 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 10 |
| 35-49 | 8 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 354 | 2.2 | 4 | <1 | <1 | 2 | 3 | 4 | 6 | 8 |
| 20-34 | 32 | 3.3 | 8 | 1 | 1 | 3 | 4 | 6 | 8 | 16 |
| 35-49 | 4 | 1.5 | | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 728 | 1.9 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 6 |
| **" MULTIPLE DX** | 390 | 2.3 | 5 | <1 | <1 | 2 | 3 | 4 | 6 | 8 |
| TOTAL 0-19 YRS | 990 | 2.0 | 2 | <1 | <1 | 2 | 2 | 3 | 4 | 7 |
| 20-34 | 115 | 2.4 | 4 | 1 | 1 | 2 | 3 | 4 | 5 | 13 |
| 35-49 | 12 | 1.7 | 1 | <1 | <1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **1118** | **2.0** | **2** | **<1** | **<1** | **2** | **2** | **4** | **4** | **7** |

## 654: Miscellaneous operations on testis (62.0-62.12; 62.2; 62.6-62.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 107 | 1.8 | 2 | 1 | 1 | 1 | 2 | 3 | 5 | 10 |
| 20-34 | 90 | 1.9 | 1 | <1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 35-49 | 25 | 2.2 | 2 | <1 | 1 | 2 | 3 | 5 | 6 | 6 |
| 50-64 | 8 | 2.3 | 2 | <1 | <1 | 2 | 3 | 5 | 5 | 5 |
| 65+ | 1 | 6.0 | | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 39 | 2.0 | 3 | <1 | <1 | 1 | 3 | 4 | 5 | 10 |
| 20-34 | 46 | 3.3 | 14 | <1 | <1 | 2 | 4 | 8 | 10 | 21 |
| 35-49 | 20 | 3.2 | 2 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 50-64 | 10 | 4.5 | 14 | 1 | 1 | 3 | 6 | 10 | 13 | 13 |
| 65+ | 10 | 5.9 | 15 | 1 | 1 | 6 | 8 | 11 | 14 | 14 |
| **TOT. SINGLE DX** | 231 | 1.9 | 2 | <1 | 1 | 1 | 2 | 4 | 5 | 7 |
| **" MULTIPLE DX** | 125 | 3.2 | 10 | <1 | 1 | 2 | 4 | 7 | 9 | 16 |
| TOTAL 0-19 YRS | 146 | 1.9 | 3 | <1 | 1 | 1 | 2 | 3 | 5 | 10 |
| 20-34 | 136 | 2.4 | 6 | <1 | 1 | 2 | 3 | 4 | 6 | 15 |
| 35-49 | 45 | 2.6 | 3 | <1 | 1 | 2 | 3 | 5 | 6 | 7 |
| 50-64 | 18 | 3.5 | 10 | <1 | 1 | 2 | 5 | 7 | 11 | 13 |
| 65+ | 11 | 5.9 | 14 | 1 | 1 | 6 | 8 | 11 | 14 | 14 |
| **GRAND TOTAL** | **356** | **2.4** | **5** | **<1** | **1** | **2** | **3** | **5** | **6** | **14** |

## 655: Interruption of vas deferens (63.6-63.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 43 | 1.3 | 1 | <1 | <1 | <1 | 1 | 2 | 3 | 7 |
| 35-49 | 37 | 1.3 | | <1 | <1 | <1 | 1 | 2 | 3 | 4 |
| 50-64 | 5 | 4.0 | 14 | <1 | <1 | 2 | 7 | 9 | 9 | 9 |
| 65+ | 2 | 3.5 | 5 | 2 | 2 | 3 | 5 | 5 | 5 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 20-34 | 21 | 2.9 | 26 | <1 | <1 | 2 | 3 | 3 | 13 | 25 |
| 35-49 | 18 | 4.7 | 46 | <1 | <1 | 2 | 4 | 17 | 23 | 24 |
| 50-64 | 10 | 6.3 | 28 | 1 | 1 | 4 | 12 | 14 | 15 | 15 |
| 65+ | 8 | 12.9 | 213 | 3 | 3 | 8 | 14 | 39 | 47 | 47 |
| **TOT. SINGLE DX** | 87 | 1.5 | 2 | <1 | <1 | <1 | 1 | 2 | 4 | 8 |
| **" MULTIPLE DX** | 59 | 5.3 | 64 | <1 | <1 | 3 | 5 | 15 | 23 | 45 |
| TOTAL 0-19 YRS | 2 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 20-34 | 64 | 1.8 | 10 | <1 | <1 | 1 | 2 | 3 | 3 | 22 |
| 35-49 | 55 | 2.4 | 18 | <1 | <1 | <1 | 2 | 4 | 8 | 24 |
| 50-64 | 15 | 5.5 | 23 | <1 | 1 | 3 | 8 | 14 | 15 | 15 |
| 65+ | 10 | 11.0 | 182 | 2 | 2 | 7 | 8 | 33 | 47 | 47 |
| **GRAND TOTAL** | **146** | **3.1** | **30** | **<1** | **<1** | **1** | **3** | **7** | **12** | **26** |

## 656: Miscellaneous operations on spermatic cord, epididymis, and vas deferens (63.01; 63.1-63.5; 63.8-63.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 307 | 1.7 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 5 |
| 20-34 | 352 | 2.0 | 1 | <1 | 1 | 2 | 3 | 3 | 4 | 7 |
| 35-49 | 297 | 1.9 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 50-64 | 251 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 65+ | 133 | 2.6 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 7 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 227 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 20-34 | 207 | 2.3 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 10 |
| 35-49 | 190 | 2.7 | 5 | 1 | 1 | 2 | 3 | 5 | 6 | 13 |
| 50-64 | 234 | 3.1 | 3 | 1 | 1 | 3 | 4 | 5 | 7 | 12 |
| 65+ | 177 | 4.5 | 17 | 1 | 2 | 3 | 5 | 8 | 13 | 36 |
| **TOT. SINGLE DX** | 1340 | 2.0 | 1 | <1 | <1 | 2 | 3 | 3 | 4 | 6 |
| **" MULTIPLE DX** | 1035 | 2.8 | 7 | <1 | 1 | 2 | 3 | 5 | 7 | 13 |
| TOTAL 0-19 YRS | 534 | 1.7 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 20-34 | 559 | 2.1 | 2 | <1 | 1 | 2 | 3 | 3 | 4 | 7 |
| 35-49 | 487 | 2.2 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 50-64 | 485 | 2.7 | 3 | 1 | 1 | 2 | 3 | 4 | 5 | 10 |
| 65+ | 310 | 3.7 | 11 | 1 | 2 | 3 | 4 | 6 | 9 | 17 |
| **GRAND TOTAL** | **2375** | **2.4** | **4** | **<1** | **1** | **2** | **3** | **4** | **5** | **11** |

## 657: Circumcision, other than newborn (64.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 248 | 1.5 | 8 | <1 | <1 | 1 | 1 | 2 | 2 | 15 |
| 20-34 | 84 | 1.6 | | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 35-49 | 59 | 1.5 | | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 50-64 | 54 | 1.7 | 1 | <1 | <1 | 2 | 2 | 2 | 3 | 7 |
| 65+ | 39 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 4 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 104 | 2.4 | 21 | <1 | <1 | 1 | 2 | 3 | 7 | 32 |
| 20-34 | 47 | 1.9 | 1 | <1 | 1 | 2 | 2 | 3 | 3 | 7 |
| 35-49 | 64 | 2.9 | 8 | <1 | <1 | 2 | 3 | 6 | 8 | 16 |
| 50-64 | 75 | 3.4 | 30 | <1 | <1 | 2 | 3 | 5 | 8 | 37 |
| 65+ | 95 | 5.5 | 39 | <1 | 1 | 3 | 7 | 13 | 18 | 34 |
| **TOT. SINGLE DX** | 658 | 1.5 | 4 | <1 | <1 | 1 | 2 | 2 | 3 | 6 |
| **" MULTIPLE DX** | 440 | 3.8 | 35 | <1 | <1 | 2 | 3 | 8 | 15 | 38 |
| TOTAL 0-19 YRS | 484 | 1.7 | 10 | <1 | <1 | 1 | 1 | 2 | 2 | 18 |
| 20-34 | 152 | 1.7 | 2 | <1 | <1 | 1 | 2 | 3 | 3 | 7 |
| 35-49 | 143 | 2.1 | 5 | <1 | <1 | 1 | 2 | 4 | 6 | 11 |
| 50-64 | 150 | 2.6 | 16 | <1 | <1 | 2 | 2 | 4 | 6 | 24 |
| 65+ | 169 | 5.2 | 54 | <1 | <1 | 3 | 5 | 13 | 20 | 52 |
| **GRAND TOTAL** | **1098** | **2.4** | **17** | **<1** | **<1** | **1** | **2** | **4** | **7** | **24** |

## 658: Miscellaneous operations on penis (64.11; 64.2-64.5; 64.92-64.93; 64.95-64.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 60 | 3.2 | 9 | <1 | <1 | 2 | 5 | 7 | 9 | 15 |
| 20-34 | 34 | 2.9 | 6 | <1 | <1 | 2 | 4 | 7 | 9 | 11 |
| 35-49 | 46 | 2.5 | 3 | <1 | <1 | 2 | 3 | 5 | 7 | 9 |
| 50-64 | 118 | 4.1 | 8 | 1 | 1 | 3 | 5 | 7 | 8 | 17 |
| 65+ | 84 | 3.8 | 7 | 1 | 1 | 3 | 5 | 6 | 10 | 16 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 39 | 2.7 | 6 | <1 | <1 | 2 | 3 | 5 | 10 | 12 |
| 20-34 | 54 | 6.1 | 28 | <1 | 2 | 4 | 8 | 14 | 17 | 26 |
| 35-49 | 86 | 6.6 | 83 | 1 | 2 | 4 | 7 | 14 | 17 | 61 |
| 50-64 | 207 | 5.0 | 20 | 1 | 2 | 4 | 6 | 9 | 12 | 28 |
| 65+ | 164 | 5.3 | 33 | 2 | 2 | 4 | 6 | 10 | 16 | 51 |
| **TOT. SINGLE DX** | 342 | 3.5 | 8 | <1 | 1 | 3 | 5 | 7 | 9 | 15 |
| **" MULTIPLE DX** | 550 | 5.3 | 35 | 1 | 2 | 4 | 6 | 10 | 14 | 35 |
| TOTAL 0-19 YRS | 99 | 3.0 | 9 | <1 | <1 | 2 | 4 | 7 | 9 | 15 |
| 20-34 | 88 | 4.9 | 22 | <1 | <1 | 3 | 6 | 11 | 15 | 23 |
| 35-49 | 132 | 5.2 | 60 | <1 | 1 | 3 | 6 | 10 | 15 | 44 |
| 50-64 | 325 | 4.6 | 16 | 1 | 2 | 4 | 5 | 8 | 12 | 22 |
| 65+ | 248 | 4.8 | 25 | 1 | 2 | 3 | 5 | 9 | 14 | 26 |
| **GRAND TOTAL** | **892** | **4.6** | **25** | **<1** | **1** | **3** | **5** | **9** | **13** | **24** |

## 659: Miscellaneous diagnostic and therapeutic procedures on male genital tract (60.18-60.19; 61.19; 62.19; 63.09; 64.0; 64.19; 64.91; 64.94)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 660: Local excision or destruction of ovarian lesion or tissue (65.12; 65.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 192 | 4.7 | 3 | 2 | 3 | 4 | 6 | 7 | 8 | 12 |
| 20-34 | 686 | 4.6 | 2 | 2 | 3 | 5 | 5 | 6 | 7 | 11 |
| 35-49 | 81 | 4.8 | 3 | 2 | 3 | 5 | 6 | 7 | 8 | 11 |
| 50-64 | 12 | 6.0 | 4 | 3 | 3 | 6 | 7 | 9 | 10 | 10 |
| 65+ | 9 | 6.7 | 6 | 3 | 3 | 6 | 9 | 10 | 10 | 10 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 214 | 5.3 | 6 | 2 | 3 | 5 | 6 | 8 | 10 | 14 |
| 20-34 | 1244 | 5.0 | 4 | 2 | 3 | 5 | 6 | 7 | 8 | 14 |
| 35-49 | 172 | 6.5 | 30 | 3 | 4 | 5 | 7 | 9 | 12 | 27 |
| 50-64 | 37 | 10.3 | 41 | 5 | 5 | 9 | 11 | 19 | 24 | 36 |
| 65+ | 42 | 15.9 | 154 | 6 | 7 | 13 | 19 | 24 | 51 | 64 |
| **TOT. SINGLE DX** | 980 | 4.7 | 3 | 2 | 3 | 5 | 5 | 6 | 7 | 11 |
| **" MULTIPLE DX** | 1709 | 5.6 | 15 | 2 | 3 | 5 | 6 | 8 | 11 | 19 |
| TOTAL 0-19 YRS | 406 | 5.0 | 5 | 2 | 3 | 5 | 6 | 7 | 9 | 13 |
| 20-34 | 1930 | 4.9 | 4 | 2 | 3 | 5 | 6 | 7 | 8 | 13 |
| 35-49 | 253 | 6.0 | 22 | 2 | 3 | 5 | 6 | 9 | 12 | 18 |
| 50-64 | 49 | 9.3 | 36 | 4 | 5 | 7 | 10 | 18 | 20 | 36 |
| 65+ | 51 | 14.3 | 140 | 5 | 6 | 11 | 16 | 22 | 46 | 64 |
| **GRAND TOTAL** | **2689** | **5.3** | **11** | **2** | **3** | **5** | **6** | **7** | **9** | **18** |

## 661: Unilateral oophorectomy or salpingo-oophorectomy (65.3-65.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 106 | 5.1 | 3 | 3 | 3 | 5 | 6 | 7 | 8 | 13 |
| 20-34 | 683 | 5.0 | 2 | 3 | 4 | 5 | 6 | 7 | 7 | 10 |
| 35-49 | 222 | 5.6 | 6 | 3 | 4 | 5 | 6 | 7 | 8 | 19 |
| 50-64 | 45 | 6.5 | 4 | 4 | 4 | 6 | 7 | 8 | 11 | 16 |
| 65+ | 21 | 8.0 | 6 | 5 | 5 | 8 | 9 | 11 | 13 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 126 | 5.9 | 7 | 3 | 3 | 5 | 7 | 9 | 13 | 17 |
| 20-34 | 1393 | 5.6 | 5 | 3 | 4 | 5 | 6 | 8 | 9 | 16 |
| 35-49 | 526 | 6.3 | 7 | 4 | 4 | 6 | 7 | 9 | 12 | 17 |
| 50-64 | 103 | 9.4 | 35 | 4 | 5 | 8 | 11 | 15 | 19 | 38 |
| 65+ | 86 | 14.1 | 164 | 6 | 6 | 10 | 17 | 26 | 34 | 84 |
| **TOT. SINGLE DX** | 1077 | 5.2 | 4 | 3 | 4 | 5 | 6 | 7 | 8 | 14 |
| **" MULTIPLE DX** | 2234 | 6.3 | 16 | 3 | 4 | 5 | 7 | 9 | 12 | 22 |
| TOTAL 0-19 YRS | 232 | 5.5 | 6 | 3 | 3 | 5 | 6 | 8 | 9 | 17 |
| 20-34 | 2076 | 5.4 | 4 | 3 | 4 | 5 | 6 | 7 | 9 | 15 |
| 35-49 | 748 | 6.1 | 7 | 4 | 4 | 5 | 7 | 8 | 11 | 17 |
| 50-64 | 148 | 8.5 | 28 | 4 | 5 | 7 | 10 | 14 | 17 | 34 |
| 65+ | 107 | 12.9 | 139 | 5 | 6 | 9 | 14 | 23 | 30 | 79 |
| **GRAND TOTAL** | **3311** | **5.9** | **13** | **3** | **4** | **5** | **6** | **8** | **11** | **19** |

## 662: Bilateral oophorectomy or salpingo-oophorectomy (65.5-65.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 5.5 | 3 | 4 | 4 | 5 | 7 | 8 | 8 | 8 |
| 20-34 | 96 | 5.2 | 3 | 3 | 3 | 5 | 6 | 7 | 8 | 13 |
| 35-49 | 133 | 5.8 | 2 | 4 | 4 | 5 | 6 | 8 | 10 | 13 |
| 50-64 | 86 | 6.4 | 7 | 4 | 4 | 6 | 7 | 9 | 11 | 18 |
| 65+ | 26 | 7.7 | 3 | 5 | 5 | 7 | 10 | 10 | 11 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 6.0 | 10 | 3 | 3 | 5 | 7 | 11 | 12 | 12 |
| 20-34 | 314 | 6.0 | 5 | 4 | 4 | 5 | 7 | 8 | 10 | 18 |
| 35-49 | 588 | 6.6 | 10 | 4 | 4 | 6 | 7 | 9 | 12 | 19 |
| 50-64 | 216 | 8.9 | 29 | 5 | 5 | 7 | 10 | 14 | 20 | 31 |
| 65+ | 143 | 12.2 | 92 | 5 | 6 | 10 | 14 | 21 | 29 | 63 |
| **TOT. SINGLE DX** | 345 | 5.9 | 5 | 3 | 4 | 6 | 7 | 8 | 10 | 15 |
| **" MULTIPLE DX** | 1267 | 7.5 | 25 | 4 | 4 | 6 | 8 | 12 | 15 | 30 |
| TOTAL 0-19 YRS | 10 | 5.8 | 7 | 3 | 3 | 5 | 7 | 10 | 12 | 12 |
| 20-34 | 410 | 5.8 | 5 | 3 | 4 | 5 | 7 | 8 | 9 | 16 |
| 35-49 | 721 | 6.5 | 9 | 4 | 4 | 6 | 7 | 9 | 11 | 19 |
| 50-64 | 302 | 8.2 | 25 | 4 | 5 | 7 | 9 | 13 | 18 | 30 |
| 65+ | 169 | 11.5 | 81 | 5 | 5 | 9 | 13 | 19 | 24 | 60 |
| **GRAND TOTAL** | **1612** | **7.2** | **21** | **4** | **4** | **6** | **8** | **11** | **14** | **28** |

## 663: Miscellaneous operations on ovary (65.0-65.11; 65.7-65.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 45 | 3.2 | 5 | <1 | 1 | 3 | 4 | 6 | 7 | 12 |
| 20-34 | 164 | 3.2 | 4 | <1 | <1 | 3 | 4 | 6 | 7 | 10 |
| 35-49 | 23 | 3.3 | 5 | <1 | <1 | 3 | 5 | 6 | 8 | 11 |
| 50-64 | 2 | 3.5 | 13 | 1 | 1 | 3 | 6 | 6 | 6 | 6 |
| 65+ | 2 | 9.0 | 18 | 6 | 6 | 9 | 12 | 12 | 12 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 48 | 4.4 | 5 | 1 | 2 | 4 | 6 | 7 | 9 | 11 |
| 20-34 | 412 | 4.4 | 8 | <1 | 1 | 4 | 5 | 7 | 8 | 16 |
| 35-49 | 75 | 5.7 | 63 | <1 | 1 | 5 | 6 | 7 | 12 | 56 |
| 50-64 | 4 | 15.8 | 213 | 3 | 3 | 13 | 27 | 34 | 34 | 34 |
| 65+ | 7 | 9.3 | 23 | 5 | 5 | 9 | 11 | 17 | 19 | 19 |
| **TOT. SINGLE DX** | 236 | 3.2 | 5 | <1 | <1 | 3 | 4 | 6 | 7 | 12 |
| **" MULTIPLE DX** | 546 | 4.7 | 18 | <1 | 1 | 4 | 6 | 7 | 9 | 17 |
| TOTAL 0-19 YRS | 93 | 3.8 | 5 | 1 | 1 | 3 | 5 | 6 | 9 | 12 |
| 20-34 | 576 | 4.0 | 8 | <1 | 1 | 4 | 5 | 6 | 8 | 15 |
| 35-49 | 98 | 5.1 | 51 | <1 | 1 | 5 | 6 | 7 | 12 | 44 |
| 50-64 | 6 | 11.7 | 170 | 1 | 1 | 5 | 21 | 33 | 34 | 34 |
| 65+ | 9 | 9.2 | 19 | 5 | 5 | 9 | 11 | 16 | 19 | 19 |
| **GRAND TOTAL** | **782** | **4.3** | **15** | **<1** | **1** | **4** | **5** | **7** | **9** | **16** |

## 664: Bilateral endoscopic destruction or occlusion of fallopian tubes (66.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 1.5 | | <1 | <1 | 1 | 2 | 2 | 3 | 3 |
| 20-34 | 1637 | 1.1 | | <1 | <1 | <1 | 1 | 1 | 2 | 3 |
| 35-49 | 860 | 1.1 | | <1 | <1 | <1 | 1 | 1 | 2 | 2 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 1.8 | 1 | <1 | <1 | 1 | 2 | 4 | 4 | 4 |
| 20-34 | 651 | 1.8 | 5 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| 35-49 | 368 | 1.4 | 1 | <1 | <1 | 1 | 1 | 2 | 3 | 7 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 2510 | 1.1 | | <1 | <1 | <1 | 1 | 1 | 2 | 2 |
| **" MULTIPLE DX** | 1027 | 1.6 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| TOTAL 0-19 YRS | 21 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 4 |
| 20-34 | 2288 | 1.3 | 2 | <1 | <1 | <1 | 1 | 2 | 2 | 6 |
| 35-49 | 1228 | 1.2 | 1 | <1 | <1 | <1 | 1 | 2 | 2 | 4 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **3537** | **1.2** | **2** | **<1** | **<1** | **<1** | **1** | **2** | **2** | **5** |

### 665: Other bilateral tubal interruption or salpingectomy (66.3; 66.5; 66.63)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 26 | 2.1 | 2 | <1 | 1 | 2 | 2 | 5 | 5 | 6 |
| 20-34 | 1486 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| 35-49 | 614 | 1.7 | | <1 | <1 | 1 | 2 | 3 | 3 | 5 |
| 50-64 | 2 | 1.0 | | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 21 | 2.6 | 1 | <1 | 1 | 3 | 3 | 4 | 5 | 5 |
| 20-34 | 2345 | 2.9 | 2 | 1 | 1 | 3 | 4 | 4 | 5 | 8 |
| 35-49 | 712 | 2.9 | 4 | <1 | 1 | 3 | 4 | 5 | 6 | 11 |
| 50-64 | 7 | 11.9 | 56 | 4 | 5 | 8 | 18 | 23 | 24 | 24 |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 2128 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| **" MULTIPLE DX** | 3085 | 2.9 | 3 | 1 | 1 | 3 | 4 | 4 | 5 | 9 |
| TOTAL 0-19 YRS | 47 | 2.3 | 2 | <1 | 1 | 2 | 3 | 5 | 5 | 6 |
| 20-34 | 3831 | 2.5 | 2 | <1 | 1 | 2 | 3 | 4 | 5 | 7 |
| 35-49 | 1326 | 2.4 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 50-64 | 9 | 9.4 | 65 | 1 | 1 | 7 | 15 | 22 | 24 | 24 |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **5213** | **2.5** | **3** | **<1** | **1** | **2** | **3** | **5** | **5** | **7** |

### 666: Miscellaneous operations on fallopian tube (66.0-66.11; 66.4; 66.61-66.62; 66.69-66.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 117 | 4.2 | 1 | 3 | 3 | 4 | 5 | 6 | 6 | 7 |
| 20-34 | 1476 | 4.2 | 2 | 2 | 3 | 4 | 5 | 6 | 6 | 9 |
| 35-49 | 201 | 4.0 | 3 | <1 | 1 | 4 | 5 | 6 | 6 | 9 |
| 50-64 | 2 | 5.5 | 1 | 5 | 5 | 5 | 6 | 6 | 6 | 6 |
| 65+ | 2 | 3.5 | 5 | 2 | 2 | 3 | 5 | 5 | 5 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 111 | 4.9 | 4 | 2 | 3 | 4 | 5 | 7 | 8 | 13 |
| 20-34 | 1742 | 4.5 | 7 | 2 | 3 | 4 | 5 | 6 | 7 | 10 |
| 35-49 | 233 | 4.7 | 8 | 1 | 2 | 5 | 6 | 7 | 8 | 11 |
| 50-64 | 5 | 5.8 | 11 | 2 | 2 | 5 | 8 | 11 | 11 | 11 |
| 65+ | 2 | 8.5 | 5 | 7 | 7 | 8 | 10 | 10 | 10 | 10 |
| **TOT. SINGLE DX** | 1798 | 4.2 | 2 | 1 | 3 | 4 | 5 | 6 | 6 | 9 |
| **" MULTIPLE DX** | 2093 | 4.6 | 8 | 2 | 3 | 4 | 5 | 6 | 7 | 11 |
| TOTAL 0-19 YRS | 228 | 4.5 | 3 | 2 | 3 | 4 | 5 | 6 | 7 | 12 |
| 20-34 | 3218 | 4.4 | 5 | 2 | 3 | 4 | 5 | 6 | 7 | 9 |
| 35-49 | 434 | 4.4 | 6 | 1 | 1 | 4 | 5 | 6 | 7 | 11 |
| 50-64 | 7 | 5.7 | 8 | 2 | 2 | 5 | 7 | 10 | 11 | 11 |
| 65+ | 4 | 6.0 | 11 | 2 | 2 | 6 | 8 | 10 | 10 | 10 |
| **GRAND TOTAL** | **3891** | **4.4** | **5** | **2** | **3** | **4** | **5** | **6** | **7** | **10** |

### 667: Conization of cervix (67.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 21 | 1.3 | | <1 | <1 | 1 | 2 | 2 | 3 | 3 |
| 20-34 | 537 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| 35-49 | 255 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 6 |
| 50-64 | 93 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| 65+ | 39 | 1.8 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 21 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| 20-34 | 329 | 1.8 | 3 | <1 | <1 | 1 | 2 | 4 | 5 | 9 |
| 35-49 | 192 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 4 | 8 |
| 50-64 | 85 | 2.0 | 2 | <1 | <1 | 2 | 2 | 4 | 5 | 9 |
| 65+ | 38 | 3.0 | 7 | 1 | 1 | 2 | 3 | 8 | 10 | 13 |
| **TOT. SINGLE DX** | 945 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| **" MULTIPLE DX** | 665 | 1.9 | 3 | <1 | <1 | 1 | 2 | 3 | 5 | 9 |
| TOTAL 0-19 YRS | 42 | 1.5 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 5 |
| 20-34 | 866 | 1.6 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| 35-49 | 447 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| 50-64 | 178 | 1.9 | 2 | <1 | <1 | 1 | 2 | 3 | 5 | 9 |
| 65+ | 77 | 2.4 | 5 | <1 | <1 | 2 | 2 | 5 | 9 | 12 |
| **GRAND TOTAL** | **1610** | **1.7** | **2** | **<1** | **<1** | **1** | **2** | **3** | **4** | **8** |

### 668: Other excision or destruction of lesion or tissue of cervix (67.11-67.12; 67.3-67.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 1.4 | | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| 20-34 | 106 | 1.4 | | <1 | <1 | 1 | 1 | 2 | 3 | 6 |
| 35-49 | 68 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 50-64 | 70 | 2.3 | 6 | <1 | <1 | 1 | 2 | 5 | 6 | 15 |
| 65+ | 47 | 3.7 | 12 | <1 | <1 | 2 | 6 | 7 | 8 | 20 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 1.8 | 1 | <1 | <1 | 1 | 2 | 4 | 4 | 4 |
| 20-34 | 159 | 2.4 | 9 | <1 | <1 | 1 | 3 | 5 | 7 | 21 |
| 35-49 | 164 | 2.6 | 8 | <1 | <1 | 1 | 3 | 6 | 8 | 19 |
| 50-64 | 133 | 4.2 | 40 | <1 | <1 | 2 | 5 | 10 | 18 | 31 |
| 65+ | 132 | 6.4 | 38 | 1 | 1 | 6 | 8 | 12 | 17 | 31 |
| **TOT. SINGLE DX** | 298 | 2.0 | 5 | <1 | <1 | 1 | 2 | 5 | 7 | 8 |
| **" MULTIPLE DX** | 596 | 3.8 | 25 | <1 | <1 | 2 | 5 | 8 | 13 | 23 |
| TOTAL 0-19 YRS | 15 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 4 |
| 20-34 | 265 | 2.0 | 6 | <1 | <1 | 1 | 2 | 4 | 6 | 13 |
| 35-49 | 232 | 2.3 | 7 | <1 | <1 | 1 | 2 | 5 | 7 | 14 |
| 50-64 | 203 | 3.6 | 30 | <1 | <1 | 2 | 4 | 8 | 14 | 27 |
| 65+ | 179 | 5.7 | 33 | 1 | 1 | 5 | 7 | 11 | 16 | 25 |
| **GRAND TOTAL** | **894** | **3.2** | **19** | **<1** | **<1** | **2** | **4** | **7** | **10** | **21** |

## 669: Repair and plastic operations on cervix (67.5-67.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 27 | 2.3 | 2 | <1 | <1 | 2 | 3 | 4 | 6 | 7 |
| 20-34 | 517 | 2.1 | 11 | <1 | <1 | 2 | 3 | 3 | 4 | 7 |
| 35-49 | 48 | 2.3 | 1 | 1 | 1 | 2 | 3 | 5 | 5 | 6 |
| 50-64 | 1 | 1.0 | | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 2.3 | 2 | 1 | 1 | 2 | 4 | 4 | 4 | 4 |
| 20-34 | 148 | 3.3 | 11 | 1 | 1 | 2 | 4 | 6 | 9 | 23 |
| 35-49 | 24 | 2.5 | 2 | <1 | 1 | 2 | 3 | 6 | 6 | 7 |
| 50-64 | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 65+ | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| **TOT. SINGLE DX** | 593 | 2.1 | 10 | <1 | <1 | 2 | 3 | 4 | 4 | 7 |
| **" MULTIPLE DX** | 180 | 3.1 | 10 | 1 | 1 | 2 | 4 | 6 | 8 | 21 |
| TOTAL 0-19 YRS | 33 | 2.3 | 2 | <1 | <1 | 2 | 3 | 4 | 6 | 7 |
| 20-34 | 665 | 2.4 | 12 | <1 | 1 | 2 | 3 | 4 | 6 | 10 |
| 35-49 | 72 | 2.3 | 2 | 1 | 1 | 2 | 3 | 5 | 6 | 7 |
| 50-64 | 2 | 1.5 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| 65+ | 1 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| **GRAND TOTAL** | **773** | **2.4** | **10** | **<1** | **1** | **2** | **3** | **4** | **6** | **9** |

## 670: Hysterotomy and excision of lesion of uterus (68.0; 68.13-68.14; 68.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 7.8 | 86 | 1 | 1 | 5 | 7 | 23 | 30 | 30 |
| 20-34 | 183 | 4.4 | 4 | <1 | 1 | 5 | 6 | 7 | 8 | 9 |
| 35-49 | 57 | 3.7 | 4 | <1 | 1 | 4 | 6 | 6 | 6 | 7 |
| 50-64 | 14 | 2.3 | 7 | <1 | <1 | 1 | 2 | 8 | 9 | 9 |
| 65+ | 11 | 1.9 | 1 | <1 | <1 | 2 | 2 | 4 | 4 | 4 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 3.5 | 10 | <1 | <1 | 2 | 6 | 8 | 10 | 10 |
| 20-34 | 318 | 5.1 | 7 | 1 | 1 | 5 | 6 | 7 | 9 | 14 |
| 35-49 | 107 | 5.1 | 8 | 1 | 1 | 5 | 6 | 9 | 10 | 14 |
| 50-64 | 31 | 5.8 | 46 | <1 | <1 | 3 | 8 | 16 | 22 | 27 |
| 65+ | 23 | 5.8 | 41 | 1 | 1 | 4 | 8 | 13 | 24 | 25 |
| **TOT. SINGLE DX** | 273 | 4.2 | 7 | <1 | 1 | 4 | 6 | 7 | 7 | 9 |
| **" MULTIPLE DX** | 492 | 5.1 | 12 | <1 | 1 | 5 | 6 | 8 | 10 | 22 |
| TOTAL 0-19 YRS | 21 | 5.1 | 41 | <1 | <1 | 4 | 7 | 9 | 19 | 30 |
| 20-34 | 501 | 4.9 | 6 | <1 | 1 | 5 | 6 | 7 | 8 | 13 |
| 35-49 | 164 | 4.6 | 8 | <1 | 1 | 5 | 6 | 8 | 10 | 12 |
| 50-64 | 45 | 4.7 | 37 | <1 | <1 | 2 | 6 | 12 | 21 | 27 |
| 65+ | 34 | 4.5 | 32 | 1 | 1 | 2 | 4 | 9 | 21 | 25 |
| **GRAND TOTAL** | **765** | **4.8** | **10** | **<1** | **1** | **5** | **6** | **8** | **9** | **20** |

## 671: Abdominal hysterectomy (68.3-68.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 5.2 | 1 | 4 | 4 | 5 | 6 | 7 | 7 | 7 |
| 20-34 | 856 | 5.9 | 2 | 4 | 4 | 6 | 6 | 7 | 8 | 12 |
| 35-49 | 1449 | 6.0 | 3 | 4 | 4 | 6 | 6 | 8 | 8 | 12 |
| 50-64 | 518 | 6.5 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 14 |
| 65+ | 242 | 7.6 | 6 | 5 | 5 | 7 | 8 | 11 | 13 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 53 | 6.8 | 9 | 4 | 4 | 6 | 7 | 11 | 12 | 21 |
| 20-34 | 4760 | 6.4 | 5 | 4 | 5 | 6 | 7 | 8 | 10 | 16 |
| 35-49 | 9732 | 6.5 | 5 | 4 | 5 | 6 | 7 | 8 | 10 | 15 |
| 50-64 | 2884 | 7.2 | 13 | 5 | 5 | 6 | 8 | 9 | 12 | 21 |
| 65+ | 1079 | 9.5 | 41 | 5 | 6 | 8 | 10 | 16 | 20 | 37 |
| **TOT. SINGLE DX** | 3075 | 6.2 | 4 | 4 | 5 | 6 | 7 | 8 | 9 | 14 |
| **" MULTIPLE DX** | 18508 | 6.8 | 9 | 4 | 5 | 6 | 7 | 9 | 11 | 19 |
| TOTAL 0-19 YRS | 63 | 6.5 | 8 | 4 | 4 | 6 | 7 | 11 | 12 | 20 |
| 20-34 | 5616 | 6.3 | 5 | 4 | 5 | 6 | 7 | 8 | 10 | 15 |
| 35-49 | 11181 | 6.4 | 5 | 4 | 5 | 6 | 7 | 8 | 10 | 15 |
| 50-64 | 3402 | 7.1 | 12 | 5 | 5 | 6 | 8 | 9 | 12 | 20 |
| 65+ | 1321 | 9.2 | 35 | 5 | 5 | 8 | 10 | 14 | 19 | 33 |
| **GRAND TOTAL** | **21583** | **6.7** | **9** | **4** | **5** | **6** | **7** | **9** | **11** | **18** |

## 672: Vaginal hysterectomy (68.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 4.5 | 4 | 2 | 2 | 4 | 6 | 7 | 7 | 7 |
| 20-34 | 615 | 5.0 | 1 | 3 | 4 | 5 | 6 | 6 | 7 | 9 |
| 35-49 | 627 | 5.2 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 10 |
| 50-64 | 196 | 6.5 | 2 | 4 | 5 | 6 | 7 | 8 | 9 | 12 |
| 65+ | 156 | 7.1 | 3 | 5 | 5 | 7 | 8 | 9 | 10 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 4.3 | | 4 | 4 | 4 | 5 | 5 | 5 | 5 |
| 20-34 | 2245 | 5.4 | 3 | 3 | 4 | 5 | 6 | 7 | 8 | 12 |
| 35-49 | 3769 | 5.7 | 3 | 3 | 4 | 5 | 6 | 8 | 9 | 12 |
| 50-64 | 1056 | 6.6 | 7 | 4 | 4 | 6 | 8 | 9 | 10 | 14 |
| 65+ | 646 | 7.7 | 10 | 5 | 5 | 7 | 9 | 11 | 13 | 20 |
| **TOT. SINGLE DX** | 1598 | 5.5 | 3 | 3 | 4 | 5 | 6 | 7 | 8 | 10 |
| **" MULTIPLE DX** | 7719 | 5.9 | 5 | 3 | 4 | 6 | 7 | 8 | 9 | 14 |
| TOTAL 0-19 YRS | 7 | 4.4 | 2 | 2 | 2 | 4 | 5 | 7 | 7 | 7 |
| 20-34 | 2860 | 5.3 | 3 | 3 | 4 | 5 | 6 | 7 | 8 | 11 |
| 35-49 | 4396 | 5.6 | 3 | 3 | 4 | 5 | 6 | 8 | 9 | 12 |
| 50-64 | 1252 | 6.6 | 7 | 4 | 4 | 6 | 7 | 9 | 10 | 14 |
| 65+ | 802 | 7.6 | 9 | 5 | 5 | 7 | 8 | 10 | 12 | 19 |
| **GRAND TOTAL** | **9317** | **5.8** | **5** | **3** | **4** | **6** | **7** | **8** | **9** | **13** |

## 673: Radical hysterectomy and pelvic evisceration (68.6-68.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 13 | 9.4 | 11 | 3 | 5 | 10 | 11 | 14 | 15 | 15 |
| 35-49 | 15 | 9.4 | 8 | 4 | 8 | 9 | 10 | 13 | 15 | 16 |
| 50-64 | 6 | 13.0 | 14 | 9 | 9 | 13 | 15 | 19 | 19 | 19 |
| 65+ | 6 | 13.8 | 8 | 10 | 10 | 13 | 16 | 18 | 18 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 30 | 12.1 | 45 | 4 | 5 | 11 | 15 | 18 | 20 | 38 |
| 35-49 | 29 | 14.9 | 96 | 7 | 8 | 12 | 16 | 25 | 32 | 55 |
| 50-64 | 40 | 16.8 | 147 | 6 | 8 | 12 | 17 | 36 | 45 | 60 |
| 65+ | 21 | 17.7 | 140 | 7 | 7 | 16 | 21 | 33 | 49 | 52 |
| **TOT. SINGLE DX** | 40 | 10.6 | 13 | 4 | 7 | 10 | 13 | 15 | 17 | 19 |
| **" MULTIPLE DX** | 120 | 15.3 | 110 | 6 | 7 | 12 | 17 | 25 | 43 | 56 |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | 43 | 11.3 | 36 | 4 | 5 | 11 | 14 | 18 | 19 | 38 |
| 35-49 | 44 | 13.0 | 73 | 6 | 8 | 10 | 14 | 22 | 27 | 55 |
| 50-64 | 46 | 16.3 | 131 | 7 | 8 | 12 | 16 | 30 | 45 | 60 |
| 65+ | 27 | 16.9 | 112 | 7 | 7 | 14 | 20 | 24 | 47 | 52 |
| **GRAND TOTAL** | **160** | **14.1** | **90** | **6** | **7** | **11** | **16** | **23** | **35** | **55** |

## 674: Dilation and curettage of uterus for termination of pregnancy (69.01)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 78 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 3 | 5 |
| 20-34 | 357 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 3 | 4 |
| 35-49 | 49 | 1.3 | | <1 | <1 | 1 | 1 | 3 | 3 | 4 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 48 | 1.9 | 3 | <1 | <1 | 1 | 2 | 4 | 6 | 10 |
| 20-34 | 269 | 2.4 | 13 | <1 | <1 | 1 | 2 | 5 | 8 | 16 |
| 35-49 | 60 | 2.1 | 3 | <1 | <1 | 1 | 2 | 4 | 6 | 9 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 484 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 3 | 4 |
| **" MULTIPLE DX** | 377 | 2.3 | 11 | <1 | <1 | 1 | 2 | 5 | 7 | 15 |
| TOTAL 0-19 YRS | 126 | 1.5 | 2 | <1 | <1 | 1 | 1 | 3 | 4 | 8 |
| 20-34 | 626 | 1.8 | 7 | <1 | <1 | 1 | 1 | 3 | 5 | 13 |
| 35-49 | 109 | 1.7 | 2 | <1 | <1 | 1 | 2 | 3 | 5 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **861** | **1.7** | **5** | **<1** | **<1** | **1** | **1** | **3** | **5** | **12** |

## 675: Other dilation and curettage (69.02; 69.09)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1360 | 1.5 | 1 | <1 | <1 | 1 | 2 | 2 | 3 | 6 |
| 20-34 | 7189 | 1.3 | 2 | <1 | <1 | 1 | 1 | 2 | 3 | 5 |
| 35-49 | 2592 | 1.3 | | <1 | <1 | 1 | 1 | 2 | 2 | 5 |
| 50-64 | 1076 | 1.3 | | <1 | <1 | 1 | 1 | 2 | 2 | 5 |
| 65+ | 416 | 1.9 | 5 | <1 | <1 | 1 | 2 | 3 | 5 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 401 | 2.3 | 4 | <1 | <1 | 2 | 3 | 5 | 7 | 11 |
| 20-34 | 3471 | 2.1 | 7 | <1 | <1 | 1 | 2 | 4 | 6 | 11 |
| 35-49 | 2749 | 2.0 | 7 | <1 | <1 | 1 | 2 | 4 | 6 | 12 |
| 50-64 | 1359 | 2.0 | 6 | <1 | <1 | 1 | 2 | 3 | 6 | 15 |
| 65+ | 662 | 3.5 | 26 | <1 | <1 | 2 | 3 | 8 | 12 | 27 |
| **TOT. SINGLE DX** | 12633 | 1.4 | 2 | <1 | <1 | 1 | 1 | 2 | 3 | 5 |
| **" MULTIPLE DX** | 8642 | 2.2 | 9 | <1 | <1 | 1 | 2 | 4 | 6 | 13 |
| TOTAL 0-19 YRS | 1761 | 1.6 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 7 |
| 20-34 | 10660 | 1.6 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 7 |
| 35-49 | 5341 | 1.6 | 4 | <1 | <1 | 1 | 2 | 2 | 5 | 9 |
| 50-64 | 2435 | 1.7 | 4 | <1 | <1 | 1 | 2 | 2 | 4 | 10 |
| 65+ | 1078 | 2.9 | 19 | <1 | <1 | 2 | 2 | 6 | 10 | 24 |
| **GRAND TOTAL** | **21275** | **1.7** | **5** | **<1** | **<1** | **1** | **2** | **3** | **4** | **9** |

## 676: Aspiration curettage for termination of pregnancy (69.51)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 363 | 1.1 | | <1 | <1 | 1 | 1 | 1 | 1 | 3 |
| 20-34 | 862 | 1.1 | | <1 | <1 | 1 | 1 | 1 | 2 | 4 |
| 35-49 | 98 | 1.2 | 1 | <1 | <1 | 1 | 1 | 2 | 2 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 56 | 2.1 | 16 | <1 | <1 | <1 | 2 | 4 | 5 | 29 |
| 20-34 | 465 | 2.1 | 7 | <1 | <1 | 1 | 2 | 5 | 7 | 15 |
| 35-49 | 137 | 1.6 | 4 | <1 | <1 | 1 | 1 | 2 | 4 | 9 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 1323 | 1.1 | | <1 | <1 | 1 | 1 | 1 | 2 | 4 |
| **" MULTIPLE DX** | 658 | 2.0 | 7 | <1 | <1 | 1 | 2 | 4 | 7 | 15 |
| TOTAL 0-19 YRS | 419 | 1.2 | 2 | <1 | <1 | 1 | 1 | 1 | 2 | 4 |
| 20-34 | 1327 | 1.5 | 3 | <1 | <1 | 1 | 1 | 2 | 4 | 10 |
| 35-49 | 235 | 1.5 | 3 | <1 | <1 | 1 | 1 | 2 | 3 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **1981** | **1.4** | **3** | **<1** | **<1** | **1** | **1** | **2** | **3** | **9** |

## 677: Other aspiration curettage (69.52; 69.59)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 200 | 1.3 | | <1 | <1 | 1 | 1 | 2 | 2 | 4 |
| 20-34 | 1019 | 1.3 | 1 | <1 | <1 | 1 | 1 | 2 | 2 | 5 |
| 35-49 | 124 | 1.1 | | <1 | <1 | 1 | 1 | 1 | 2 | 4 |
| 50-64 | 3 | 3.3 | 16 | <1 | <1 | 1 | 6 | 8 | 8 | 8 |
| 65+ | 3 | 1.3 | | 1 | 1 | 1 | 2 | 2 | 2 | 2 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 37 | 2.8 | 32 | <1 | <1 | 1 | 2 | 4 | 10 | 34 |
| 20-34 | 325 | 1.9 | 2 | <1 | <1 | 1 | 2 | 4 | 5 | 8 |
| 35-49 | 89 | 1.7 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| 50-64 | 4 | 8.5 | 48 | 2 | 2 | 7 | 13 | 18 | 18 | 18 |
| 65+ | 4 | 1.5 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| **TOT. SINGLE DX** | 1349 | 1.3 | 1 | <1 | <1 | 1 | 1 | 2 | 2 | 4 |
| **" MULTIPLE DX** | 459 | 2.0 | 5 | <1 | <1 | 1 | 2 | 4 | 5 | 9 |
| TOTAL 0-19 YRS | 237 | 1.5 | 6 | <1 | <1 | 1 | 1 | 2 | 3 | 9 |
| 20-34 | 1344 | 1.4 | 2 | <1 | <1 | 1 | 1 | 2 | 3 | 7 |
| 35-49 | 213 | 1.4 | 1 | <1 | <1 | 1 | 1 | 2 | 3 | 5 |
| 50-64 | 7 | 6.3 | 37 | <1 | <1 | 5 | 9 | 16 | 18 | 18 |
| 65+ | 7 | 1.4 | 1 | <1 | <1 | 1 | 2 | 3 | 3 | 3 |
| **GRAND TOTAL** | **1808** | **1.4** | **2** | **<1** | **<1** | **1** | **1** | **2** | **3** | **7** |

## 678: Repair and plastic operations on supporting structures of uterus (69.2-69.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 28 | 4.1 | 2 | <1 | 1 | 4 | 5 | 6 | 6 | 7 |
| 35-49 | 3 | 4.3 | 9 | <1 | <1 | 5 | 6 | 7 | 7 | 7 |
| 50-64 | 3 | 6.0 | 3 | 4 | 4 | 7 | 7 | 7 | 7 | 7 |
| 65+ | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 4.3 | 6 | 2 | 2 | 4 | 5 | 9 | 9 | 9 |
| 20-34 | 82 | 4.4 | 2 | 2 | 3 | 4 | 5 | 6 | 7 | 9 |
| 35-49 | 11 | 5.2 | 6 | <1 | 1 | 5 | 7 | 8 | 8 | 8 |
| 50-64 | 6 | 11.7 | 51 | 6 | 6 | 10 | 13 | 24 | 25 | 25 |
| 65+ | 5 | 9.2 | 43 | <1 | <1 | 8 | 13 | 19 | 19 | 19 |
| **TOT. SINGLE DX** | 35 | 4.3 | 3 | <1 | 1 | 4 | 5 | 7 | 7 | 7 |
| **" MULTIPLE DX** | 110 | 5.1 | 10 | 1 | 2 | 5 | 6 | 8 | 9 | 21 |
| TOTAL 0-19 YRS | 6 | 4.3 | 6 | 2 | 2 | 4 | 5 | 9 | 9 | 9 |
| 20-34 | 110 | 4.3 | 2 | 1 | 2 | 4 | 5 | 6 | 7 | 9 |
| 35-49 | 14 | 5.0 | 7 | <1 | <1 | 5 | 7 | 8 | 8 | 8 |
| 50-64 | 9 | 9.8 | 41 | 4 | 5 | 7 | 12 | 20 | 25 | 25 |
| 65+ | 6 | 8.5 | 38 | <1 | <1 | 7 | 11 | 18 | 19 | 19 |
| **GRAND TOTAL** | **145** | **4.9** | **8** | **1** | **2** | **5** | **6** | **7** | **9** | **19** |

## 679: Miscellaneous operations on uterus, cervix, and supporting structures (67.0; 68.12; 69.1; 69.4; 69.91; 69.93-69.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 29 | 2.2 | 3 | <1 | <1 | 1 | 3 | 5 | 6 | 7 |
| 20-34 | 104 | 2.6 | 3 | <1 | <1 | 2 | 4 | 6 | 6 | 7 |
| 35-49 | 26 | 3.1 | 2 | <1 | <1 | 3 | 4 | 5 | 5 | 6 |
| 50-64 | 11 | 4.2 | 2 | 3 | 3 | 3 | 6 | 7 | 7 | 7 |
| 65+ | 11 | 3.1 | | 2 | 2 | 3 | 4 | 4 | 4 | 4 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 20 | 4.7 | 32 | 1 | 1 | 3 | 5 | 13 | 20 | 21 |
| 20-34 | 180 | 3.4 | 9 | 1 | 1 | 3 | 5 | 6 | 8 | 11 |
| 35-49 | 45 | 3.9 | 13 | <1 | <1 | 3 | 6 | 7 | 14 | 15 |
| 50-64 | 10 | 3.4 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 6 |
| 65+ | 29 | 4.2 | 6 | 2 | 2 | 3 | 5 | 8 | 9 | 14 |
| **TOT. SINGLE DX** | 181 | 2.8 | 3 | <1 | 1 | 2 | 4 | 5 | 6 | 7 |
| **" MULTIPLE DX** | 284 | 3.6 | 11 | <1 | 1 | 3 | 5 | 7 | 8 | 18 |
| TOTAL 0-19 YRS | 49 | 3.2 | 16 | <1 | 1 | 2 | 4 | 6 | 8 | 21 |
| 20-34 | 284 | 3.1 | 8 | <1 | 1 | 2 | 5 | 6 | 7 | 10 |
| 35-49 | 71 | 3.6 | 9 | <1 | <1 | 3 | 5 | 7 | 8 | 15 |
| 50-64 | 21 | 3.8 | 2 | 1 | 2 | 3 | 5 | 6 | 7 | 7 |
| 65+ | 40 | 3.9 | 5 | 2 | 2 | 3 | 4 | 6 | 8 | 14 |
| **GRAND TOTAL** | **465** | **3.3** | **8** | **<1** | **1** | **3** | **5** | **6** | **7** | **15** |

## 680: Plastic repair of cystocele and rectocele (70.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 65 | 4.2 | 6 | 1 | 1 | 4 | 6 | 8 | 9 | 11 |
| 35-49 | 173 | 5.4 | 3 | 2 | 3 | 5 | 7 | 8 | 8 | 10 |
| 50-64 | 325 | 5.5 | 3 | 2 | 3 | 5 | 7 | 8 | 8 | 11 |
| 65+ | 240 | 6.0 | 4 | 3 | 4 | 6 | 7 | 8 | 9 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 96 | 5.4 | 4 | 2 | 3 | 5 | 7 | 8 | 10 | 12 |
| 35-49 | 380 | 6.5 | 11 | 3 | 4 | 6 | 7 | 9 | 11 | 20 |
| 50-64 | 533 | 6.7 | 6 | 3 | 4 | 6 | 8 | 10 | 12 | 16 |
| 65+ | 428 | 6.9 | 9 | 4 | 4 | 6 | 8 | 10 | 12 | 17 |
| **TOT. SINGLE DX** | 804 | 5.5 | 4 | 2 | 3 | 5 | 7 | 8 | 9 | 11 |
| **" MULTIPLE DX** | 1438 | 6.6 | 9 | 3 | 4 | 6 | 8 | 10 | 11 | 17 |
| TOTAL 0-19 YRS | 2 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| 20-34 | 161 | 5.0 | 6 | 1 | 2 | 5 | 6 | 8 | 9 | 11 |
| 35-49 | 553 | 6.1 | 9 | 3 | 3 | 6 | 7 | 9 | 10 | 17 |
| 50-64 | 858 | 6.2 | 6 | 3 | 4 | 6 | 7 | 9 | 11 | 14 |
| 65+ | 668 | 6.6 | 8 | 3 | 4 | 6 | 8 | 10 | 11 | 17 |
| **GRAND TOTAL** | **2242** | **6.2** | **7** | **3** | **4** | **6** | **7** | **9** | **10** | **16** |

## 681: Other colporrhaphy (70.6-70.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 4.1 | 12 | <1 | <1 | 3 | 7 | 9 | 11 | 11 |
| 20-34 | 128 | 3.0 | 5 | <1 | <1 | 3 | 4 | 5 | 8 | 12 |
| 35-49 | 57 | 3.2 | 4 | <1 | 1 | 3 | 4 | 6 | 7 | 10 |
| 50-64 | 33 | 5.1 | 7 | <1 | 1 | 6 | 7 | 8 | 8 | 14 |
| 65+ | 19 | 7.5 | 10 | 1 | 2 | 7 | 9 | 12 | 17 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 30 | 4.4 | 96 | <1 | <1 | 1 | 4 | 6 | 15 | 54 |
| 20-34 | 78 | 3.8 | 16 | <1 | 1 | 3 | 4 | 9 | 10 | 25 |
| 35-49 | 53 | 4.7 | 12 | <1 | 1 | 4 | 7 | 9 | 12 | 16 |
| 50-64 | 50 | 5.7 | 9 | 1 | 1 | 6 | 7 | 9 | 10 | 15 |
| 65+ | 48 | 11.6 | 193 | 2 | 2 | 7 | 12 | 27 | 32 | 85 |
| **TOT. SINGLE DX** | 250 | 3.7 | 9 | <1 | 1 | 3 | 5 | 7 | 9 | 14 |
| **" MULTIPLE DX** | 259 | 5.8 | 63 | <1 | 1 | 4 | 7 | 10 | 16 | 40 |
| TOTAL 0-19 YRS | 43 | 4.3 | 70 | <1 | <1 | 2 | 5 | 8 | 12 | 54 |
| 20-34 | 206 | 3.3 | 10 | <1 | 1 | 3 | 4 | 6 | 9 | 13 |
| 35-49 | 110 | 3.9 | 9 | <1 | 1 | 3 | 6 | 8 | 10 | 14 |
| 50-64 | 83 | 5.5 | 9 | 1 | 1 | 6 | 7 | 9 | 10 | 15 |
| 65+ | 67 | 10.4 | 147 | 2 | 2 | 7 | 11 | 22 | 30 | 77 |
| **GRAND TOTAL** | **509** | **4.8** | **37** | **<1** | **1** | **4** | **6** | **9** | **12** | **30** |

## 682: Miscellaneous operations on vagina (70.0-70.1; 70.22-70.24; 70.3-70.4; 70.8-70.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 82 | 3.5 | 12 | <1 | <1 | 2 | 5 | 10 | 12 | 14 |
| 20-34 | 260 | 2.9 | 8 | <1 | <1 | 2 | 4 | 6 | 9 | 15 |
| 35-49 | 63 | 3.3 | 8 | <1 | <1 | 2 | 5 | 7 | 8 | 15 |
| 50-64 | 60 | 3.1 | 7 | <1 | <1 | 2 | 5 | 6 | 7 | 14 |
| 65+ | 86 | 4.9 | 42 | 1 | 1 | 4 | 6 | 8 | 10 | 45 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 63 | 4.1 | 76 | <1 | 1 | 2 | 5 | 7 | 8 | 62 |
| 20-34 | 245 | 4.2 | 12 | <1 | 1 | 3 | 6 | 9 | 11 | 15 |
| 35-49 | 134 | 6.3 | 48 | <1 | 1 | 5 | 7 | 11 | 22 | 40 |
| 50-64 | 144 | 6.5 | 32 | 1 | 2 | 5 | 7 | 12 | 18 | 34 |
| 65+ | 174 | 8.2 | 74 | 2 | 2 | 6 | 9 | 14 | 21 | 51 |
| **TOT. SINGLE DX** | 551 | 3.4 | 15 | <1 | <1 | 2 | 5 | 7 | 9 | 15 |
| **" MULTIPLE DX** | 760 | 5.9 | 44 | <1 | 1 | 5 | 7 | 11 | 15 | 37 |
| TOTAL 0-19 YRS | 145 | 3.7 | 40 | <1 | <1 | 2 | 5 | 8 | 11 | 17 |
| 20-34 | 505 | 3.6 | 11 | <1 | <1 | 2 | 5 | 8 | 10 | 15 |
| 35-49 | 197 | 5.3 | 38 | <1 | <1 | 4 | 7 | 9 | 13 | 38 |
| 50-64 | 204 | 5.5 | 27 | <1 | 1 | 4 | 7 | 11 | 16 | 31 |
| 65+ | 260 | 7.1 | 66 | 1 | 2 | 5 | 8 | 13 | 19 | 51 |
| **GRAND TOTAL** | **1311** | **4.9** | **33** | **<1** | **<1** | **4** | **6** | **9** | **13** | **27** |

## 683: Incision and excision of lesion of vulva and perineum (71.0-71.11; 71.2-71.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 73 | 1.4 | | <1 | <1 | 1 | 1 | 3 | 4 | 5 |
| 20-34 | 310 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 6 |
| 35-49 | 196 | 1.8 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| 50-64 | 60 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 5 | 8 |
| 65+ | 48 | 2.6 | 3 | <1 | <1 | 2 | 3 | 5 | 6 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 32 | 2.1 | 1 | <1 | <1 | 2 | 3 | 4 | 4 | 5 |
| 20-34 | 150 | 2.8 | 7 | <1 | <1 | 2 | 3 | 5 | 9 | 16 |
| 35-49 | 68 | 3.3 | 16 | <1 | <1 | 2 | 3 | 7 | 9 | 27 |
| 50-64 | 86 | 4.7 | 69 | <1 | 1 | 2 | 4 | 7 | 18 | 53 |
| 65+ | 61 | 7.9 | 68 | 1 | 1 | 5 | 11 | 17 | 24 | 44 |
| **TOT. SINGLE DX** | 687 | 1.7 | 2 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| **" MULTIPLE DX** | 397 | 4.0 | 34 | <1 | <1 | 2 | 4 | 9 | 14 | 30 |
| TOTAL 0-19 YRS | 105 | 1.6 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 5 |
| 20-34 | 460 | 2.0 | 4 | <1 | <1 | 1 | 2 | 4 | 5 | 12 |
| 35-49 | 264 | 2.2 | 7 | <1 | <1 | 1 | 2 | 4 | 6 | 13 |
| 50-64 | 146 | 3.5 | 44 | <1 | <1 | 2 | 3 | 6 | 8 | 48 |
| 65+ | 109 | 5.6 | 46 | <1 | 1 | 3 | 6 | 14 | 18 | 37 |
| **GRAND TOTAL** | **1084** | **2.6** | **15** | **<1** | **<1** | **1** | **3** | **5** | **8** | **18** |

## 684: Vulvectomy (71.5-71.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 1.0 | | <1 | <1 | <1 | <1 | <1 | <1 | <1 |
| 20-34 | 11 | 3.0 | 7 | <1 | <1 | 2 | 4 | 8 | 8 | 8 |
| 35-49 | 10 | 4.8 | 24 | <1 | <1 | 3 | 6 | 12 | 18 | 18 |
| 50-64 | 19 | 5.1 | 23 | <1 | 1 | 3 | 6 | 11 | 16 | 21 |
| 65+ | 14 | 7.6 | 21 | 2 | 3 | 6 | 11 | 14 | 16 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 5 | 5.6 | 27 | 1 | 1 | 4 | 9 | 14 | 14 | 14 |
| 35-49 | 8 | 10.6 | 52 | 2 | 2 | 10 | 17 | 20 | 21 | 21 |
| 50-64 | 20 | 9.9 | 68 | 1 | 2 | 9 | 12 | 17 | 28 | 38 |
| 65+ | 36 | 13.1 | 115 | 2 | 3 | 10 | 17 | 31 | 36 | 44 |
| **TOT. SINGLE DX** | 55 | 5.2 | 21 | <1 | 1 | 3 | 7 | 11 | 16 | 21 |
| **" MULTIPLE DX** | 69 | 11.3 | 90 | 2 | 2 | 9 | 15 | 22 | 35 | 43 |
| TOTAL 0-19 YRS | 1 | 1.0 | | <1 | <1 | <1 | <1 | <1 | <1 | <1 |
| 20-34 | 16 | 3.8 | 14 | <1 | <1 | 2 | 6 | 8 | 12 | 14 |
| 35-49 | 18 | 7.4 | 44 | <1 | 1 | 5 | 10 | 19 | 20 | 21 |
| 50-64 | 39 | 7.5 | 51 | 1 | 1 | 6 | 11 | 15 | 20 | 38 |
| 65+ | 50 | 11.6 | 95 | 2 | 3 | 9 | 16 | 25 | 35 | 44 |
| **GRAND TOTAL** | **124** | **8.6** | **69** | **1** | **1** | **6** | **12** | **18** | **24** | **40** |

### 685: Miscellaneous operations on vulva and perineum (71.7-71.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 17 | 1.7 | 2 | <1 | <1 | 1 | 2 | 4 | 5 | 6 |
| 20-34 | 35 | 2.6 | 2 | 1 | 1 | 2 | 3 | 5 | 6 | 7 |
| 35-49 | 17 | 3.8 | 5 | <1 | 1 | 4 | 5 | 7 | 9 | 10 |
| 50-64 | 17 | 3.1 | 5 | <1 | 1 | 2 | 4 | 7 | 8 | 8 |
| 65+ | 6 | 4.3 | 15 | 1 | 1 | 3 | 7 | 11 | 11 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 30 | 2.2 | 11 | 1 | 1 | 1 | 2 | 4 | 6 | 19 |
| 20-34 | 38 | 3.6 | 8 | 1 | 1 | 3 | 4 | 7 | 9 | 17 |
| 35-49 | 18 | 4.4 | 13 | 1 | 1 | 3 | 6 | 7 | 13 | 17 |
| 50-64 | 6 | 2.5 | 1 | 1 | 1 | 3 | 3 | 4 | 4 | 4 |
| 65+ | 4 | 4.0 | 10 | 1 | 1 | 3 | 6 | 8 | 8 | 8 |
| **TOT. SINGLE DX** | 92 | 2.9 | 5 | <1 | 1 | 2 | 4 | 6 | 7 | 11 |
| **" MULTIPLE DX** | 96 | 3.3 | 10 | 1 | 1 | 2 | 4 | 7 | 8 | 18 |
| TOTAL 0-19 YRS | 47 | 2.0 | 8 | <1 | 1 | 1 | 2 | 4 | 6 | 19 |
| 20-34 | 73 | 3.1 | 6 | 1 | 1 | 3 | 3 | 6 | 7 | 15 |
| 35-49 | 35 | 4.1 | 10 | 1 | 1 | 3 | 6 | 7 | 9 | 17 |
| 50-64 | 23 | 2.9 | 4 | <1 | 1 | 2 | 4 | 6 | 8 | 8 |
| 65+ | 10 | 4.2 | 12 | 1 | 1 | 3 | 7 | 9 | 11 | 11 |
| **GRAND TOTAL** | **188** | **3.1** | **8** | **1** | **1** | **2** | **4** | **6** | **8** | **17** |

### 686: Miscellaneous diagnostic and therapeutic procedures on female genital system (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 11 | 1.1 | | <1 | <1 | <1 | 1 | 2 | 2 | 2 |
| 20-34 | 4 | 2.8 | 12 | 1 | 1 | 1 | 5 | 8 | 8 | 8 |
| 35-49 | 2 | 1.0 | | <1 | <1 | <1 | 1 | 1 | 1 | 1 |
| 50-64 | | | | | | | | | | |
| 65+ | 2 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 10.7 | 280 | 1 | 1 | 1 | 23 | 30 | 30 | 30 |
| 20-34 | 11 | 2.0 | 2 | <1 | <1 | 1 | 3 | 5 | 6 | 6 |
| 35-49 | 5 | 4.2 | 12 | 1 | 1 | 2 | 7 | 9 | 9 | 9 |
| 50-64 | 5 | 12.6 | 316 | 1 | 1 | 3 | 21 | 43 | 43 | 43 |
| 65+ | 2 | 19.5 | 481 | 4 | 4 | 19 | 35 | 35 | 35 | 35 |
| **TOT. SINGLE DX** | 19 | 2.1 | 6 | <1 | <1 | 1 | 1 | 7 | 8 | 8 |
| **" MULTIPLE DX** | 26 | 6.8 | 128 | <1 | 1 | 2 | 6 | 28 | 37 | 43 |
| TOTAL 0-19 YRS | 14 | 3.1 | 60 | <1 | <1 | 1 | 1 | 5 | 24 | 30 |
| 20-34 | 15 | 2.2 | 5 | <1 | <1 | 1 | 3 | 6 | 7 | 8 |
| 35-49 | 7 | 3.3 | 11 | <1 | <1 | 2 | 6 | 9 | 9 | 9 |
| 50-64 | 5 | 12.6 | 316 | 1 | 1 | 3 | 21 | 43 | 43 | 43 |
| 65+ | 4 | 13.3 | 212 | 4 | 4 | 7 | 21 | 35 | 35 | 35 |
| **GRAND TOTAL** | **45** | **4.8** | **81** | **<1** | **<1** | **1** | **4** | **9** | **31** | **43** |

### 687: Low forceps with and without episiotomy (72.0-72.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 857 | 2.3 | | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 20-34 | 5571 | 2.5 | | 1 | 1 | 2 | 3 | 4 | 4 | 5 |
| 35-49 | 309 | 2.8 | | 1 | 2 | 3 | 3 | 4 | 4 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1698 | 2.7 | 2 | 1 | 2 | 3 | 3 | 4 | 4 | 7 |
| 20-34 | 9482 | 2.8 | 2 | 1 | 2 | 3 | 3 | 4 | 5 | 7 |
| 35-49 | 619 | 3.2 | 4 | 2 | 2 | 3 | 4 | 4 | 5 | 10 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 6737 | 2.5 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 5 |
| **" MULTIPLE DX** | 11799 | 2.8 | 2 | 1 | 2 | 3 | 3 | 4 | 5 | 7 |
| TOTAL 0-19 YRS | 2555 | 2.6 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 7 |
| 20-34 | 15053 | 2.7 | 2 | 1 | 2 | 3 | 3 | 4 | 4 | 6 |
| 35-49 | 928 | 3.0 | 3 | 1 | 2 | 3 | 3 | 4 | 5 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **18536** | **2.7** | **2** | **1** | **2** | **3** | **3** | **4** | **4** | **6** |

### 688: Breech extraction (72.5-72.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 2.1 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 20-34 | 109 | 2.0 | | <1 | 1 | 2 | 2 | 3 | 4 | 5 |
| 35-49 | 5 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 3 | 3 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 44 | 2.4 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 5 |
| 20-34 | 367 | 2.7 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 35-49 | 35 | 2.3 | | <1 | 1 | 2 | 3 | 3 | 4 | 4 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 124 | 2.0 | 1 | <1 | 1 | 2 | 2 | 3 | 4 | 5 |
| **" MULTIPLE DX** | 446 | 2.6 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| TOTAL 0-19 YRS | 54 | 2.3 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 5 |
| 20-34 | 476 | 2.5 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 35-49 | 40 | 2.3 | 1 | <1 | 1 | 2 | 3 | 3 | 4 | 4 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **570** | **2.5** | **2** | **1** | **1** | **2** | **3** | **4** | **5** | **9** |

### 689: Other extraction procedures to assist delivery (72.2-72.4; 72.7-72.9; 73.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 185 | 2.4 | | 1 | 1 | 2 | 3 | 4 | 4 | 5 |
| 20-34 | 1100 | 2.6 | 1 | 1 | 1 | 3 | 3 | 4 | 4 | 6 |
| 35-49 | 63 | 2.6 | 1 | 1 | 1 | 2 | 3 | 4 | 5 | 5 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 555 | 2.8 | 1 | 1 | 2 | 3 | 3 | 4 | 4 | 6 |
| 20-34 | 3644 | 2.9 | 1 | 1 | 2 | 3 | 3 | 4 | 5 | 7 |
| 35-49 | 301 | 3.2 | 2 | 1 | 2 | 3 | 4 | 5 | 5 | 9 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 1348 | 2.6 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| **" MULTIPLE DX** | 4500 | 2.9 | 2 | 1 | 2 | 3 | 3 | 4 | 5 | 7 |
| TOTAL 0-19 YRS | 740 | 2.7 | 1 | 1 | 2 | 3 | 3 | 4 | 4 | 6 |
| 20-34 | 4744 | 2.8 | 2 | 1 | 2 | 3 | 3 | 4 | 5 | 7 |
| 35-49 | 364 | 3.1 | 2 | 1 | 2 | 3 | 4 | 4 | 5 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **5848** | **2.8** | **2** | **1** | **2** | **3** | **3** | **4** | **5** | **7** |

### 690: Other procedures to assist delivery (73.0-73.1; 73.3-73.6; 73.91-73.92)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 6405 | 2.1 | | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 20-34 | 42787 | 2.1 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 35-49 | 2418 | 2.3 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6943 | 2.4 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 7 |
| 20-34 | 46545 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 35-49 | 3279 | 2.7 | 2 | 1 | 1 | 3 | 3 | 4 | 5 | 7 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 51610 | 2.1 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| **" MULTIPLE DX** | 56767 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| TOTAL 0-19 YRS | 13348 | 2.2 | 1 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 20-34 | 89332 | 2.3 | 2 | 1 | 1 | 2 | 3 | 3 | 4 | 5 |
| 35-49 | 5697 | 2.5 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 7 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **108377** | **2.3** | **2** | **1** | **1** | **2** | **3** | **3** | **4** | **6** |

### 691: Low cervical cesarean section (74.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 677 | 4.5 | 1 | 3 | 3 | 4 | 5 | 6 | 6 | 10 |
| 20-34 | 6775 | 4.6 | 2 | 3 | 3 | 5 | 5 | 6 | 6 | 8 |
| 35-49 | 489 | 5.0 | 3 | 3 | 3 | 5 | 6 | 7 | 7 | 13 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2461 | 5.3 | 8 | 3 | 4 | 5 | 6 | 7 | 9 | 14 |
| 20-34 | 22744 | 5.2 | 8 | 3 | 3 | 5 | 6 | 7 | 8 | 15 |
| 35-49 | 2402 | 5.7 | 16 | 3 | 4 | 5 | 6 | 7 | 9 | 19 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 7941 | 4.6 | 3 | 3 | 3 | 5 | 5 | 6 | 6 | 8 |
| **" MULTIPLE DX** | 27607 | 5.2 | 9 | 3 | 4 | 5 | 6 | 7 | 8 | 15 |
| TOTAL 0-19 YRS | 3138 | 5.1 | 7 | 3 | 3 | 5 | 5 | 7 | 8 | 13 |
| 20-34 | 29519 | 5.0 | 7 | 3 | 3 | 5 | 5 | 6 | 7 | 13 |
| 35-49 | 2891 | 5.5 | 15 | 3 | 4 | 5 | 6 | 7 | 8 | 19 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **35548** | **5.1** | **8** | **3** | **3** | **5** | **5** | **6** | **8** | **14** |

### 692: Other cesarean section (74.0; 74.2; 74.4; 74.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARI-ANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 48 | 4.9 | 2 | 3 | 3 | 5 | 5 | 6 | 7 | 12 |
| 20-34 | 366 | 4.8 | 6 | 3 | 3 | 5 | 5 | 6 | 7 | 12 |
| 35-49 | 17 | 4.5 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 7 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 158 | 5.4 | 6 | 3 | 3 | 5 | 6 | 8 | 10 | 16 |
| 20-34 | 1196 | 5.3 | 14 | 3 | 3 | 5 | 6 | 7 | 8 | 17 |
| 35-49 | 140 | 5.9 | 17 | 3 | 4 | 5 | 6 | 8 | 12 | 30 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 431 | 4.8 | 6 | 3 | 3 | 5 | 5 | 6 | 7 | 12 |
| **" MULTIPLE DX** | 1494 | 5.3 | 14 | 3 | 3 | 5 | 6 | 7 | 8 | 17 |
| TOTAL 0-19 YRS | 206 | 5.3 | 6 | 3 | 3 | 5 | 6 | 7 | 9 | 16 |
| 20-34 | 1562 | 5.2 | 13 | 3 | 3 | 5 | 6 | 7 | 8 | 16 |
| 35-49 | 157 | 5.7 | 16 | 3 | 3 | 5 | 6 | 8 | 11 | 28 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **1925** | **5.2** | **12** | **3** | **3** | **5** | **6** | **7** | **8** | **16** |

### 693: Other abdominal operation to remove embryo or fetus (74.3; 74.91)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 8 | 4.3 | 1 | 2 | 2 | 4 | 5 | 6 | 6 | 6 |
| 20-34 | 48 | 4.2 | 1 | 3 | 3 | 4 | 5 | 6 | 6 | 6 |
| 35-49 | 1 | 3.0 | | 3 | 3 | 3 | 3 | 3 | 3 | 3 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 4.0 | | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 20-34 | 47 | 5.2 | 6 | 2 | 3 | 4 | 6 | 8 | 10 | 14 |
| 35-49 | 4 | 4.3 | 2 | 3 | 3 | 4 | 5 | 6 | 6 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 57 | 4.2 | 1 | 2 | 3 | 4 | 5 | 6 | 6 | 6 |
| **" MULTIPLE DX** | 53 | 5.1 | 6 | 2 | 3 | 4 | 6 | 8 | 9 | 14 |
| TOTAL 0-19 YRS | 10 | 4.2 | 1 | 2 | 3 | 4 | 5 | 6 | 6 | 6 |
| 20-34 | 95 | 4.7 | 4 | 2 | 3 | 4 | 5 | 7 | 8 | 14 |
| 35-49 | 5 | 4.0 | 2 | 3 | 3 | 3 | 5 | 6 | 6 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **110** | **4.6** | **3** | **2** | **3** | **4** | **5** | **6** | **8** | **13** |

### 694: Intra-amniotic injection for abortion (75.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 157 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 2 |
| 20-34 | 216 | 1.3 | | <1 | <1 | 1 | 1 | 2 | 2 | 5 |
| 35-49 | 23 | 1.3 | | <1 | <1 | 1 | 2 | 2 | 3 | 3 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 70 | 1.3 | | 1 | 1 | 1 | 1 | 2 | 2 | 4 |
| 20-34 | 117 | 1.7 | 2 | <1 | 1 | 1 | 2 | 3 | 4 | 9 |
| 35-49 | 27 | 1.6 | 1 | 1 | 1 | 1 | 2 | 3 | 4 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 396 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 4 |
| **" MULTIPLE DX** | 214 | 1.6 | 2 | 1 | 1 | 1 | 2 | 3 | 4 | 7 |
| TOTAL 0-19 YRS | 227 | 1.2 | | <1 | <1 | 1 | 1 | 2 | 2 | 3 |
| 20-34 | 333 | 1.4 | 1 | <1 | <1 | 1 | 1 | 2 | 3 | 7 |
| 35-49 | 50 | 1.5 | 1 | <1 | 1 | 1 | 2 | 3 | 3 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **610** | **1.3** | **1** | **<1** | **<1** | **1** | **1** | **2** | **3** | **6** |

### 695: Diagnostic amniocentesis (75.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 47 | 3.6 | 15 | 1 | 1 | 2 | 5 | 6 | 10 | 20 |
| 20-34 | 213 | 3.5 | 17 | <1 | 1 | 2 | 4 | 7 | 11 | 28 |
| 35-49 | 15 | 2.7 | 4 | <1 | 1 | 2 | 4 | 5 | 7 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 53 | 4.0 | 8 | 1 | 1 | 4 | 5 | 9 | 11 | 12 |
| 20-34 | 209 | 4.6 | 28 | <1 | 1 | 3 | 5 | 10 | 13 | 28 |
| 35-49 | 20 | 3.6 | 9 | <1 | 1 | 3 | 5 | 7 | 11 | 14 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 275 | 3.5 | 17 | <1 | 1 | 2 | 4 | 7 | 9 | 26 |
| **" MULTIPLE DX** | 282 | 4.4 | 24 | <1 | 1 | 3 | 5 | 9 | 12 | 25 |
| TOTAL 0-19 YRS | 100 | 3.8 | 11 | 1 | 1 | 3 | 5 | 7 | 10 | 19 |
| 20-34 | 422 | 4.1 | 24 | <1 | 1 | 3 | 5 | 8 | 12 | 28 |
| 35-49 | 35 | 3.2 | 7 | <1 | 1 | 3 | 5 | 6 | 8 | 14 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **557** | **4.0** | **20** | **<1** | **1** | **3** | **5** | **8** | **12** | **25** |

### 696: Intra-uterine procedures on fetus and amnion (75.2-75.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 364 | 1.9 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 8 |
| 20-34 | 1582 | 2.0 | 3 | <1 | <1 | 2 | 2 | 3 | 4 | 6 |
| 35-49 | 101 | 2.0 | 1 | <1 | <1 | 2 | 2 | 4 | 5 | 7 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 610 | 2.7 | 3 | 1 | 1 | 2 | 3 | 4 | 5 | 11 |
| 20-34 | 3247 | 2.7 | 3 | 1 | 1 | 3 | 3 | 4 | 5 | 8 |
| 35-49 | 245 | 3.1 | 12 | 1 | 1 | 3 | 3 | 4 | 6 | 17 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 2047 | 2.0 | 3 | <1 | <1 | 2 | 2 | 3 | 4 | 7 |
| **" MULTIPLE DX** | 4102 | 2.7 | 4 | 1 | 1 | 3 | 3 | 4 | 5 | 9 |
| TOTAL 0-19 YRS | 974 | 2.4 | 3 | <1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 4829 | 2.5 | 3 | <1 | 1 | 2 | 3 | 4 | 4 | 8 |
| 35-49 | 346 | 2.8 | 9 | <1 | <1 | 2 | 3 | 4 | 6 | 16 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **6149** | **2.5** | **4** | **<1** | **1** | **2** | **3** | **4** | **4** | **8** |

## 697: Miscellaneous obstetric procedures (73.8; 73.93-73.94; 73.99; 75.4-75.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1017 | 2.0 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 20-34 | 6435 | 2.0 | | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| 35-49 | 437 | 2.0 | | <1 | 1 | 2 | 2 | 3 | 3 | 5 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1824 | 2.4 | 6 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 20-34 | 11474 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 35-49 | 906 | 2.6 | 2 | 1 | 1 | 2 | 3 | 4 | 5 | 8 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | 7889 | 2.0 | 1 | 1 | 1 | 2 | 2 | 3 | 3 | 4 |
| **" MULTIPLE DX** | 14204 | 2.4 | 3 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| TOTAL 0-19 YRS | 2841 | 2.2 | 4 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 20-34 | 17909 | 2.2 | 2 | 1 | 1 | 2 | 3 | 3 | 4 | 6 |
| 35-49 | 1343 | 2.4 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 6 |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | **22093** | **2.3** | **2** | **1** | **1** | **2** | **3** | **3** | **4** | **6** |

## 698: Excision of facial bone (76.11; 76.2-76.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 42 | 2.6 | 3 | <1 | 1 | 2 | 3 | 5 | 7 | 9 |
| 20-34 | 42 | 2.8 | 5 | <1 | 1 | 2 | 3 | 4 | 7 | 15 |
| 35-49 | 23 | 3.7 | 6 | <1 | 1 | 3 | 5 | 8 | 9 | 10 |
| 50-64 | 32 | 4.1 | 13 | 1 | 1 | 3 | 5 | 9 | 12 | 16 |
| 65+ | 16 | 8.3 | 50 | 2 | 2 | 6 | 12 | 17 | 25 | 28 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 9 | 3.9 | 24 | <1 | <1 | 2 | 4 | 12 | 16 | 16 |
| 20-34 | 25 | 5.1 | 49 | 1 | 1 | 3 | 4 | 13 | 23 | 31 |
| 35-49 | 16 | 5.9 | 23 | 1 | 2 | 4 | 10 | 14 | 15 | 15 |
| 50-64 | 45 | 11.2 | 168 | 2 | 2 | 7 | 13 | 27 | 40 | 64 |
| 65+ | 41 | 15.8 | 235 | 2 | 2 | 11 | 22 | 41 | 61 | 99+ |
| **TOT. SINGLE DX** | 155 | 3.7 | 14 | 1 | 1 | 2 | 4 | 9 | 10 | 17 |
| **" MULTIPLE DX** | 136 | 10.4 | 157 | 1 | 2 | 5 | 13 | 27 | 37 | 71 |
| TOTAL 0-19 YRS | 51 | 2.9 | 7 | <1 | 1 | 2 | 3 | 6 | 9 | 16 |
| 20-34 | 67 | 3.7 | 23 | 1 | 1 | 2 | 3 | 7 | 13 | 29 |
| 35-49 | 39 | 4.6 | 14 | <1 | 1 | 3 | 6 | 11 | 13 | 15 |
| 50-64 | 77 | 8.2 | 116 | 1 | 1 | 4 | 10 | 19 | 27 | 60 |
| 65+ | 57 | 13.7 | 193 | 2 | 2 | 9 | 17 | 33 | 55 | 97 |
| **GRAND TOTAL** | **291** | **6.8** | **91** | **1** | **1** | **3** | **8** | **16** | **27** | **61** |

## 699: Repair and plastic operations on facial bones (76.4-76.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 288 | 3.4 | | 2 | 2 | 3 | 4 | 5 | 5 | 6 |
| 20-34 | 528 | 3.2 | 1 | 2 | 2 | 3 | 4 | 5 | 5 | 7 |
| 35-49 | 190 | 3.4 | 2 | 2 | 2 | 3 | 4 | 5 | 6 | 10 |
| 50-64 | 41 | 3.8 | 2 | 1 | 2 | 3 | 5 | 6 | 7 | 8 |
| 65+ | 13 | 6.0 | 28 | 2 | 2 | 4 | 6 | 15 | 20 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 224 | 3.9 | 2 | 2 | 2 | 4 | 4 | 6 | 7 | 10 |
| 20-34 | 335 | 4.1 | 3 | 2 | 2 | 4 | 5 | 6 | 7 | 12 |
| 35-49 | 132 | 4.6 | 15 | 2 | 2 | 4 | 5 | 7 | 9 | 19 |
| 50-64 | 43 | 7.0 | 67 | 1 | 2 | 4 | 6 | 25 | 29 | 32 |
| 65+ | 11 | 9.7 | 128 | 2 | 2 | 5 | 19 | 56 | 94 | 99+ |
| **TOT. SINGLE DX** | 1060 | 3.4 | 2 | 2 | 2 | 3 | 4 | 5 | 5 | 7 |
| **" MULTIPLE DX** | 745 | 4.4 | 11 | 2 | 2 | 4 | 5 | 6 | 8 | 25 |
| TOTAL 0-19 YRS | 512 | 3.6 | 2 | 2 | 2 | 3 | 4 | 5 | 6 | 8 |
| 20-34 | 863 | 3.6 | 2 | 2 | 2 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 322 | 3.9 | 8 | 2 | 2 | 3 | 4 | 6 | 7 | 13 |
| 50-64 | 84 | 5.4 | 38 | 1 | 2 | 4 | 5 | 8 | 25 | 31 |
| 65+ | 24 | 7.7 | 75 | 2 | 2 | 4 | 11 | 23 | 53 | 99+ |
| **GRAND TOTAL** | **1805** | **3.8** | **6** | **2** | **2** | **3** | **4** | **5** | **6** | **12** |

## 700: Closed reduction of maxillofacial fractures (76.70; 76.71; 76.73; 76.75; 76.78)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 57 | 2.1 | 2 | 1 | 1 | 2 | 2 | 4 | 5 | 10 |
| 20-34 | 87 | 2.1 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 8 |
| 35-49 | 31 | 2.5 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 5 |
| 50-64 | 5 | 2.0 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 65+ | 3 | 6.7 | 30 | 3 | 3 | 4 | 11 | 13 | 13 | 13 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 182 | 3.5 | 32 | 1 | 1 | 2 | 3 | 6 | 8 | 43 |
| 20-34 | 304 | 3.9 | 17 | 1 | 1 | 3 | 5 | 7 | 10 | 29 |
| 35-49 | 78 | 4.1 | 15 | 1 | 1 | 3 | 5 | 8 | 14 | 20 |
| 50-64 | 23 | 6.2 | 32 | <1 | 1 | 4 | 9 | 15 | 18 | 23 |
| 65+ | 17 | 6.5 | 13 | 2 | 2 | 6 | 9 | 12 | 13 | 13 |
| **TOT. SINGLE DX** | 183 | 2.2 | 2 | 1 | 1 | 2 | 3 | 4 | 4 | 10 |
| **" MULTIPLE DX** | 604 | 3.9 | 23 | 1 | 1 | 2 | 4 | 8 | 11 | 29 |
| TOTAL 0-19 YRS | 239 | 3.1 | 26 | 1 | 1 | 2 | 3 | 5 | 7 | 37 |
| 20-34 | 391 | 3.5 | 14 | 1 | 1 | 2 | 4 | 7 | 10 | 28 |
| 35-49 | 109 | 3.6 | 12 | 1 | 1 | 2 | 4 | 7 | 9 | 19 |
| 50-64 | 28 | 5.4 | 30 | 1 | 1 | 3 | 8 | 14 | 16 | 23 |
| 65+ | 20 | 6.5 | 15 | 2 | 2 | 6 | 9 | 12 | 13 | 13 |
| **GRAND TOTAL** | **787** | **3.5** | **18** | **1** | **1** | **2** | **4** | **7** | **10** | **25** |

## 701: Open reduction of maxillofacial fractures (76.72; 76.74; 76.76-76.77; 76.79)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 72 | 2.6 | 3 | 1 | 1 | 2 | 3 | 4 | 6 | 12 |
| 20-34 | 175 | 2.9 | 21 | <1 | 1 | 2 | 3 | 5 | 6 | 17 |
| 35-49 | 56 | 2.4 | 3 | 1 | 1 | 2 | 3 | 5 | 7 | 9 |
| 50-64 | 22 | 2.8 | 3 | <1 | 1 | 2 | 4 | 5 | 7 | 8 |
| 65+ | 12 | 4.8 | 24 | 1 | 2 | 3 | 4 | 14 | 17 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 254 | 4.3 | 21 | 1 | 1 | 3 | 5 | 9 | 12 | 24 |
| 20-34 | 684 | 5.0 | 43 | 1 | 1 | 3 | 5 | 10 | 15 | 34 |
| 35-49 | 249 | 5.1 | 34 | 1 | 1 | 3 | 6 | 12 | 18 | 36 |
| 50-64 | 94 | 7.0 | 132 | 1 | 1 | 4 | 6 | 15 | 26 | 71 |
| 65+ | 64 | 10.3 | 74 | 2 | 2 | 8 | 13 | 23 | 26 | 44 |
| **TOT. SINGLE DX** | 337 | 2.8 | 14 | <1 | 1 | 2 | 3 | 5 | 6 | 14 |
| **" MULTIPLE DX** | 1345 | 5.3 | 47 | 1 | 1 | 3 | 6 | 11 | 16 | 36 |
| TOTAL 0-19 YRS | 326 | 3.9 | 18 | 1 | 1 | 3 | 5 | 8 | 12 | 24 |
| 20-34 | 859 | 4.6 | 39 | 1 | 1 | 3 | 5 | 9 | 13 | 33 |
| 35-49 | 305 | 4.6 | 30 | 1 | 1 | 3 | 5 | 11 | 15 | 33 |
| 50-64 | 116 | 6.2 | 110 | 1 | 1 | 3 | 5 | 13 | 19 | 64 |
| 65+ | 76 | 9.4 | 70 | 2 | 2 | 8 | 12 | 21 | 25 | 43 |
| **GRAND TOTAL** | **1682** | **4.8** | **41** | **1** | **1** | **3** | **5** | **10** | **14** | **34** |

## 702: Miscellaneous maxillofacial surgery (76.0; 76.91-76.95; 76.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 2.6 | 1 | <1 | 1 | 3 | 3 | 4 | 5 | 5 |
| 20-34 | 23 | 3.9 | 9 | 1 | 1 | 3 | 5 | 9 | 10 | 12 |
| 35-49 | 13 | 4.6 | 8 | 1 | 1 | 5 | 6 | 9 | 10 | 10 |
| 50-64 | 15 | 4.4 | 3 | 2 | 2 | 4 | 6 | 7 | 8 | 8 |
| 65+ | 4 | 3.8 | 4 | 2 | 2 | 4 | 5 | 6 | 6 | 6 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 13 | 4.2 | 8 | 1 | 1 | 4 | 5 | 8 | 11 | 12 |
| 20-34 | 22 | 6.4 | 73 | <1 | 1 | 4 | 8 | 11 | 24 | 42 |
| 35-49 | 17 | 5.5 | 33 | 1 | 1 | 4 | 6 | 10 | 20 | 25 |
| 50-64 | 15 | 5.4 | 25 | 1 | 2 | 4 | 6 | 13 | 17 | 19 |
| 65+ | 7 | 9.3 | 237 | 1 | 1 | 4 | 6 | 36 | 44 | 44 |
| **TOT. SINGLE DX** | 71 | 3.8 | 6 | 1 | 1 | 3 | 5 | 8 | 9 | 12 |
| **" MULTIPLE DX** | 74 | 5.9 | 56 | 1 | 1 | 4 | 7 | 11 | 18 | 44 |
| TOTAL 0-19 YRS | 29 | 3.3 | 5 | 1 | 1 | 3 | 4 | 6 | 7 | 12 |
| 20-34 | 45 | 5.1 | 41 | 1 | 1 | 3 | 7 | 9 | 12 | 42 |
| 35-49 | 30 | 5.1 | 22 | 1 | 1 | 4 | 6 | 10 | 10 | 25 |
| 50-64 | 30 | 4.9 | 14 | 2 | 2 | 4 | 6 | 9 | 13 | 19 |
| 65+ | 11 | 7.3 | 151 | 1 | 1 | 4 | 6 | 21 | 42 | 44 |
| **GRAND TOTAL** | **145** | **4.9** | **32** | **1** | **1** | **4** | **6** | **9** | **12** | **42** |

## 703: Incision and division of bones except facial (77.0-77.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 278 | 4.8 | 33 | 1 | 1 | 3 | 5 | 9 | 15 | 35 |
| 20-34 | 262 | 4.4 | 16 | 1 | 1 | 3 | 5 | 9 | 11 | 23 |
| 35-49 | 186 | 4.7 | 23 | <1 | 1 | 4 | 6 | 8 | 10 | 33 |
| 50-64 | 217 | 5.0 | 10 | 1 | 1 | 4 | 6 | 9 | 11 | 17 |
| 65+ | 73 | 7.6 | 23 | 2 | 2 | 7 | 9 | 13 | 19 | 23 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 179 | 6.8 | 61 | 1 | 1 | 4 | 7 | 16 | 25 | 40 |
| 20-34 | 250 | 6.5 | 80 | 1 | 1 | 4 | 7 | 11 | 27 | 43 |
| 35-49 | 236 | 5.6 | 28 | 1 | 1 | 4 | 7 | 11 | 16 | 33 |
| 50-64 | 303 | 7.1 | 75 | 1 | 2 | 5 | 8 | 13 | 20 | 64 |
| 65+ | 167 | 12.1 | 207 | 2 | 3 | 7 | 14 | 27 | 44 | 92 |
| **TOT. SINGLE DX** | 1016 | 4.9 | 22 | 1 | 1 | 4 | 6 | 9 | 12 | 24 |
| **" MULTIPLE DX** | 1135 | 7.4 | 88 | 1 | 2 | 4 | 8 | 15 | 25 | 55 |
| TOTAL 0-19 YRS | 457 | 5.6 | 45 | 1 | 1 | 4 | 6 | 12 | 20 | 39 |
| 20-34 | 512 | 5.4 | 49 | 1 | 1 | 4 | 6 | 10 | 17 | 39 |
| 35-49 | 422 | 5.2 | 26 | 1 | 1 | 4 | 6 | 10 | 14 | 33 |
| 50-64 | 520 | 6.2 | 50 | 1 | 2 | 5 | 7 | 11 | 16 | 52 |
| 65+ | 240 | 10.7 | 155 | 2 | 2 | 7 | 12 | 22 | 38 | 87 |
| **GRAND TOTAL** | **2151** | **6.2** | **58** | **1** | **1** | **4** | **7** | **12** | **19** | **43** |

## 704: Bunionectomy (77.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 184 | 2.9 | 1 | 1 | 1 | 3 | 4 | 5 | 6 | 6 |
| 20-34 | 329 | 2.9 | 2 | <1 | 1 | 3 | 4 | 5 | 5 | 7 |
| 35-49 | 325 | 3.2 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 50-64 | 430 | 3.5 | 3 | 1 | 1 | 3 | 4 | 6 | 7 | 9 |
| 65+ | 219 | 3.6 | 2 | 1 | 2 | 3 | 5 | 6 | 7 | 9 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 222 | 3.4 | 2 | 1 | 2 | 3 | 4 | 5 | 6 | 8 |
| 20-34 | 460 | 3.6 | 11 | 1 | 2 | 3 | 4 | 5 | 6 | 10 |
| 35-49 | 616 | 3.7 | 8 | 1 | 2 | 3 | 4 | 6 | 7 | 9 |
| 50-64 | 1156 | 4.4 | 20 | 2 | 2 | 4 | 5 | 7 | 9 | 16 |
| 65+ | 743 | 4.7 | 15 | 2 | 2 | 4 | 5 | 7 | 9 | 23 |
| **TOT. SINGLE DX** | 1487 | 3.3 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| **" MULTIPLE DX** | 3197 | 4.1 | 15 | 1 | 2 | 4 | 5 | 6 | 8 | 14 |
| TOTAL 0-19 YRS | 406 | 3.2 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 7 |
| 20-34 | 789 | 3.3 | 8 | 1 | 1 | 3 | 4 | 5 | 6 | 9 |
| 35-49 | 941 | 3.5 | 7 | 1 | 2 | 3 | 4 | 5 | 7 | 9 |
| 50-64 | 1586 | 4.2 | 16 | 1 | 2 | 4 | 5 | 6 | 8 | 14 |
| 65+ | 962 | 4.4 | 13 | 2 | 2 | 4 | 5 | 7 | 8 | 19 |
| **GRAND TOTAL** | **4684** | **3.9** | **11** | **1** | **2** | **3** | **5** | **6** | **7** | **12** |

## 705: Excision of bone except facial (77.4; 77.6-77.9; 79.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 999 | 3.2 | 18 | <1 | 1 | 2 | 3 | 5 | 8 | 25 |
| 20-34 | 1366 | 3.3 | 14 | <1 | 1 | 2 | 4 | 6 | 8 | 19 |
| 35-49 | 886 | 3.5 | 15 | <1 | 1 | 3 | 4 | 7 | 9 | 21 |
| 50-64 | 614 | 3.8 | 18 | <1 | 1 | 3 | 4 | 7 | 10 | 24 |
| 65+ | 320 | 4.7 | 25 | <1 | 1 | 3 | 6 | 10 | 14 | 28 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 565 | 7.1 | 140 | 1 | 1 | 3 | 6 | 14 | 32 | 65 |
| 20-34 | 1283 | 6.9 | 109 | 1 | 1 | 3 | 7 | 16 | 29 | 57 |
| 35-49 | 992 | 6.8 | 122 | 1 | 1 | 3 | 6 | 15 | 30 | 74 |
| 50-64 | 1039 | 7.4 | 102 | 1 | 1 | 4 | 8 | 17 | 28 | 60 |
| 65+ | 843 | 12.6 | 223 | 1 | 2 | 7 | 16 | 31 | 47 | 71 |
| **TOT. SINGLE DX** | 4185 | 3.5 | 17 | <1 | 1 | 2 | 4 | 6 | 9 | 23 |
| **" MULTIPLE DX** | 4722 | 8.0 | 139 | 1 | 1 | 4 | 8 | 20 | 33 | 66 |
| TOTAL 0-19 YRS | 1564 | 4.6 | 66 | <1 | 1 | 2 | 4 | 8 | 14 | 45 |
| 20-34 | 2649 | 5.0 | 63 | <1 | 1 | 3 | 5 | 9 | 18 | 44 |
| 35-49 | 1878 | 5.2 | 75 | 1 | 1 | 3 | 5 | 10 | 18 | 48 |
| 50-64 | 1653 | 6.0 | 74 | 1 | 1 | 3 | 6 | 13 | 22 | 50 |
| 65+ | 1163 | 10.4 | 181 | 1 | 1 | 5 | 12 | 25 | 39 | 69 |
| **GRAND TOTAL** | **8907** | **5.9** | **87** | **1** | **1** | **3** | **5** | **13** | **23** | **54** |

## 706: Repair and plastic operation on bone except facial (78.0-78.5; 78.7-78.9, 79.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 312 | 5.2 | 37 | 1 | 1 | 4 | 6 | 9 | 13 | 45 |
| 20-34 | 351 | 4.3 | 13 | 1 | 1 | 4 | 5 | 8 | 10 | 21 |
| 35-49 | 141 | 4.7 | 13 | 1 | 1 | 4 | 6 | 10 | 13 | 17 |
| 50-64 | 71 | 7.3 | 33 | 2 | 3 | 5 | 9 | 13 | 16 | 35 |
| 65+ | 113 | 10.7 | 48 | 2 | 3 | 10 | 14 | 19 | 23 | 41 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 207 | 9.1 | 129 | 1 | 1 | 4 | 11 | 23 | 32 | 58 |
| 20-34 | 246 | 9.0 | 150 | 1 | 2 | 5 | 10 | 21 | 31 | 62 |
| 35-49 | 131 | 11.2 | 259 | 1 | 2 | 6 | 12 | 25 | 63 | 99+ |
| 50-64 | 165 | 13.1 | 167 | 2 | 3 | 8 | 16 | 29 | 42 | 61 |
| 65+ | 426 | 15.8 | 118 | 5 | 7 | 13 | 19 | 29 | 41 | 80 |
| **TOT. SINGLE DX** | 988 | 5.6 | 30 | 1 | 1 | 4 | 7 | 11 | 15 | 28 |
| **" MULTIPLE DX** | 1175 | 12.3 | 158 | 2 | 2 | 9 | 16 | 27 | 40 | 71 |
| TOTAL 0-19 YRS | 519 | 6.7 | 77 | 1 | 1 | 4 | 7 | 16 | 24 | 55 |
| 20-34 | 597 | 6.2 | 75 | 1 | 1 | 4 | 6 | 12 | 20 | 45 |
| 35-49 | 272 | 7.9 | 142 | 1 | 2 | 4 | 8 | 15 | 25 | 93 |
| 50-64 | 236 | 11.4 | 134 | 2 | 3 | 7 | 13 | 26 | 38 | 59 |
| 65+ | 539 | 14.7 | 108 | 3 | 6 | 12 | 18 | 27 | 39 | 67 |
| **GRAND TOTAL** | **2163** | **9.2** | **111** | **1** | **2** | **6** | **12** | **21** | **29** | **61** |

## 707: Removal of internal fixation device (76.97; 78.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 714 | 2.2 | 7 | <1 | <1 | 2 | 2 | 4 | 5 | 16 |
| 20-34 | 1261 | 2.2 | 4 | <1 | <1 | 2 | 3 | 4 | 6 | 11 |
| 35-49 | 562 | 2.5 | 8 | <1 | <1 | 2 | 3 | 4 | 6 | 14 |
| 50-64 | 396 | 3.1 | 13 | <1 | <1 | 2 | 3 | 6 | 9 | 17 |
| 65+ | 317 | 4.8 | 37 | <1 | 1 | 3 | 5 | 11 | 16 | 30 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 185 | 4.4 | 48 | <1 | <1 | 2 | 4 | 9 | 21 | 34 |
| 20-34 | 413 | 5.4 | 62 | <1 | 1 | 3 | 6 | 13 | 21 | 42 |
| 35-49 | 218 | 4.9 | 53 | <1 | 1 | 3 | 5 | 10 | 16 | 36 |
| 50-64 | 241 | 7.7 | 122 | 1 | 1 | 4 | 9 | 15 | 24 | 62 |
| 65+ | 363 | 12.0 | 185 | 2 | 2 | 8 | 15 | 26 | 39 | 77 |
| **TOT. SINGLE DX** | 3250 | 2.6 | 11 | <1 | <1 | 2 | 3 | 5 | 7 | 16 |
| **" MULTIPLE DX** | 1420 | 7.3 | 109 | <1 | 1 | 4 | 8 | 17 | 25 | 56 |
| TOTAL 0-19 YRS | 899 | 2.7 | 16 | <1 | <1 | 2 | 3 | 4 | 6 | 26 |
| 20-34 | 1674 | 3.0 | 21 | <1 | <1 | 2 | 3 | 6 | 9 | 23 |
| 35-49 | 780 | 3.1 | 22 | <1 | <1 | 2 | 3 | 6 | 9 | 22 |
| 50-64 | 637 | 4.8 | 59 | <1 | 1 | 2 | 5 | 11 | 15 | 45 |
| 65+ | 680 | 8.6 | 129 | <1 | 1 | 4 | 11 | 19 | 30 | 57 |
| **GRAND TOTAL** | **4670** | **4.0** | **45** | **<1** | **<1** | **2** | **4** | **8** | **14** | **36** |

## 708: Closed reduction of fracture of femur with internal fixation (79.15)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 29 | 14.4 | 82 | 2 | 4 | 13 | 21 | 26 | 30 | 39 |
| 20-34 | 20 | 12.5 | 87 | 4 | 5 | 9 | 14 | 25 | 36 | 42 |
| 35-49 | 11 | 10.0 | 57 | 3 | 4 | 7 | 12 | 21 | 29 | 30 |
| 50-64 | 29 | 13.1 | 192 | 4 | 5 | 11 | 13 | 18 | 43 | 75 |
| 65+ | 99 | 12.6 | 27 | 5 | 6 | 12 | 15 | 19 | 23 | 29 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 100 | 18.7 | 147 | 3 | 5 | 16 | 24 | 38 | 44 | 49 |
| 20-34 | 97 | 18.6 | 215 | 6 | 8 | 14 | 20 | 39 | 52 | 77 |
| 35-49 | 37 | 19.9 | 256 | 5 | 6 | 14 | 25 | 52 | 57 | 99+ |
| 50-64 | 100 | 15.7 | 137 | 5 | 6 | 12 | 20 | 30 | 47 | 85 |
| 65+ | 522 | 16.4 | 93 | 6 | 8 | 14 | 19 | 27 | 35 | 62 |
| **TOT. SINGLE DX** | 180 | 12.8 | 69 | 4 | 5 | 11 | 15 | 21 | 27 | 42 |
| **" MULTIPLE DX** | 856 | 17.0 | 127 | 6 | 7 | 14 | 20 | 31 | 45 | 63 |
| TOTAL 0-19 YRS | 129 | 17.7 | 135 | 3 | 5 | 16 | 23 | 37 | 42 | 49 |
| 20-34 | 117 | 17.5 | 198 | 6 | 7 | 13 | 18 | 38 | 51 | 71 |
| 35-49 | 48 | 17.7 | 226 | 5 | 5 | 12 | 20 | 50 | 53 | 99+ |
| 50-64 | 129 | 15.1 | 150 | 5 | 6 | 12 | 17 | 29 | 47 | 80 |
| 65+ | 621 | 15.8 | 85 | 6 | 8 | 14 | 18 | 26 | 33 | 60 |
| **GRAND TOTAL** | **1044** | **16.2** | **119** | **5** | **7** | **14** | **19** | **29** | **41** | **62** |

## 709: Open reduction of fracture of femur with internal fixation (79.35; 79.55)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 59 | 11.1 | 131 | 2 | 3 | 8 | 13 | 20 | 38 | 58 |
| 20-34 | 56 | 10.3 | 39 | 3 | 3 | 9 | 13 | 20 | 24 | 28 |
| 35-49 | 35 | 12.7 | 149 | 3 | 5 | 9 | 13 | 22 | 47 | 59 |
| 50-64 | 105 | 10.8 | 20 | 5 | 6 | 10 | 13 | 16 | 18 | 28 |
| 65+ | 588 | 14.3 | 38 | 6 | 8 | 13 | 17 | 22 | 26 | 37 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 174 | 14.8 | 121 | 4 | 5 | 12 | 18 | 30 | 36 | 55 |
| 20-34 | 308 | 16.4 | 174 | 5 | 6 | 12 | 20 | 32 | 43 | 77 |
| 35-49 | 218 | 15.7 | 131 | 5 | 6 | 13 | 20 | 28 | 35 | 65 |
| 50-64 | 554 | 15.3 | 82 | 6 | 7 | 13 | 18 | 27 | 34 | 61 |
| 65+ | 3862 | 17.5 | 107 | 7 | 9 | 15 | 20 | 29 | 38 | 63 |
| **TOT. SINGLE DX** | 843 | 13.3 | 50 | 5 | 6 | 12 | 16 | 21 | 25 | 38 |
| **" MULTIPLE DX** | 5116 | 17.1 | 111 | 7 | 8 | 14 | 20 | 29 | 38 | 64 |
| TOTAL 0-19 YRS | 233 | 13.9 | 126 | 3 | 4 | 11 | 17 | 27 | 36 | 58 |
| 20-34 | 364 | 15.4 | 158 | 5 | 6 | 12 | 19 | 30 | 40 | 73 |
| 35-49 | 253 | 15.3 | 134 | 5 | 6 | 12 | 20 | 28 | 35 | 59 |
| 50-64 | 659 | 14.6 | 75 | 6 | 7 | 13 | 18 | 25 | 32 | 60 |
| 65+ | 4450 | 17.1 | 99 | 7 | 9 | 15 | 20 | 28 | 37 | 61 |
| **GRAND TOTAL** | **5959** | **16.5** | **104** | **6** | **8** | **14** | **19** | **28** | **36** | **62** |

## 710: Reduction of fracture of femur without internal fixation (79.05; 79.25; 79.45)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 82 | 10.1 | 112 | 1 | 2 | 6 | 13 | 26 | 30 | 53 |
| 20-34 | 11 | 17.5 | 781 | <1 | 2 | 7 | 22 | 54 | 94 | 98 |
| 35-49 | 2 | 7.5 | 25 | 4 | 4 | 7 | 11 | 11 | 11 | 11 |
| 50-64 | 7 | 16.3 | 238 | 2 | 2 | 10 | 26 | 41 | 44 | 44 |
| 65+ | 29 | 17.0 | 277 | 5 | 6 | 12 | 17 | 49 | 61 | 65 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 184 | 13.3 | 178 | 1 | 1 | 8 | 19 | 34 | 39 | 54 |
| 20-34 | 40 | 16.6 | 297 | 1 | 2 | 12 | 29 | 51 | 88 | 99+ |
| 35-49 | 17 | 19.9 | 538 | 3 | 3 | 9 | 21 | 62 | 70 | 74 |
| 50-64 | 31 | 19.4 | 400 | 1 | 2 | 10 | 27 | 58 | 61 | 65 |
| 65+ | 105 | 18.6 | 221 | 3 | 5 | 13 | 22 | 44 | 55 | 82 |
| **TOT. SINGLE DX** | 131 | 12.5 | 213 | 1 | 2 | 8 | 15 | 27 | 48 | 71 |
| **" MULTIPLE DX** | 377 | 15.9 | 241 | 1 | 2 | 11 | 22 | 39 | 55 | 79 |
| TOTAL 0-19 YRS | 266 | 12.3 | 160 | 1 | 2 | 7 | 17 | 30 | 37 | 54 |
| 20-34 | 51 | 16.8 | 389 | 1 | 2 | 11 | 25 | 51 | 95 | 99+ |
| 35-49 | 19 | 18.6 | 495 | 3 | 3 | 9 | 16 | 62 | 69 | 74 |
| 50-64 | 38 | 18.8 | 364 | 1 | 2 | 10 | 28 | 54 | 60 | 65 |
| 65+ | 134 | 18.3 | 232 | 3 | 5 | 13 | 20 | 47 | 55 | 73 |
| **GRAND TOTAL** | **508** | **15.1** | **235** | **1** | **2** | **10** | **19** | **37** | **52** | **78** |

## 711: Closed reduction of fracture with internal fixation, except maxillofacial and femur (79.10-79.14; 79.16-79.19)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 143 | 3.9 | 26 | <1 | 1 | 2 | 3 | 9 | 18 | 22 |
| 20-34 | 144 | 2.6 | 5 | <1 | <1 | 2 | 3 | 5 | 7 | 13 |
| 35-49 | 78 | 2.6 | 3 | <1 | <1 | 2 | 4 | 5 | 6 | 8 |
| 50-64 | 74 | 3.7 | 30 | <1 | 1 | 2 | 4 | 7 | 10 | 37 |
| 65+ | 62 | 3.7 | 32 | 1 | 1 | 3 | 4 | 6 | 8 | 41 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 219 | 5.8 | 50 | 1 | 1 | 3 | 8 | 15 | 20 | 35 |
| 20-34 | 253 | 6.7 | 117 | <1 | 1 | 3 | 7 | 14 | 24 | 64 |
| 35-49 | 136 | 5.4 | 67 | 1 | 1 | 3 | 6 | 11 | 15 | 52 |
| 50-64 | 157 | 7.3 | 145 | 1 | 1 | 4 | 6 | 17 | 27 | 68 |
| 65+ | 169 | 10.5 | 142 | 1 | 2 | 6 | 14 | 23 | 31 | 59 |
| **TOT. SINGLE DX** | 501 | 3.3 | 18 | <1 | 1 | 2 | 4 | 6 | 9 | 21 |
| **" MULTIPLE DX** | 934 | 7.1 | 106 | 1 | 1 | 3 | 8 | 17 | 25 | 56 |
| TOTAL 0-19 YRS | 362 | 5.0 | 42 | <1 | 1 | 2 | 5 | 14 | 19 | 32 |
| 20-34 | 397 | 5.2 | 81 | <1 | 1 | 3 | 5 | 11 | 17 | 59 |
| 35-49 | 214 | 4.4 | 46 | <1 | 1 | 3 | 5 | 9 | 12 | 51 |
| 50-64 | 231 | 6.2 | 111 | 1 | 1 | 3 | 6 | 13 | 25 | 54 |
| 65+ | 231 | 8.7 | 121 | 1 | 1 | 4 | 11 | 20 | 29 | 55 |
| **GRAND TOTAL** | **1435** | **5.7** | **79** | **<1** | **1** | **3** | **6** | **13** | **20** | **51** |

## 712: Open reduction of fracture with internal fixation, except maxillofacial and femur (79.30-79.34; 79.36-79.39; 79.50-79.54; 79.56-79.59)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 838 | 2.9 | 6 | <1 | 1 | 2 | 4 | 5 | 7 | 14 |
| 20-34 | 1181 | 3.4 | 6 | <1 | 1 | 3 | 4 | 6 | 8 | 13 |
| 35-49 | 613 | 4.0 | 7 | 1 | 1 | 4 | 5 | 7 | 9 | 14 |
| 50-64 | 413 | 5.1 | 13 | 1 | 2 | 4 | 6 | 9 | 11 | 20 |
| 65+ | 279 | 6.5 | 21 | 2 | 2 | 5 | 8 | 13 | 15 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1371 | 4.4 | 35 | 1 | 1 | 3 | 5 | 8 | 13 | 32 |
| 20-34 | 2409 | 5.9 | 70 | 1 | 1 | 4 | 6 | 11 | 19 | 50 |
| 35-49 | 1287 | 5.9 | 53 | 1 | 1 | 4 | 6 | 10 | 16 | 36 |
| 50-64 | 1219 | 7.1 | 63 | 1 | 2 | 5 | 8 | 13 | 19 | 43 |
| 65+ | 1017 | 10.0 | 80 | 2 | 3 | 7 | 13 | 20 | 28 | 48 |
| **TOT. SINGLE DX** | 3324 | 3.9 | 10 | 1 | 1 | 3 | 5 | 7 | 10 | 17 |
| **" MULTIPLE DX** | 7303 | 6.4 | 64 | 1 | 1 | 4 | 7 | 13 | 19 | 45 |
| TOTAL 0-19 YRS | 2209 | 3.8 | 25 | 1 | 1 | 3 | 4 | 7 | 10 | 28 |
| 20-34 | 3590 | 5.1 | 51 | 1 | 1 | 3 | 5 | 9 | 14 | 40 |
| 35-49 | 1900 | 5.3 | 40 | 1 | 1 | 4 | 6 | 9 | 13 | 32 |
| 50-64 | 1632 | 6.6 | 51 | 1 | 2 | 5 | 7 | 12 | 17 | 40 |
| 65+ | 1296 | 9.3 | 70 | 2 | 3 | 7 | 12 | 19 | 26 | 46 |
| **GRAND TOTAL** | **10627** | **5.6** | **48** | **1** | **1** | **4** | **6** | **11** | **16** | **36** |

### 713: Other reduction of fracture without internal fixation, except maxillofacial and femur (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1114 | 2.0 | 6 | <1 | <1 | 1 | 2 | 4 | 6 | 14 |
| 20-34 | 278 | 3.1 | 9 | <1 | 1 | 2 | 4 | 6 | 8 | 20 |
| 35-49 | 192 | 3.4 | 19 | <1 | 1 | 2 | 4 | 7 | 9 | 28 |
| 50-64 | 185 | 3.2 | 6 | 1 | 1 | 2 | 4 | 6 | 8 | 13 |
| 65+ | 255 | 3.9 | 15 | <1 | 1 | 2 | 5 | 8 | 11 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1665 | 3.0 | 31 | <1 | 1 | 1 | 3 | 6 | 10 | 27 |
| 20-34 | 780 | 5.8 | 84 | 1 | 1 | 3 | 6 | 11 | 21 | 53 |
| 35-49 | 457 | 5.3 | 48 | 1 | 1 | 3 | 6 | 11 | 18 | 39 |
| 50-64 | 525 | 5.7 | 53 | 1 | 1 | 3 | 7 | 11 | 20 | 39 |
| 65+ | 774 | 7.8 | 82 | 1 | 1 | 5 | 10 | 18 | 25 | 47 |
| **TOT. SINGLE DX** | 2024 | 2.6 | 10 | <1 | 1 | 2 | 3 | 5 | 8 | 17 |
| **" MULTIPLE DX** | 4201 | 5.0 | 58 | 1 | 1 | 3 | 5 | 11 | 17 | 40 |
| TOTAL 0-19 YRS | 2779 | 2.6 | 21 | <1 | 1 | 1 | 2 | 5 | 8 | 20 |
| 20-34 | 1058 | 5.1 | 66 | 1 | 1 | 3 | 5 | 10 | 17 | 44 |
| 35-49 | 649 | 4.7 | 41 | 1 | 1 | 3 | 5 | 9 | 14 | 38 |
| 50-64 | 710 | 5.0 | 43 | 1 | 1 | 3 | 6 | 10 | 15 | 38 |
| 65+ | 1029 | 6.8 | 69 | 1 | 1 | 4 | 8 | 16 | 21 | 42 |
| **GRAND TOTAL** | **6225** | **4.2** | **44** | **<1** | **1** | **2** | **5** | **9** | **14** | **36** |

### 714: Closed reduction of dislocation (79.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 114 | 5.2 | 58 | <1 | <1 | 2 | 7 | 14 | 21 | 36 |
| 20-34 | 104 | 2.6 | 11 | <1 | <1 | 2 | 3 | 4 | 6 | 22 |
| 35-49 | 63 | 4.0 | 45 | <1 | 1 | 2 | 3 | 11 | 17 | 40 |
| 50-64 | 79 | 3.5 | 18 | <1 | 1 | 2 | 3 | 8 | 14 | 22 |
| 65+ | 110 | 4.4 | 24 | <1 | 1 | 2 | 5 | 12 | 14 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 157 | 5.4 | 51 | <1 | 1 | 3 | 7 | 16 | 22 | 44 |
| 20-34 | 189 | 6.2 | 109 | <1 | 1 | 2 | 7 | 14 | 22 | 62 |
| 35-49 | 103 | 5.1 | 108 | <1 | 1 | 2 | 5 | 11 | 16 | 59 |
| 50-64 | 107 | 8.5 | 156 | 1 | 1 | 4 | 9 | 21 | 32 | 68 |
| 65+ | 240 | 7.3 | 75 | 1 | 1 | 4 | 10 | 17 | 27 | 44 |
| **TOT. SINGLE DX** | 470 | 4.0 | 32 | <1 | <1 | 2 | 4 | 11 | 15 | 25 |
| **" MULTIPLE DX** | 796 | 6.5 | 95 | 1 | 1 | 3 | 8 | 15 | 24 | 62 |
| TOTAL 0-19 YRS | 271 | 5.3 | 54 | <1 | 1 | 2 | 7 | 16 | 21 | 43 |
| 20-34 | 293 | 4.9 | 78 | <1 | <1 | 2 | 4 | 12 | 17 | 50 |
| 35-49 | 166 | 4.7 | 84 | <1 | 1 | 2 | 5 | 11 | 16 | 40 |
| 50-64 | 186 | 6.4 | 104 | <1 | 1 | 3 | 6 | 16 | 24 | 63 |
| 65+ | 350 | 6.4 | 61 | 1 | 1 | 4 | 8 | 14 | 23 | 41 |
| **GRAND TOTAL** | **1266** | **5.6** | **73** | **<1** | **1** | **3** | **6** | **14** | **21** | **43** |

### 715: Open reduction of dislocation (79.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 80 | 3.9 | 19 | 1 | 1 | 3 | 4 | 7 | 10 | 29 |
| 20-34 | 179 | 3.0 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 35-49 | 47 | 3.3 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 50-64 | 26 | 5.0 | 14 | <1 | 1 | 4 | 6 | 10 | 14 | 16 |
| 65+ | 26 | 10.9 | 156 | 2 | 3 | 7 | 12 | 22 | 36 | 63 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 75 | 4.1 | 21 | 1 | 1 | 3 | 5 | 9 | 13 | 29 |
| 20-34 | 197 | 4.3 | 35 | 1 | 1 | 3 | 4 | 7 | 13 | 52 |
| 35-49 | 85 | 3.8 | 8 | 1 | 1 | 3 | 5 | 7 | 9 | 16 |
| 50-64 | 60 | 9.6 | 325 | 1 | 1 | 5 | 8 | 20 | 66 | 99+ |
| 65+ | 59 | 12.7 | 180 | 1 | 1 | 10 | 18 | 25 | 33 | 80 |
| **TOT. SINGLE DX** | 358 | 3.9 | 22 | 1 | 1 | 3 | 4 | 6 | 10 | 22 |
| **" MULTIPLE DX** | 476 | 5.9 | 92 | 1 | 1 | 3 | 6 | 14 | 19 | 78 |
| TOTAL 0-19 YRS | 155 | 3.9 | 20 | 1 | 1 | 3 | 5 | 8 | 10 | 32 |
| 20-34 | 376 | 3.6 | 21 | 1 | 1 | 3 | 4 | 6 | 9 | 22 |
| 35-49 | 132 | 3.6 | 6 | 1 | 1 | 3 | 4 | 6 | 8 | 15 |
| 50-64 | 86 | 8.2 | 235 | 1 | 1 | 4 | 8 | 16 | 33 | 99 |
| 65+ | 85 | 12.1 | 172 | 1 | 2 | 9 | 17 | 23 | 31 | 77 |
| **GRAND TOTAL** | **834** | **5.0** | **63** | **1** | **1** | **3** | **5** | **10** | **16** | **41** |

### 716: Arthrotomy and division of joint capsule (80.0-80.1; 80.4)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 717 | 3.1 | 16 | <1 | 1 | 2 | 3 | 5 | 8 | 23 |
| 20-34 | 754 | 2.8 | 5 | <1 | 1 | 2 | 3 | 5 | 7 | 15 |
| 35-49 | 320 | 3.2 | 9 | <1 | 1 | 2 | 4 | 6 | 8 | 16 |
| 50-64 | 212 | 5.2 | 66 | <1 | 1 | 3 | 5 | 10 | 14 | 48 |
| 65+ | 97 | 8.6 | 72 | 1 | 2 | 6 | 12 | 19 | 23 | 48 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 451 | 4.6 | 36 | 1 | 1 | 3 | 5 | 10 | 16 | 29 |
| 20-34 | 637 | 4.2 | 49 | <1 | 1 | 3 | 4 | 8 | 12 | 23 |
| 35-49 | 358 | 5.2 | 62 | 1 | 1 | 3 | 5 | 10 | 17 | 41 |
| 50-64 | 284 | 8.7 | 140 | 1 | 1 | 4 | 9 | 23 | 38 | 61 |
| 65+ | 298 | 15.4 | 236 | 1 | 3 | 9 | 23 | 39 | 50 | 78 |
| **TOT. SINGLE DX** | 2100 | 3.5 | 21 | <1 | 1 | 2 | 4 | 6 | 10 | 22 |
| **" MULTIPLE DX** | 2028 | 6.8 | 104 | 1 | 1 | 3 | 6 | 17 | 26 | 55 |
| TOTAL 0-19 YRS | 1168 | 3.7 | 25 | <1 | 1 | 2 | 4 | 7 | 11 | 28 |
| 20-34 | 1391 | 3.5 | 26 | <1 | 1 | 2 | 4 | 6 | 9 | 19 |
| 35-49 | 678 | 4.2 | 39 | <1 | 1 | 3 | 4 | 8 | 12 | 30 |
| 50-64 | 496 | 7.2 | 112 | 1 | 1 | 4 | 7 | 18 | 28 | 57 |
| 65+ | 395 | 13.8 | 205 | 1 | 2 | 8 | 19 | 35 | 48 | 74 |
| **GRAND TOTAL** | **4128** | **5.1** | **64** | **<1** | **1** | **3** | **5** | **10** | **19** | **47** |

## 717: Excision of intervertebral disc (80.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 82 | 7.6 | 17 | 2 | 3 | 7 | 10 | 13 | 14 | 25 |
| 20-34 | 1604 | 8.4 | 20 | 3 | 4 | 7 | 10 | 14 | 17 | 24 |
| 35-49 | 2581 | 8.8 | 20 | 4 | 4 | 8 | 11 | 15 | 17 | 23 |
| 50-64 | 1263 | 9.2 | 22 | 4 | 5 | 8 | 11 | 15 | 18 | 24 |
| 65+ | 305 | 11.3 | 34 | 4 | 6 | 10 | 14 | 19 | 22 | 32 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 29 | 13.3 | 277 | 2 | 4 | 7 | 12 | 29 | 50 | 83 |
| 20-34 | 631 | 11.7 | 69 | 4 | 5 | 9 | 14 | 21 | 28 | 47 |
| 35-49 | 1282 | 11.3 | 54 | 4 | 5 | 9 | 14 | 20 | 25 | 43 |
| 50-64 | 1127 | 11.7 | 59 | 4 | 5 | 10 | 14 | 19 | 24 | 38 |
| 65+ | 646 | 15.6 | 106 | 6 | 7 | 13 | 19 | 28 | 35 | 65 |
| **TOT. SINGLE DX** | 5835 | 8.9 | 22 | 4 | 4 | 8 | 11 | 15 | 18 | 24 |
| **" MULTIPLE DX** | 3715 | 12.3 | 72 | 4 | 5 | 10 | 15 | 22 | 27 | 47 |
| TOTAL 0-19 YRS | 111 | 9.1 | 90 | 2 | 3 | 7 | 10 | 14 | 22 | 62 |
| 20-34 | 2235 | 9.3 | 37 | 4 | 4 | 8 | 11 | 16 | 20 | 35 |
| 35-49 | 3863 | 9.6 | 33 | 4 | 5 | 8 | 11 | 16 | 20 | 32 |
| 50-64 | 2390 | 10.4 | 42 | 4 | 5 | 9 | 12 | 18 | 21 | 33 |
| 65+ | 951 | 14.2 | 87 | 5 | 6 | 12 | 17 | 26 | 32 | 60 |
| **GRAND TOTAL** | **9550** | **10.2** | **44** | **4** | **5** | **8** | **12** | **18** | **22** | **36** |

## 718: Excision of semilunar cartilage of knee joint (80.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 629 | 2.2 | 1 | <1 | <1 | 2 | 3 | 4 | 5 | 7 |
| 20-34 | 1909 | 2.4 | 2 | <1 | <1 | 2 | 3 | 5 | 5 | 8 |
| 35-49 | 1162 | 2.6 | 4 | <1 | <1 | 2 | 3 | 5 | 6 | 11 |
| 50-64 | 685 | 2.8 | 5 | <1 | 1 | 2 | 4 | 5 | 6 | 9 |
| 65+ | 217 | 3.1 | 5 | 1 | 1 | 2 | 4 | 6 | 7 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 578 | 2.9 | 5 | <1 | 1 | 2 | 4 | 5 | 6 | 12 |
| 20-34 | 1984 | 2.8 | 5 | <1 | <1 | 2 | 4 | 5 | 6 | 12 |
| 35-49 | 1199 | 2.8 | 7 | <1 | <1 | 2 | 3 | 5 | 6 | 13 |
| 50-64 | 1235 | 3.0 | 6 | <1 | 1 | 2 | 4 | 6 | 8 | 14 |
| 65+ | 582 | 3.9 | 14 | 1 | 1 | 3 | 5 | 7 | 10 | 21 |
| **TOT. SINGLE DX** | 4602 | 2.5 | 4 | <1 | <1 | 2 | 3 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 5578 | 3.0 | 7 | <1 | 1 | 2 | 4 | 6 | 7 | 13 |
| TOTAL 0-19 YRS | 1207 | 2.5 | 4 | <1 | <1 | 2 | 3 | 5 | 6 | 8 |
| 20-34 | 3893 | 2.6 | 4 | <1 | <1 | 2 | 3 | 5 | 6 | 10 |
| 35-49 | 2361 | 2.7 | 6 | <1 | <1 | 2 | 3 | 5 | 6 | 12 |
| 50-64 | 1920 | 3.0 | 6 | <1 | 1 | 2 | 4 | 5 | 7 | 13 |
| 65+ | 799 | 3.7 | 12 | 1 | 1 | 3 | 4 | 7 | 9 | 19 |
| **GRAND TOTAL** | **10180** | **2.8** | **6** | **<1** | **1** | **2** | **4** | **5** | **6** | **12** |

## 719: Miscellaneous excisions of joint structures (80.2-80.3; 80.7-80.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 915 | 2.0 | 7 | <1 | <1 | 1 | 2 | 3 | 4 | 12 |
| 20-34 | 1658 | 2.0 | 3 | <1 | <1 | 2 | 2 | 4 | 5 | 9 |
| 35-49 | 786 | 2.3 | 4 | <1 | <1 | 2 | 3 | 4 | 5 | 12 |
| 50-64 | 538 | 2.9 | 7 | <1 | 1 | 2 | 3 | 6 | 8 | 16 |
| 65+ | 226 | 3.9 | 34 | <1 | 1 | 2 | 4 | 8 | 12 | 27 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 547 | 2.6 | 14 | <1 | 1 | 2 | 3 | 4 | 6 | 17 |
| 20-34 | 1259 | 2.9 | 13 | <1 | <1 | 2 | 3 | 5 | 8 | 22 |
| 35-49 | 695 | 3.3 | 16 | <1 | <1 | 2 | 4 | 6 | 8 | 25 |
| 50-64 | 592 | 4.4 | 38 | 1 | 1 | 3 | 4 | 9 | 15 | 38 |
| 65+ | 386 | 7.4 | 105 | 1 | 1 | 3 | 8 | 18 | 28 | 54 |
| **TOT. SINGLE DX** | 4123 | 2.3 | 7 | <1 | <1 | 2 | 3 | 4 | 6 | 13 |
| **" MULTIPLE DX** | 3479 | 3.7 | 31 | <1 | 1 | 2 | 4 | 7 | 11 | 30 |
| TOTAL 0-19 YRS | 1462 | 2.2 | 10 | <1 | <1 | 2 | 2 | 4 | 5 | 17 |
| 20-34 | 2917 | 2.4 | 8 | <1 | <1 | 2 | 3 | 4 | 6 | 15 |
| 35-49 | 1481 | 2.8 | 10 | <1 | <1 | 2 | 3 | 5 | 7 | 17 |
| 50-64 | 1130 | 3.7 | 25 | <1 | 1 | 2 | 4 | 7 | 11 | 24 |
| 65+ | 612 | 6.1 | 82 | 1 | 1 | 3 | 6 | 15 | 23 | 50 |
| **GRAND TOTAL** | **7602** | **2.9** | **19** | **<1** | **<1** | **2** | **3** | **5** | **8** | **23** |

## 720: Spinal fusion (81.0)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 149 | 10.1 | 18 | 4 | 5 | 9 | 12 | 16 | 18 | 25 |
| 20-34 | 170 | 9.4 | 22 | 4 | 4 | 8 | 12 | 15 | 18 | 25 |
| 35-49 | 261 | 8.9 | 24 | 4 | 4 | 8 | 11 | 14 | 17 | 25 |
| 50-64 | 131 | 9.1 | 32 | 4 | 4 | 8 | 11 | 17 | 20 | 36 |
| 65+ | 36 | 10.7 | 36 | 5 | 5 | 9 | 14 | 20 | 26 | 27 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 92 | 14.4 | 100 | 5 | 7 | 11 | 16 | 26 | 33 | 63 |
| 20-34 | 219 | 16.0 | 185 | 5 | 6 | 12 | 18 | 30 | 40 | 82 |
| 35-49 | 310 | 12.0 | 86 | 4 | 5 | 9 | 14 | 21 | 28 | 54 |
| 50-64 | 233 | 12.4 | 122 | 4 | 5 | 9 | 14 | 25 | 29 | 90 |
| 65+ | 89 | 19.9 | 288 | 4 | 5 | 15 | 26 | 47 | 66 | 99+ |
| **TOT. SINGLE DX** | 747 | 9.4 | 25 | 4 | 4 | 8 | 11 | 15 | 18 | 26 |
| **" MULTIPLE DX** | 943 | 14.0 | 144 | 4 | 5 | 10 | 16 | 26 | 36 | 78 |
| TOTAL 0-19 YRS | 241 | 11.7 | 54 | 4 | 6 | 10 | 13 | 18 | 25 | 39 |
| 20-34 | 389 | 13.1 | 125 | 4 | 5 | 10 | 15 | 22 | 32 | 74 |
| 35-49 | 571 | 10.6 | 61 | 4 | 5 | 9 | 12 | 18 | 23 | 46 |
| 50-64 | 364 | 11.2 | 92 | 4 | 4 | 9 | 13 | 22 | 28 | 64 |
| 65+ | 125 | 17.3 | 233 | 4 | 5 | 13 | 20 | 44 | 61 | 99+ |
| **GRAND TOTAL** | **1690** | **11.9** | **97** | **4** | **5** | **9** | **14** | **21** | **28** | **67** |

## 721: Arthrodesis and stabilization of joint (81.1-81.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 75 | 3.7 | 6 | 1 | 1 | 3 | 5 | 6 | 8 | 14 |
| 20-34 | 192 | 4.4 | 21 | <1 | 1 | 3 | 5 | 8 | 10 | 27 |
| 35-49 | 197 | 4.2 | 9 | <1 | 1 | 3 | 6 | 9 | 10 | 14 |
| 50-64 | 275 | 4.2 | 19 | <1 | 1 | 3 | 5 | 8 | 10 | 27 |
| 65+ | 140 | 4.2 | 23 | <1 | <1 | 3 | 5 | 7 | 11 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 74 | 4.4 | 5 | 1 | 2 | 4 | 6 | 7 | 10 | 12 |
| 20-34 | 198 | 5.3 | 22 | 1 | 1 | 4 | 6 | 11 | 15 | 25 |
| 35-49 | 193 | 5.3 | 28 | 1 | 1 | 4 | 6 | 9 | 15 | 31 |
| 50-64 | 246 | 6.2 | 61 | 1 | 2 | 4 | 7 | 12 | 17 | 60 |
| 65+ | 208 | 6.9 | 81 | 1 | 2 | 4 | 7 | 12 | 21 | 51 |
| **TOT. SINGLE DX** | 879 | 4.2 | 17 | <1 | 1 | 3 | 5 | 8 | 10 | 23 |
| **" MULTIPLE DX** | 919 | 5.8 | 47 | 1 | 2 | 4 | 7 | 11 | 15 | 36 |
| TOTAL 0-19 YRS | 149 | 4.1 | 6 | 1 | 1 | 3 | 5 | 7 | 9 | 14 |
| 20-34 | 390 | 4.8 | 22 | <1 | 1 | 4 | 6 | 9 | 13 | 25 |
| 35-49 | 390 | 4.8 | 19 | <1 | 1 | 4 | 6 | 9 | 12 | 19 |
| 50-64 | 521 | 5.2 | 40 | 1 | 1 | 4 | 6 | 10 | 13 | 31 |
| 65+ | 348 | 5.8 | 60 | <1 | 1 | 4 | 6 | 11 | 15 | 42 |
| **GRAND TOTAL** | **1798** | **5.0** | **33** | **<1** | **1** | **4** | **6** | **9** | **13** | **31** |

## 722: Total hip replacement (81.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2 | 9.5 | 1 | 9 | 9 | 9 | 10 | 10 | 10 | 10 |
| 20-34 | 37 | 13.1 | 57 | 6 | 7 | 10 | 15 | 27 | 33 | 35 |
| 35-49 | 107 | 11.9 | 19 | 7 | 8 | 11 | 14 | 16 | 19 | 30 |
| 50-64 | 488 | 12.4 | 28 | 7 | 8 | 11 | 14 | 18 | 20 | 32 |
| 65+ | 1048 | 13.7 | 30 | 8 | 8 | 13 | 16 | 20 | 23 | 30 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 4 | 16.3 | 64 | 10 | 10 | 13 | 21 | 28 | 28 | 28 |
| 20-34 | 52 | 14.5 | 46 | 8 | 8 | 13 | 16 | 26 | 32 | 34 |
| 35-49 | 121 | 14.2 | 72 | 7 | 8 | 12 | 17 | 24 | 29 | 83 |
| 50-64 | 613 | 15.3 | 64 | 7 | 9 | 13 | 17 | 25 | 31 | 53 |
| 65+ | 1799 | 17.6 | 95 | 8 | 10 | 15 | 20 | 29 | 37 | 63 |
| **TOT. SINGLE DX** | 1682 | 13.2 | 30 | 7 | 8 | 12 | 15 | 19 | 22 | 32 |
| **" MULTIPLE DX** | 2589 | 16.8 | 87 | 8 | 9 | 14 | 19 | 27 | 35 | 58 |
| TOTAL 0-19 YRS | 6 | 14.0 | 51 | 9 | 9 | 12 | 14 | 27 | 28 | 28 |
| 20-34 | 89 | 13.9 | 51 | 7 | 8 | 12 | 16 | 27 | 32 | 35 |
| 35-49 | 228 | 13.1 | 49 | 7 | 8 | 11 | 15 | 20 | 25 | 48 |
| 50-64 | 1101 | 14.0 | 50 | 7 | 8 | 12 | 16 | 22 | 27 | 42 |
| 65+ | 2847 | 16.2 | 75 | 8 | 9 | 14 | 18 | 25 | 31 | 52 |
| **GRAND TOTAL** | **4271** | **15.4** | **68** | **8** | **9** | **13** | **17** | **24** | **30** | **51** |

## 723: Other arthroplasty of hip (81.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 14 | 8.6 | 68 | <1 | 1 | 6 | 13 | 24 | 26 | 26 |
| 20-34 | 10 | 8.1 | 7 | 3 | 5 | 8 | 9 | 12 | 14 | 14 |
| 35-49 | 19 | 13.4 | 72 | 3 | 7 | 11 | 13 | 29 | 33 | 36 |
| 50-64 | 89 | 11.8 | 20 | 7 | 8 | 11 | 14 | 17 | 20 | 29 |
| 65+ | 440 | 14.0 | 29 | 7 | 8 | 13 | 16 | 20 | 22 | 36 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 6 | 15.8 | 65 | 9 | 9 | 13 | 22 | 28 | 29 | 29 |
| 20-34 | 13 | 11.4 | 38 | 2 | 3 | 13 | 16 | 19 | 21 | 21 |
| 35-49 | 34 | 15.7 | 42 | 7 | 7 | 15 | 18 | 25 | 28 | 32 |
| 50-64 | 216 | 15.4 | 83 | 6 | 8 | 13 | 17 | 24 | 37 | 52 |
| 65+ | 2107 | 17.2 | 96 | 8 | 9 | 15 | 20 | 27 | 34 | 63 |
| **TOT. SINGLE DX** | 572 | 13.4 | 31 | 7 | 8 | 13 | 16 | 19 | 22 | 35 |
| **" MULTIPLE DX** | 2376 | 17.0 | 94 | 8 | 9 | 15 | 20 | 27 | 35 | 62 |
| TOTAL 0-19 YRS | 20 | 10.8 | 75 | <1 | 1 | 9 | 15 | 25 | 27 | 29 |
| 20-34 | 23 | 10.0 | 27 | 2 | 3 | 9 | 14 | 17 | 19 | 21 |
| 35-49 | 53 | 14.9 | 53 | 6 | 7 | 13 | 17 | 27 | 30 | 36 |
| 50-64 | 305 | 14.3 | 67 | 7 | 8 | 12 | 16 | 22 | 34 | 47 |
| 65+ | 2547 | 16.7 | 86 | 8 | 9 | 15 | 19 | 26 | 33 | 60 |
| **GRAND TOTAL** | **2948** | **16.3** | **84** | **7** | **9** | **14** | **19** | **26** | **33** | **59** |

## 724: Arthroplasty of knee (81.41-81.47)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 426 | 3.9 | 4 | 1 | 1 | 4 | 5 | 6 | 8 | 12 |
| 20-34 | 797 | 4.5 | 11 | 1 | 2 | 4 | 5 | 7 | 9 | 14 |
| 35-49 | 264 | 6.1 | 24 | 1 | 2 | 5 | 8 | 13 | 15 | 24 |
| 50-64 | 424 | 11.8 | 39 | 2 | 5 | 11 | 14 | 20 | 24 | 33 |
| 65+ | 842 | 14.0 | 35 | 7 | 8 | 13 | 16 | 21 | 24 | 35 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 602 | 4.7 | 12 | 1 | 2 | 4 | 5 | 7 | 9 | 17 |
| 20-34 | 1208 | 5.0 | 16 | 2 | 2 | 4 | 6 | 8 | 9 | 16 |
| 35-49 | 435 | 5.6 | 22 | 1 | 2 | 4 | 6 | 10 | 14 | 23 |
| 50-64 | 501 | 12.2 | 64 | 2 | 3 | 11 | 15 | 22 | 28 | 40 |
| 65+ | 1099 | 16.6 | 76 | 7 | 9 | 15 | 19 | 27 | 33 | 53 |
| **TOT. SINGLE DX** | 2753 | 8.6 | 43 | 1 | 2 | 7 | 12 | 17 | 20 | 30 |
| **" MULTIPLE DX** | 3845 | 9.3 | 67 | 2 | 3 | 6 | 13 | 19 | 24 | 39 |
| TOTAL 0-19 YRS | 1028 | 4.4 | 9 | 1 | 2 | 4 | 5 | 7 | 8 | 15 |
| 20-34 | 2005 | 4.8 | 15 | 1 | 2 | 4 | 6 | 8 | 9 | 15 |
| 35-49 | 699 | 5.8 | 23 | 1 | 2 | 4 | 7 | 11 | 15 | 23 |
| 50-64 | 925 | 12.1 | 53 | 2 | 4 | 11 | 15 | 21 | 26 | 39 |
| 65+ | 1941 | 15.5 | 60 | 7 | 9 | 14 | 18 | 24 | 30 | 47 |
| **GRAND TOTAL** | **6598** | **9.0** | **57** | **2** | **2** | **6** | **13** | **18** | **22** | **36** |

## 725: Repair of other joint (81.3; 81.48-81.49; 81.7-81.8; 81.93-81.96; 81.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 580 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 5 | 8 |
| 20-34 | 1224 | 3.2 | 2 | 1 | 1 | 3 | 4 | 5 | 6 | 8 |
| 35-49 | 482 | 3.4 | 5 | 1 | 1 | 3 | 4 | 6 | 7 | 11 |
| 50-64 | 449 | 4.0 | 12 | 1 | 1 | 3 | 5 | 7 | 9 | 26 |
| 65+ | 220 | 5.3 | 23 | 1 | 2 | 4 | 6 | 11 | 14 | 24 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 266 | 3.1 | 4 | 1 | 1 | 3 | 4 | 6 | 7 | 10 |
| 20-34 | 602 | 3.7 | 22 | 1 | 1 | 3 | 4 | 6 | 7 | 20 |
| 35-49 | 390 | 3.8 | 7 | 1 | 1 | 3 | 5 | 7 | 8 | 14 |
| 50-64 | 505 | 4.8 | 15 | 1 | 2 | 4 | 5 | 8 | 12 | 21 |
| 65+ | 356 | 7.1 | 45 | 2 | 2 | 5 | 9 | 15 | 20 | 28 |
| **TOT. SINGLE DX** | 2955 | 3.4 | 7 | 1 | 1 | 3 | 4 | 6 | 7 | 13 |
| **" MULTIPLE DX** | 2119 | 4.5 | 21 | 1 | 1 | 3 | 5 | 8 | 12 | 22 |
| TOTAL 0-19 YRS | 846 | 3.0 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 9 |
| 20-34 | 1826 | 3.3 | 9 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 35-49 | 872 | 3.6 | 7 | 1 | 1 | 3 | 4 | 6 | 7 | 13 |
| 50-64 | 954 | 4.4 | 15 | 1 | 2 | 4 | 5 | 8 | 10 | 22 |
| 65+ | 576 | 6.4 | 38 | 2 | 2 | 4 | 8 | 14 | 18 | 28 |
| **GRAND TOTAL** | **5074** | **3.9** | **13** | **1** | **1** | **3** | **4** | **6** | **9** | **18** |

## 726: Incision and division of muscle, tendon, fascia, and bursa (82.0; 83.0-83.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 282 | 3.4 | 15 | <1 | <1 | 2 | 4 | 7 | 11 | 24 |
| 20-34 | 259 | 3.1 | 8 | <1 | <1 | 2 | 4 | 6 | 8 | 13 |
| 35-49 | 250 | 2.7 | 8 | <1 | <1 | 2 | 3 | 5 | 8 | 14 |
| 50-64 | 199 | 2.3 | 6 | <1 | <1 | 2 | 3 | 4 | 6 | 16 |
| 65+ | 82 | 2.8 | 8 | <1 | <1 | 2 | 3 | 6 | 9 | 17 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 195 | 5.6 | 61 | 1 | 1 | 3 | 6 | 11 | 17 | 49 |
| 20-34 | 258 | 6.3 | 69 | <1 | 1 | 4 | 8 | 13 | 18 | 45 |
| 35-49 | 191 | 5.7 | 42 | <1 | 1 | 3 | 8 | 13 | 18 | 32 |
| 50-64 | 186 | 6.4 | 66 | 1 | 1 | 3 | 7 | 17 | 22 | 38 |
| 65+ | 137 | 12.2 | 218 | <1 | 1 | 7 | 15 | 30 | 48 | 73 |
| **TOT. SINGLE DX** | 1072 | 2.9 | 10 | <1 | <1 | 2 | 3 | 6 | 8 | 18 |
| **" MULTIPLE DX** | 967 | 6.9 | 87 | 1 | 1 | 4 | 9 | 15 | 22 | 54 |
| TOTAL 0-19 YRS | 477 | 4.3 | 35 | <1 | <1 | 3 | 5 | 9 | 14 | 29 |
| 20-34 | 517 | 4.7 | 41 | <1 | <1 | 3 | 5 | 10 | 14 | 31 |
| 35-49 | 441 | 4.0 | 25 | <1 | <1 | 2 | 5 | 9 | 14 | 24 |
| 50-64 | 385 | 4.3 | 39 | <1 | <1 | 2 | 4 | 10 | 17 | 34 |
| 65+ | 219 | 8.7 | 161 | <1 | <1 | 4 | 11 | 20 | 34 | 71 |
| **GRAND TOTAL** | **2039** | **4.8** | **51** | **<1** | **<1** | **3** | **5** | **11** | **16** | **36** |

## 727: Excision of lesion of muscle, tendon, fascia, and bursa (82.2; 83.21; 83.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 402 | 2.0 | 10 | <1 | <1 | 1 | 2 | 3 | 5 | 22 |
| 20-34 | 687 | 1.9 | 5 | <1 | <1 | 1 | 2 | 3 | 4 | 11 |
| 35-49 | 478 | 2.0 | 7 | <1 | <1 | 1 | 2 | 4 | 5 | 14 |
| 50-64 | 373 | 2.3 | 7 | <1 | <1 | 2 | 3 | 4 | 5 | 9 |
| 65+ | 151 | 2.8 | 6 | <1 | <1 | 2 | 3 | 6 | 9 | 12 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 65 | 3.9 | 35 | <1 | <1 | 2 | 4 | 9 | 14 | 38 |
| 20-34 | 179 | 3.4 | 36 | <1 | <1 | 2 | 4 | 7 | 9 | 60 |
| 35-49 | 207 | 4.7 | 50 | <1 | 1 | 3 | 4 | 10 | 17 | 35 |
| 50-64 | 293 | 6.3 | 101 | <1 | 1 | 3 | 6 | 13 | 23 | 49 |
| 65+ | 228 | 8.6 | 94 | 1 | 1 | 5 | 11 | 23 | 29 | 57 |
| **TOT. SINGLE DX** | 2091 | 2.1 | 7 | <1 | <1 | 1 | 2 | 4 | 5 | 12 |
| **" MULTIPLE DX** | 972 | 5.8 | 76 | <1 | 1 | 3 | 6 | 13 | 23 | 49 |
| TOTAL 0-19 YRS | 467 | 2.3 | 14 | <1 | <1 | 1 | 2 | 4 | 6 | 23 |
| 20-34 | 866 | 2.2 | 12 | <1 | <1 | 1 | 2 | 4 | 6 | 12 |
| 35-49 | 685 | 2.9 | 22 | <1 | <1 | 2 | 3 | 5 | 8 | 26 |
| 50-64 | 666 | 4.0 | 53 | <1 | <1 | 2 | 4 | 8 | 12 | 38 |
| 65+ | 379 | 6.3 | 67 | <1 | 1 | 3 | 7 | 15 | 25 | 47 |
| **GRAND TOTAL** | **3063** | **3.3** | **32** | **<1** | **<1** | **2** | **3** | **6** | **11** | **29** |

## 728: Division and resection of muscle, tendon, and fascia of hand (82.1; 82.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 22 | 1.5 | | <1 | <1 | 1 | 2 | 3 | 3 | 4 |
| 20-34 | 49 | 2.5 | 9 | <1 | <1 | 2 | 2 | 5 | 9 | 19 |
| 35-49 | 85 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 4 | 4 |
| 50-64 | 285 | 1.9 | | <1 | <1 | 2 | 2 | 3 | 3 | 4 |
| 65+ | 217 | 2.1 | 1 | <1 | <1 | 2 | 3 | 3 | 4 | 7 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 20 | 7.9 | 104 | <1 | 1 | 3 | 10 | 21 | 32 | 40 |
| 20-34 | 38 | 3.9 | 18 | <1 | <1 | 2 | 5 | 9 | 14 | 20 |
| 35-49 | 48 | 2.3 | 2 | <1 | <1 | 2 | 3 | 4 | 5 | 7 |
| 50-64 | 145 | 2.8 | 8 | <1 | 1 | 2 | 3 | 5 | 8 | 20 |
| 65+ | 107 | 3.1 | 12 | 1 | 1 | 2 | 3 | 5 | 9 | 21 |
| **TOT. SINGLE DX** | 658 | 2.0 | 2 | <1 | <1 | 2 | 2 | 3 | 4 | 6 |
| **" MULTIPLE DX** | 358 | 3.2 | 16 | <1 | 1 | 2 | 3 | 6 | 10 | 23 |
| TOTAL 0-19 YRS | 42 | 4.5 | 59 | <1 | <1 | 2 | 3 | 13 | 21 | 40 |
| 20-34 | 87 | 3.1 | 13 | <1 | <1 | 2 | 3 | 6 | 11 | 20 |
| 35-49 | 133 | 2.0 | 1 | <1 | <1 | 2 | 2 | 3 | 4 | 7 |
| 50-64 | 430 | 2.2 | 4 | <1 | <1 | 2 | 2 | 3 | 4 | 11 |
| 65+ | 324 | 2.4 | 6 | <1 | <1 | 2 | 3 | 4 | 5 | 13 |
| **GRAND TOTAL** | **1016** | **2.4** | **7** | **<1** | **<1** | **2** | **3** | **4** | **5** | **16** |

### 729: Repair and plastic operations on hand (82.4-82.8; 82.91; 82.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 211 | 1.8 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 15 |
| 20-34 | 331 | 2.0 | 5 | <1 | <1 | 1 | 2 | 3 | 5 | 12 |
| 35-49 | 143 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 8 |
| 50-64 | 75 | 1.9 | 2 | <1 | <1 | 2 | 2 | 3 | 4 | 10 |
| 65+ | 40 | 2.0 | 2 | <1 | <1 | 2 | 2 | 3 | 6 | 7 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 342 | 2.0 | 3 | <1 | 1 | 1 | 2 | 3 | 5 | 15 |
| 20-34 | 598 | 2.4 | 9 | <1 | 1 | 2 | 3 | 4 | 6 | 14 |
| 35-49 | 220 | 2.5 | 10 | <1 | <1 | 2 | 3 | 5 | 6 | 19 |
| 50-64 | 161 | 3.0 | 12 | <1 | 1 | 2 | 3 | 6 | 10 | 18 |
| 65+ | 52 | 2.7 | 12 | <1 | 1 | 2 | 3 | 4 | 7 | 25 |
| **TOT. SINGLE DX** | 800 | 1.9 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 10 |
| **" MULTIPLE DX** | 1373 | 2.4 | 9 | <1 | 1 | 2 | 3 | 4 | 6 | 16 |
| TOTAL 0-19 YRS | 553 | 1.9 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 15 |
| 20-34 | 929 | 2.2 | 8 | <1 | <1 | 2 | 3 | 4 | 5 | 13 |
| 35-49 | 363 | 2.2 | 7 | <1 | <1 | 2 | 2 | 4 | 5 | 15 |
| 50-64 | 236 | 2.7 | 10 | <1 | 1 | 2 | 3 | 5 | 8 | 17 |
| 65+ | 92 | 2.4 | 8 | <1 | <1 | 2 | 3 | 4 | 7 | 18 |
| **GRAND TOTAL** | **2173** | **2.2** | **7** | **<1** | **<1** | **2** | **2** | **4** | **5** | **15** |

### 730: Resection and suture of muscle, tendon, fascia, except hand (83.4-83.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 116 | 2.2 | 4 | <1 | <1 | 2 | 3 | 4 | 6 | 13 |
| 20-34 | 248 | 2.7 | 6 | <1 | <1 | 2 | 3 | 6 | 7 | 14 |
| 35-49 | 245 | 3.1 | 3 | <1 | <1 | 3 | 4 | 5 | 6 | 11 |
| 50-64 | 355 | 3.7 | 6 | <1 | 1 | 3 | 5 | 7 | 9 | 13 |
| 65+ | 122 | 5.0 | 30 | 1 | 2 | 4 | 6 | 8 | 12 | 29 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 165 | 3.3 | 27 | <1 | 1 | 2 | 4 | 6 | 8 | 24 |
| 20-34 | 344 | 3.8 | 23 | 1 | 1 | 2 | 4 | 8 | 11 | 33 |
| 35-49 | 249 | 4.5 | 19 | 1 | 1 | 3 | 5 | 9 | 12 | 21 |
| 50-64 | 351 | 5.3 | 32 | 1 | 2 | 4 | 6 | 9 | 15 | 27 |
| 65+ | 226 | 6.9 | 50 | 1 | 2 | 5 | 8 | 13 | 18 | 36 |
| **TOT. SINGLE DX** | 1086 | 3.3 | 9 | <1 | 1 | 3 | 4 | 6 | 8 | 14 |
| **" MULTIPLE DX** | 1335 | 4.8 | 32 | 1 | 1 | 3 | 5 | 9 | 13 | 29 |
| TOTAL 0-19 YRS | 281 | 2.9 | 18 | <1 | 1 | 2 | 3 | 5 | 7 | 21 |
| 20-34 | 592 | 3.4 | 17 | <1 | 1 | 2 | 4 | 6 | 10 | 22 |
| 35-49 | 494 | 3.8 | 12 | <1 | 1 | 3 | 5 | 7 | 11 | 16 |
| 50-64 | 706 | 4.5 | 20 | 1 | 1 | 4 | 5 | 8 | 11 | 23 |
| 65+ | 348 | 6.2 | 44 | 1 | 2 | 5 | 7 | 12 | 15 | 33 |
| **GRAND TOTAL** | **2421** | **4.1** | **22** | **<1** | **1** | **3** | **5** | **8** | **11** | **23** |

### 731: Miscellaneous operations on muscle, tendon, fascia, and bursa, except hand (83.71-83.93; 83.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 361 | 3.0 | 3 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 20-34 | 327 | 3.6 | 8 | 1 | 1 | 3 | 4 | 6 | 8 | 14 |
| 35-49 | 184 | 3.6 | 14 | 1 | 1 | 3 | 4 | 6 | 8 | 21 |
| 50-64 | 132 | 3.6 | 11 | 1 | 1 | 3 | 4 | 6 | 8 | 20 |
| 65+ | 48 | 5.7 | 39 | 1 | 1 | 4 | 6 | 10 | 22 | 30 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 321 | 3.7 | 27 | 1 | 1 | 3 | 4 | 6 | 7 | 16 |
| 20-34 | 314 | 4.5 | 42 | 1 | 1 | 3 | 5 | 8 | 12 | 33 |
| 35-49 | 172 | 5.1 | 106 | 1 | 1 | 3 | 5 | 7 | 11 | 76 |
| 50-64 | 129 | 5.1 | 42 | 1 | 1 | 3 | 5 | 9 | 15 | 41 |
| 65+ | 59 | 8.2 | 120 | 2 | 2 | 5 | 9 | 16 | 21 | 76 |
| **TOT. SINGLE DX** | 1052 | 3.4 | 10 | 1 | 1 | 3 | 4 | 6 | 7 | 18 |
| **" MULTIPLE DX** | 995 | 4.6 | 54 | 1 | 1 | 3 | 5 | 7 | 12 | 42 |
| TOTAL 0-19 YRS | 682 | 3.3 | 15 | 1 | 1 | 3 | 4 | 5 | 7 | 12 |
| 20-34 | 641 | 4.0 | 25 | 1 | 1 | 3 | 5 | 7 | 9 | 22 |
| 35-49 | 356 | 4.3 | 59 | 1 | 1 | 3 | 4 | 7 | 9 | 36 |
| 50-64 | 261 | 4.3 | 27 | 1 | 1 | 3 | 5 | 7 | 12 | 29 |
| 65+ | 107 | 7.1 | 85 | 1 | 2 | 5 | 7 | 15 | 22 | 52 |
| **GRAND TOTAL** | **2047** | **4.0** | **32** | **1** | **1** | **3** | **4** | **7** | **9** | **24** |

### 732: Amputation of fingers and thumb (84.01-84.02)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 40 | 1.6 | | <1 | <1 | 1 | 2 | 3 | 3 | 4 |
| 20-34 | 67 | 2.5 | 7 | <1 | 1 | 1 | 3 | 5 | 9 | 14 |
| 35-49 | 46 | 2.6 | 16 | <1 | <1 | 2 | 2 | 4 | 7 | 27 |
| 50-64 | 23 | 2.1 | 1 | <1 | 1 | 2 | 3 | 3 | 4 | 4 |
| 65+ | 21 | 1.8 | | <1 | <1 | 2 | 2 | 3 | 4 | 4 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 63 | 2.8 | 8 | <1 | 1 | 1 | 3 | 7 | 10 | 14 |
| 20-34 | 167 | 4.3 | 92 | <1 | 1 | 2 | 3 | 8 | 13 | 88 |
| 35-49 | 95 | 3.7 | 23 | 1 | 1 | 2 | 4 | 9 | 13 | 28 |
| 50-64 | 85 | 5.0 | 62 | <1 | 1 | 2 | 4 | 11 | 27 | 36 |
| 65+ | 54 | 6.6 | 137 | 1 | 1 | 2 | 4 | 17 | 32 | 60 |
| **TOT. SINGLE DX** | 197 | 2.2 | 7 | <1 | <1 | 2 | 3 | 4 | 5 | 14 |
| **" MULTIPLE DX** | 464 | 4.4 | 67 | <1 | 1 | 2 | 4 | 9 | 16 | 51 |
| TOTAL 0-19 YRS | 103 | 2.4 | 6 | <1 | <1 | 1 | 3 | 5 | 8 | 12 |
| 20-34 | 234 | 3.8 | 68 | <1 | 1 | 2 | 3 | 7 | 13 | 64 |
| 35-49 | 141 | 3.4 | 21 | <1 | 1 | 2 | 3 | 8 | 12 | 27 |
| 50-64 | 108 | 4.4 | 51 | <1 | 1 | 2 | 4 | 9 | 26 | 35 |
| 65+ | 75 | 5.2 | 103 | 1 | 1 | 2 | 3 | 11 | 21 | 58 |
| **GRAND TOTAL** | **661** | **3.8** | **50** | **<1** | **1** | **2** | **3** | **8** | **12** | **37** |

### 733: Amputation of hand, forearm, and upper arm (84.00; 84.03-84.09)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 2.0 | 1 | 1 | 1 | 2 | 3 | 3 | 3 | 3 |
| 20-34 | 7 | 4.1 | 13 | 1 | 1 | 3 | 8 | 9 | 9 | 9 |
| 35-49 | 10 | 2.9 | 2 | <1 | 1 | 3 | 3 | 5 | 7 | 7 |
| 50-64 | 6 | 3.7 | 4 | 1 | 1 | 4 | 4 | 7 | 7 | 7 |
| 65+ | 2 | 24.0 | 288 | 12 | 12 | 24 | 36 | 36 | 36 | 36 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 7 | 8.0 | 34 | 2 | 2 | 8 | 13 | 16 | 17 | 17 |
| 20-34 | 19 | 9.5 | 248 | 1 | 1 | 3 | 10 | 28 | 50 | 65 |
| 35-49 | 3 | 30.7 | 632 | 2 | 2 | 41 | 47 | 49 | 49 | 49 |
| 50-64 | 13 | 21.6 | 236 | 4 | 6 | 20 | 39 | 54 | 90 | 99+ |
| 65+ | 13 | 8.8 | 41 | 1 | 1 | 10 | 12 | 19 | 21 | 22 |
| **TOT. SINGLE DX** | 28 | 4.8 | 45 | 1 | 1 | 3 | 5 | 9 | 14 | 36 |
| **" MULTIPLE DX** | 55 | 13.2 | 219 | 1 | 1 | 8 | 18 | 40 | 49 | 98 |
| TOTAL 0-19 YRS | 10 | 6.2 | 31 | 1 | 1 | 3 | 9 | 15 | 17 | 17 |
| 20-34 | 26 | 8.1 | 188 | 1 | 1 | 3 | 9 | 23 | 38 | 65 |
| 35-49 | 13 | 9.3 | 256 | <1 | 1 | 3 | 5 | 43 | 48 | 49 |
| 50-64 | 19 | 15.9 | 232 | 1 | 2 | 9 | 32 | 44 | 74 | 99+ |
| 65+ | 15 | 10.8 | 85 | 1 | 1 | 10 | 13 | 22 | 32 | 36 |
| **GRAND TOTAL** | **83** | **10.3** | **175** | **1** | **1** | **5** | **12** | **35** | **43** | **88** |

### 734: Amputation of toes (84.11)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 16 | 3.2 | 9 | <1 | <1 | 2 | 4 | 8 | 11 | 12 |
| 20-34 | 12 | 2.5 | 4 | <1 | <1 | 2 | 3 | 6 | 7 | 7 |
| 35-49 | 15 | 3.9 | 9 | 1 | 1 | 3 | 5 | 7 | 11 | 13 |
| 50-64 | 22 | 5.7 | 33 | <1 | 1 | 3 | 8 | 15 | 20 | 22 |
| 65+ | 50 | 8.1 | 111 | 1 | 1 | 4 | 10 | 20 | 27 | 62 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 26 | 7.2 | 297 | <1 | 1 | 3 | 5 | 12 | 28 | 90 |
| 20-34 | 42 | 11.2 | 204 | 1 | 2 | 8 | 13 | 20 | 46 | 72 |
| 35-49 | 84 | 13.5 | 120 | 1 | 2 | 11 | 18 | 28 | 34 | 54 |
| 50-64 | 198 | 17.2 | 225 | 2 | 3 | 13 | 22 | 36 | 46 | 74 |
| 65+ | 296 | 16.9 | 249 | 2 | 3 | 12 | 24 | 38 | 50 | 73 |
| **TOT. SINGLE DX** | 115 | 5.8 | 62 | <1 | 1 | 3 | 6 | 13 | 20 | 44 |
| **" MULTIPLE DX** | 646 | 15.8 | 229 | 1 | 2 | 11 | 21 | 35 | 49 | 73 |
| TOTAL 0-19 YRS | 42 | 5.6 | 189 | <1 | 1 | 2 | 5 | 11 | 12 | 90 |
| 20-34 | 54 | 9.3 | 173 | <1 | 1 | 6 | 10 | 16 | 34 | 72 |
| 35-49 | 99 | 12.1 | 115 | 1 | 2 | 9 | 16 | 27 | 34 | 53 |
| 50-64 | 220 | 16.0 | 218 | 2 | 2 | 12 | 21 | 35 | 45 | 74 |
| 65+ | 346 | 15.6 | 239 | 1 | 2 | 10 | 22 | 36 | 49 | 72 |
| **GRAND TOTAL** | **761** | **14.3** | **217** | **1** | **2** | **9** | **19** | **33** | **45** | **72** |

### 735: Below-knee amputation (84.15)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 8.0 | | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 20-34 | 8 | 11.0 | 40 | 5 | 6 | 10 | 11 | 22 | 26 | 26 |
| 35-49 | 2 | 11.5 | 25 | 8 | 8 | 11 | 15 | 15 | 15 | 15 |
| 50-64 | 13 | 14.7 | 136 | 4 | 5 | 9 | 20 | 34 | 40 | 41 |
| 65+ | 20 | 13.6 | 85 | 3 | 5 | 12 | 17 | 35 | 67 | 99+ |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 27 | 21.3 | 339 | 4 | 5 | 14 | 39 | 57 | 64 | 99+ |
| 35-49 | 45 | 26.2 | 328 | 7 | 9 | 22 | 40 | 60 | 88 | 99+ |
| 50-64 | 188 | 26.6 | 345 | 7 | 8 | 21 | 35 | 58 | 73 | 99+ |
| 65+ | 452 | 27.9 | 367 | 8 | 10 | 22 | 36 | 56 | 75 | 95 |
| **TOT. SINGLE DX** | 44 | 13.2 | 85 | 4 | 5 | 10 | 16 | 31 | 40 | 99+ |
| **" MULTIPLE DX** | 712 | 27.2 | 359 | 7 | 9 | 22 | 36 | 56 | 75 | 99+ |
| TOTAL 0-19 YRS | 1 | 8.0 | | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| 20-34 | 35 | 18.9 | 287 | 5 | 5 | 11 | 29 | 53 | 60 | 99+ |
| 35-49 | 47 | 23.6 | 324 | 7 | 8 | 22 | 38 | 58 | 87 | 99+ |
| 50-64 | 201 | 25.9 | 340 | 6 | 8 | 21 | 34 | 56 | 71 | 99+ |
| 65+ | 472 | 27.3 | 364 | 7 | 9 | 22 | 35 | 55 | 75 | 97 |
| **GRAND TOTAL** | **756** | **26.4** | **353** | **7** | **8** | **21** | **35** | **56** | **74** | **99+** |

### 736: Above-knee amputation (84.16-84.17)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 9.0 | 61 | 4 | 4 | 5 | 15 | 18 | 18 | 18 |
| 20-34 | 3 | 7.3 | 8 | 4 | 4 | 9 | 9 | 9 | 9 | 9 |
| 35-49 | 3 | 11.3 | 56 | 7 | 7 | 7 | 17 | 20 | 20 | 20 |
| 50-64 | 6 | 12.7 | 21 | 7 | 7 | 12 | 17 | 18 | 18 | 18 |
| 65+ | 22 | 19.8 | 188 | 6 | 7 | 14 | 29 | 44 | 49 | 50 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 35.6 | 739 | 6 | 6 | 48 | 53 | 67 | 67 | 67 |
| 20-34 | 12 | 20.2 | 182 | 3 | 4 | 16 | 33 | 38 | 41 | 41 |
| 35-49 | 18 | 30.1 | 329 | 8 | 12 | 26 | 51 | 64 | 86 | 99+ |
| 50-64 | 64 | 23.8 | 288 | 8 | 10 | 21 | 32 | 61 | 92 | 99+ |
| 65+ | 403 | 23.6 | 342 | 6 | 7 | 18 | 29 | 50 | 68 | 98 |
| **TOT. SINGLE DX** | 37 | 16.1 | 142 | 4 | 5 | 12 | 19 | 33 | 47 | 50 |
| **" MULTIPLE DX** | 502 | 23.9 | 335 | 6 | 8 | 18 | 30 | 51 | 70 | 99+ |
| TOTAL 0-19 YRS | 8 | 25.6 | 629 | 4 | 4 | 13 | 48 | 62 | 67 | 67 |
| 20-34 | 15 | 17.6 | 173 | 3 | 4 | 12 | 31 | 37 | 40 | 41 |
| 35-49 | 21 | 27.4 | 331 | 6 | 7 | 23 | 50 | 59 | 82 | 99+ |
| 50-64 | 70 | 22.9 | 275 | 7 | 9 | 17 | 31 | 55 | 88 | 99+ |
| 65+ | 425 | 23.4 | 335 | 6 | 7 | 18 | 29 | 50 | 67 | 98 |
| **GRAND TOTAL** | **539** | **23.4** | **326** | **6** | **7** | **18** | **30** | **50** | **68** | **99+** |

## 737: Other amputation of lower extremity (84.10; 84.12-84.14; 84.18-84.19)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 8.0 | 109 | 1 | 1 | 3 | 16 | 20 | 20 | 20 |
| 20-34 | 6 | 15.2 | 244 | 2 | 2 | 10 | 18 | 42 | 45 | 45 |
| 35-49 | 6 | 7.7 | 41 | <1 | <1 | 6 | 12 | 17 | 18 | 18 |
| 50-64 | 5 | 7.2 | 48 | 2 | 2 | 6 | 10 | 19 | 19 | 19 |
| 65+ | 7 | 10.6 | 23 | 2 | 3 | 11 | 12 | 17 | 18 | 18 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 5 | 16.4 | 201 | 1 | 1 | 10 | 31 | 32 | 32 | 32 |
| 20-34 | 16 | 24.1 | 208 | 4 | 6 | 22 | 36 | 46 | 49 | 51 |
| 35-49 | 33 | 20.2 | 270 | 4 | 6 | 15 | 26 | 44 | 63 | 68 |
| 50-64 | 76 | 20.9 | 208 | 5 | 6 | 17 | 29 | 39 | 48 | 73 |
| 65+ | 107 | 24.5 | 254 | 6 | 9 | 21 | 31 | 51 | 58 | 88 |
| **TOT. SINGLE DX** | 27 | 10.0 | 86 | 1 | 2 | 8 | 12 | 19 | 24 | 45 |
| **" MULTIPLE DX** | 237 | 22.6 | 238 | 5 | 7 | 19 | 30 | 44 | 55 | 78 |
| TOTAL 0-19 YRS | 8 | 13.3 | 165 | 1 | 1 | 9 | 25 | 32 | 32 | 32 |
| 20-34 | 22 | 21.7 | 224 | 3 | 4 | 18 | 31 | 45 | 48 | 51 |
| 35-49 | 39 | 18.3 | 254 | 3 | 4 | 13 | 23 | 43 | 57 | 68 |
| 50-64 | 81 | 20.1 | 209 | 4 | 6 | 17 | 29 | 38 | 48 | 72 |
| 65+ | 114 | 23.6 | 251 | 5 | 8 | 21 | 30 | 48 | 57 | 86 |
| **GRAND TOTAL** | **264** | **21.3** | **237** | **4** | **5** | **18** | **29** | **42** | **53** | **78** |

## 738: Reattachment of extremity (84.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 28 | 6.6 | 21 | 1 | 1 | 7 | 10 | 13 | 14 | 14 |
| 20-34 | 20 | 8.0 | 18 | 1 | 2 | 8 | 10 | 14 | 16 | 18 |
| 35-49 | 9 | 9.1 | 20 | 2 | 2 | 11 | 13 | 14 | 14 | 14 |
| 50-64 | 2 | 11.5 | 5 | 10 | 10 | 11 | 13 | 13 | 13 | 13 |
| 65+ | 4 | 7.0 | 31 | 2 | 2 | 5 | 10 | 15 | 15 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 29 | 7.5 | 40 | 1 | 1 | 6 | 11 | 17 | 18 | 25 |
| 20-34 | 30 | 12.1 | 81 | 1 | 2 | 10 | 17 | 23 | 31 | 40 |
| 35-49 | 17 | 7.2 | 42 | 1 | 1 | 7 | 10 | 11 | 22 | 28 |
| 50-64 | 10 | 9.2 | 56 | 1 | 1 | 7 | 14 | 20 | 26 | 26 |
| 65+ | 5 | 5.8 | 4 | 3 | 3 | 6 | 7 | 9 | 9 | 9 |
| **TOT. SINGLE DX** | 63 | 7.6 | 21 | 1 | 1 | 8 | 11 | 14 | 14 | 18 |
| **" MULTIPLE DX** | 91 | 9.1 | 57 | 1 | 1 | 7 | 12 | 19 | 25 | 36 |
| TOTAL 0-19 YRS | 57 | 7.1 | 31 | 1 | 1 | 6 | 11 | 14 | 17 | 25 |
| 20-34 | 50 | 10.4 | 60 | 1 | 2 | 9 | 13 | 20 | 23 | 40 |
| 35-49 | 26 | 7.9 | 35 | 1 | 1 | 8 | 11 | 13 | 17 | 28 |
| 50-64 | 12 | 9.6 | 47 | 1 | 2 | 8 | 13 | 18 | 25 | 26 |
| 65+ | 9 | 6.3 | 15 | 2 | 2 | 6 | 7 | 13 | 15 | 15 |
| **GRAND TOTAL** | **154** | **8.4** | **42** | **1** | **1** | **7** | **11** | **16** | **21** | **31** |

## 739: Revision of amputation stump (84.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 15 | 2.1 | 1 | 1 | 1 | 2 | 3 | 4 | 4 | 4 |
| 20-34 | 45 | 3.4 | 31 | <1 | <1 | 2 | 4 | 6 | 7 | 38 |
| 35-49 | 30 | 2.3 | 1 | <1 | <1 | 2 | 3 | 4 | 5 | 5 |
| 50-64 | 27 | 7.4 | 134 | <1 | <1 | 3 | 7 | 17 | 42 | 48 |
| 65+ | 34 | 7.3 | 56 | 1 | 1 | 5 | 10 | 21 | 25 | 25 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 21 | 2.3 | 4 | 1 | 1 | 1 | 3 | 5 | 7 | 9 |
| 20-34 | 58 | 5.2 | 73 | <1 | <1 | 2 | 4 | 13 | 25 | 50 |
| 35-49 | 42 | 9.5 | 218 | <1 | 1 | 4 | 9 | 23 | 50 | 70 |
| 50-64 | 59 | 11.3 | 132 | <1 | 1 | 7 | 16 | 29 | 41 | 95 |
| 65+ | 88 | 14.4 | 159 | 2 | 3 | 10 | 20 | 33 | 55 | 99+ |
| **TOT. SINGLE DX** | 151 | 4.7 | 51 | <1 | <1 | 2 | 5 | 9 | 19 | 41 |
| **" MULTIPLE DX** | 268 | 10.0 | 147 | 1 | 1 | 5 | 14 | 27 | 42 | 93 |
| TOTAL 0-19 YRS | 36 | 2.2 | 3 | 1 | 1 | 2 | 3 | 4 | 5 | 9 |
| 20-34 | 103 | 4.4 | 56 | <1 | <1 | 2 | 4 | 7 | 18 | 45 |
| 35-49 | 72 | 6.5 | 140 | <1 | <1 | 3 | 5 | 14 | 28 | 67 |
| 50-64 | 86 | 10.1 | 135 | <1 | 1 | 6 | 14 | 26 | 42 | 82 |
| 65+ | 122 | 12.5 | 140 | 1 | 2 | 8 | 19 | 30 | 42 | 99+ |
| **GRAND TOTAL** | **419** | **8.1** | **119** | **<1** | **1** | **4** | **9** | **23** | **34** | **61** |

## 740: Miscellaneous operations on musculoskeletal system (84.40; 84.44; 84.48; 84.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 1 | 4.0 | | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 20-34 | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 35-49 | 3 | 7.3 | 8 | 4 | 4 | 9 | 9 | 9 | 9 | 9 |
| 50-64 | 6 | 7.3 | 37 | 1 | 1 | 5 | 12 | 16 | 17 | 17 |
| 65+ | 3 | 4.7 | | 4 | 4 | 5 | 5 | 5 | 5 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 2 | 10.5 | 113 | 3 | 3 | 10 | 18 | 18 | 18 | 18 |
| 35-49 | 1 | 9.0 | | 9 | 9 | 9 | 9 | 9 | 9 | 9 |
| 50-64 | 3 | 16.0 | 243 | 7 | 7 | 7 | 27 | 34 | 34 | 34 |
| 65+ | 1 | 7.0 | | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| **TOT. SINGLE DX** | 14 | 6.4 | 17 | 1 | 3 | 5 | 9 | 12 | 16 | 17 |
| **" MULTIPLE DX** | 7 | 12.1 | 114 | 3 | 4 | 8 | 26 | 80 | 99+ | 99+ |
| TOTAL 0-19 YRS | 1 | 4.0 | | 4 | 4 | 52 | 99+ | 99+ | 99+ | 99+ |
| 20-34 | 3 | 8.7 | 66 | 3 | 3 | 5 | 15 | 18 | 18 | 18 |
| 35-49 | 4 | 7.8 | 6 | 4 | 4 | 9 | 9 | 9 | 9 | 9 |
| 50-64 | 9 | 10.2 | 103 | 1 | 2 | 7 | 13 | 27 | 34 | 34 |
| 65+ | 4 | 5.3 | 2 | 4 | 4 | 5 | 6 | 7 | 7 | 7 |
| **GRAND TOTAL** | **21** | **8.3** | **53** | **2** | **3** | **7** | **9** | **23** | **60** | **99+** |

## 741: Other diagnostic and therapeutic procedures on musculoskeletal structure (for *ICD-9-CM* codes, see Appendix)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| TOT. SINGLE DX | | | | | | | | | | |
| " MULTIPLE DX | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| GRAND TOTAL | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 742: Local exision or destruction of lesion of breast (85.11-85.12; 85.20-85.23)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 174 | 1.4 | | <1 | <1 | 1 | 2 | 2 | 3 | 5 |
| 20-34 | 760 | 1.6 | 3 | <1 | <1 | 1 | 2 | 2 | 3 | 7 |
| 35-49 | 1447 | 1.8 | 2 | <1 | <1 | 1 | 2 | 3 | 3 | 8 |
| 50-64 | 1007 | 2.1 | 3 | <1 | <1 | 2 | 2 | 3 | 5 | 9 |
| 65+ | 636 | 2.5 | 11 | <1 | <1 | 2 | 3 | 4 | 7 | 13 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 39 | 2.1 | 3 | <1 | <1 | 1 | 2 | 5 | 7 | 8 |
| 20-34 | 316 | 2.3 | 6 | <1 | <1 | 2 | 2 | 4 | 6 | 13 |
| 35-49 | 765 | 2.7 | 7 | <1 | <1 | 2 | 3 | 6 | 8 | 14 |
| 50-64 | 766 | 3.6 | 25 | <1 | 1 | 2 | 4 | 8 | 10 | 22 |
| 65+ | 707 | 5.3 | 44 | 1 | 1 | 3 | 6 | 11 | 17 | 37 |
| TOT. SINGLE DX | 4024 | 1.9 | 4 | <1 | <1 | 2 | 2 | 3 | 4 | 9 |
| " MULTIPLE DX | 2593 | 3.6 | 24 | <1 | <1 | 2 | 4 | 8 | 11 | 26 |
| TOTAL 0-19 YRS | 213 | 1.6 | 1 | <1 | <1 | 1 | 2 | 2 | 4 | 6 |
| 20-34 | 1076 | 1.8 | 4 | <1 | <1 | 1 | 2 | 3 | 4 | 10 |
| 35-49 | 2212 | 2.1 | 5 | <1 | <1 | 2 | 2 | 3 | 6 | 11 |
| 50-64 | 1773 | 2.7 | 13 | <1 | <1 | 2 | 3 | 6 | 8 | 15 |
| 65+ | 1343 | 4.0 | 31 | <1 | 1 | 2 | 4 | 8 | 13 | 29 |
| GRAND TOTAL | 6617 | 2.6 | 13 | <1 | <1 | 2 | 2 | 5 | 8 | 16 |

## 743: Simple mastectomy (85.41-85.42)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 26 | 2.1 | 2 | <1 | <1 | 2 | 2 | 4 | 5 | 9 |
| 20-34 | 34 | 2.6 | 4 | <1 | 1 | 2 | 3 | 5 | 8 | 10 |
| 35-49 | 53 | 3.9 | 5 | 1 | 1 | 3 | 6 | 7 | 9 | 10 |
| 50-64 | 64 | 4.5 | 7 | 1 | 2 | 4 | 6 | 8 | 9 | 16 |
| 65+ | 104 | 5.0 | 10 | 1 | 1 | 5 | 6 | 9 | 11 | 17 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 5 | 3.2 | 8 | <1 | <1 | 3 | 4 | 8 | 8 | 8 |
| 20-34 | 12 | 4.0 | 4 | 1 | 1 | 4 | 5 | 7 | 9 | 9 |
| 35-49 | 71 | 5.5 | 20 | 1 | 2 | 4 | 6 | 10 | 14 | 26 |
| 50-64 | 129 | 5.0 | 7 | 2 | 2 | 5 | 6 | 9 | 11 | 15 |
| 65+ | 191 | 8.0 | 62 | 2 | 3 | 6 | 8 | 15 | 21 | 50 |
| TOT. SINGLE DX | 281 | 4.1 | 9 | <1 | 1 | 4 | 6 | 8 | 9 | 17 |
| " MULTIPLE DX | 408 | 6.5 | 38 | 2 | 2 | 5 | 7 | 11 | 16 | 38 |
| TOTAL 0-19 YRS | 31 | 2.3 | 4 | <1 | <1 | 2 | 3 | 4 | 8 | 9 |
| 20-34 | 46 | 3.0 | 5 | <1 | 1 | 2 | 4 | 6 | 8 | 10 |
| 35-49 | 124 | 4.8 | 15 | 1 | 2 | 4 | 6 | 9 | 11 | 22 |
| 50-64 | 193 | 4.8 | 8 | 2 | 2 | 4 | 6 | 8 | 10 | 16 |
| 65+ | 295 | 7.0 | 46 | 2 | 2 | 5 | 8 | 12 | 17 | 43 |
| GRAND TOTAL | 689 | 5.5 | 27 | 1 | 2 | 5 | 6 | 10 | 13 | 26 |

## 744: Extended simple mastectomy and radical mastectomy (85.43-85.48)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 6 | 2.7 | 1 | 1 | 1 | 3 | 3 | 4 | 4 | 4 |
| 20-34 | 58 | 5.5 | 5 | 2 | 3 | 5 | 7 | 8 | 11 | 13 |
| 35-49 | 388 | 6.0 | 5 | 3 | 4 | 6 | 7 | 8 | 10 | 14 |
| 50-64 | 613 | 6.2 | 6 | 3 | 4 | 6 | 8 | 9 | 11 | 14 |
| 65+ | 634 | 6.9 | 8 | 3 | 4 | 6 | 8 | 10 | 12 | 16 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 1 | 5.0 | | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| 20-34 | 67 | 6.5 | 6 | 4 | 4 | 6 | 7 | 10 | 13 | 16 |
| 35-49 | 458 | 6.6 | 16 | 3 | 4 | 6 | 8 | 10 | 12 | 26 |
| 50-64 | 821 | 7.2 | 23 | 3 | 4 | 6 | 8 | 10 | 13 | 23 |
| 65+ | 1058 | 8.4 | 25 | 4 | 4 | 7 | 10 | 13 | 16 | 28 |
| TOT. SINGLE DX | 1699 | 6.4 | 7 | 3 | 4 | 6 | 8 | 9 | 11 | 16 |
| " MULTIPLE DX | 2405 | 7.6 | 23 | 3 | 4 | 7 | 9 | 11 | 14 | 26 |
| TOTAL 0-19 YRS | 7 | 3.0 | 2 | 1 | 1 | 3 | 4 | 5 | 5 | 5 |
| 20-34 | 125 | 6.1 | 6 | 3 | 3 | 6 | 7 | 9 | 11 | 15 |
| 35-49 | 846 | 6.3 | 11 | 3 | 4 | 6 | 7 | 9 | 10 | 21 |
| 50-64 | 1434 | 6.8 | 16 | 3 | 4 | 6 | 8 | 10 | 12 | 19 |
| 65+ | 1692 | 7.8 | 20 | 4 | 4 | 7 | 9 | 12 | 15 | 23 |
| GRAND TOTAL | 4104 | 7.1 | 17 | 3 | 4 | 6 | 8 | 10 | 13 | 21 |

### 745: Miscellaneous operations on breast (85.0; 85.24-85.25; 85.3; 85.5-85.8; 85.93-85.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 206 | 2.7 | 1 | 1 | 1 | 3 | 3 | 4 | 5 | 9 |
| 20-34 | 599 | 2.7 | 3 | <1 | <1 | 2 | 4 | 5 | 6 | 8 |
| 35-49 | 513 | 3.2 | 3 | <1 | <1 | 3 | 4 | 5 | 6 | 9 |
| 50-64 | 309 | 3.4 | 6 | 1 | 1 | 3 | 4 | 5 | 6 | 12 |
| 65+ | 83 | 3.3 | 3 | 1 | 1 | 3 | 4 | 6 | 7 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 60 | 3.7 | 6 | <1 | 2 | 3 | 4 | 5 | 8 | 16 |
| 20-34 | 359 | 3.5 | 6 | <1 | 1 | 3 | 5 | 6 | 7 | 15 |
| 35-49 | 665 | 4.0 | 12 | 1 | 1 | 4 | 5 | 6 | 8 | 13 |
| 50-64 | 486 | 4.3 | 6 | 1 | 2 | 4 | 5 | 7 | 8 | 12 |
| 65+ | 127 | 4.9 | 16 | 1 | 2 | 4 | 6 | 9 | 12 | 44 |
| **TOT. SINGLE DX** | 1710 | 3.0 | 4 | <1 | <1 | 3 | 4 | 5 | 6 | 9 |
| **" MULTIPLE DX** | 1697 | 4.1 | 10 | 1 | 1 | 4 | 5 | 7 | 8 | 14 |
| TOTAL 0-19 YRS | 266 | 3.0 | 3 | 1 | 1 | 3 | 3 | 4 | 6 | 9 |
| 20-34 | 958 | 3.0 | 5 | <1 | <1 | 3 | 4 | 5 | 6 | 9 |
| 35-49 | 1178 | 3.7 | 9 | <1 | 1 | 3 | 5 | 6 | 7 | 11 |
| 50-64 | 795 | 3.9 | 7 | 1 | 1 | 4 | 5 | 6 | 8 | 12 |
| 65+ | 210 | 4.3 | 12 | 1 | 2 | 4 | 5 | 7 | 10 | 25 |
| **GRAND TOTAL** | **3407** | **3.5** | **7** | **<1** | **1** | **3** | **4** | **6** | **7** | **12** |

### 746: Incision and excision of pilonidal cyst or sinus (86.03; 86.21)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 372 | 2.9 | 4 | 1 | 1 | 2 | 4 | 5 | 6 | 10 |
| 20-34 | 642 | 2.8 | 3 | 1 | 1 | 3 | 3 | 4 | 5 | 8 |
| 35-49 | 100 | 2.8 | 2 | <1 | 1 | 2 | 4 | 5 | 6 | 9 |
| 50-64 | 23 | 2.8 | 2 | <1 | 1 | 2 | 4 | 6 | 6 | 6 |
| 65+ | 2 | 4.0 | 2 | 3 | 3 | 4 | 5 | 5 | 5 | 5 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 45 | 4.0 | 8 | 1 | 1 | 4 | 5 | 7 | 8 | 19 |
| 20-34 | 123 | 3.7 | 8 | 1 | 2 | 3 | 4 | 6 | 9 | 19 |
| 35-49 | 32 | 5.2 | 50 | 1 | 1 | 4 | 6 | 8 | 11 | 42 |
| 50-64 | 17 | 6.1 | 27 | 1 | 2 | 4 | 7 | 15 | 18 | 20 |
| 65+ | 8 | 9.3 | 66 | 2 | 2 | 6 | 15 | 22 | 23 | 23 |
| **TOT. SINGLE DX** | 1139 | 2.8 | 4 | 1 | 1 | 2 | 3 | 5 | 6 | 8 |
| **" MULTIPLE DX** | 225 | 4.4 | 19 | 1 | 2 | 3 | 5 | 8 | 11 | 21 |
| TOTAL 0-19 YRS | 417 | 3.0 | 5 | 1 | 1 | 3 | 4 | 5 | 6 | 10 |
| 20-34 | 765 | 2.9 | 4 | 1 | 1 | 3 | 4 | 5 | 6 | 9 |
| 35-49 | 132 | 3.4 | 15 | 1 | 1 | 3 | 4 | 6 | 7 | 17 |
| 50-64 | 40 | 4.2 | 15 | <1 | 1 | 3 | 4 | 8 | 14 | 20 |
| 65+ | 10 | 8.2 | 57 | 2 | 2 | 4 | 12 | 21 | 23 | 23 |
| **GRAND TOTAL** | **1364** | **3.1** | **6** | **1** | **1** | **3** | **4** | **5** | **6** | **12** |

### 747: Local excision or destruction of lesion or tissue of skin and subcutaneous tissue (86.11; 86.3)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 617 | 1.8 | 4 | <1 | <1 | 1 | 2 | 3 | 5 | 11 |
| 20-34 | 646 | 2.6 | 13 | <1 | <1 | 2 | 3 | 5 | 8 | 20 |
| 35-49 | 612 | 2.5 | 9 | <1 | <1 | 2 | 3 | 5 | 7 | 15 |
| 50-64 | 534 | 2.7 | 12 | <1 | <1 | 2 | 3 | 5 | 9 | 15 |
| 65+ | 371 | 3.4 | 16 | <1 | <1 | 2 | 3 | 8 | 13 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 304 | 4.7 | 71 | <1 | <1 | 2 | 4 | 10 | 15 | 55 |
| 20-34 | 447 | 7.2 | 103 | <1 | 1 | 4 | 8 | 21 | 29 | 58 |
| 35-49 | 493 | 7.3 | 84 | <1 | 1 | 4 | 9 | 20 | 27 | 51 |
| 50-64 | 715 | 6.9 | 64 | <1 | 1 | 4 | 8 | 17 | 24 | 44 |
| 65+ | 948 | 10.1 | 135 | 1 | 2 | 6 | 12 | 23 | 36 | 59 |
| **TOT. SINGLE DX** | 2780 | 2.5 | 11 | <1 | <1 | 1 | 3 | 5 | 8 | 17 |
| **" MULTIPLE DX** | 2907 | 7.8 | 101 | <1 | 1 | 4 | 9 | 20 | 28 | 55 |
| TOTAL 0-19 YRS | 921 | 2.8 | 28 | <1 | <1 | 1 | 2 | 5 | 9 | 28 |
| 20-34 | 1093 | 4.5 | 55 | <1 | <1 | 2 | 4 | 10 | 19 | 49 |
| 35-49 | 1105 | 4.6 | 49 | <1 | <1 | 2 | 5 | 11 | 18 | 40 |
| 50-64 | 1249 | 5.1 | 47 | <1 | <1 | 3 | 6 | 12 | 18 | 40 |
| 65+ | 1319 | 8.2 | 111 | <1 | 1 | 4 | 10 | 20 | 29 | 57 |
| **GRAND TOTAL** | **5687** | **5.2** | **64** | **<1** | **<1** | **2** | **5** | **12** | **21** | **46** |

### 748: Other incision, excision, and suture of skin (86.04-86.09; 86.22; 86.4-86.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 553 | 3.9 | 20 | <1 | <1 | 2 | 5 | 9 | 13 | 25 |
| 20-34 | 784 | 4.6 | 25 | <1 | 1 | 3 | 6 | 10 | 14 | 27 |
| 35-49 | 472 | 5.3 | 51 | <1 | 1 | 3 | 6 | 12 | 17 | 32 |
| 50-64 | 393 | 6.1 | 61 | <1 | 1 | 4 | 7 | 14 | 20 | 44 |
| 65+ | 378 | 6.7 | 59 | 1 | 1 | 4 | 9 | 15 | 20 | 38 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 1627 | 5.6 | 82 | <1 | 1 | 3 | 6 | 13 | 23 | 53 |
| 20-34 | 2555 | 6.5 | 81 | 1 | 1 | 3 | 7 | 16 | 24 | 52 |
| 35-49 | 1403 | 8.9 | 127 | 1 | 1 | 5 | 11 | 21 | 31 | 69 |
| 50-64 | 1334 | 10.4 | 159 | 1 | 1 | 6 | 13 | 26 | 37 | 72 |
| 65+ | 1792 | 14.6 | 242 | 1 | 2 | 9 | 20 | 37 | 51 | 91 |
| **TOT. SINGLE DX** | 2530 | 5.1 | 41 | <1 | 1 | 3 | 6 | 12 | 16 | 31 |
| **" MULTIPLE DX** | 8711 | 9.0 | 145 | 1 | 1 | 5 | 11 | 23 | 34 | 69 |
| TOTAL 0-19 YRS | 2180 | 5.2 | 67 | <1 | 1 | 2 | 5 | 12 | 20 | 44 |
| 20-34 | 3339 | 6.0 | 69 | <1 | 1 | 3 | 7 | 14 | 22 | 46 |
| 35-49 | 1875 | 8.0 | 111 | 1 | 1 | 4 | 10 | 19 | 29 | 62 |
| 50-64 | 1727 | 9.4 | 141 | 1 | 1 | 5 | 12 | 24 | 32 | 68 |
| 65+ | 2170 | 13.2 | 219 | 1 | 2 | 8 | 17 | 34 | 49 | 82 |
| **GRAND TOTAL** | **11291** | **8.1** | **124** | **1** | **1** | **4** | **10** | **20** | **31** | **65** |

## 749: Free skin grafts (86.6; 86.91)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 132 | 5.2 | 32 | <1 | 1 | 3 | 7 | 12 | 15 | 29 |
| 20-34 | 147 | 6.1 | 36 | <1 | 1 | 4 | 8 | 12 | 18 | 37 |
| 35-49 | 76 | 4.9 | 25 | <1 | <1 | 3 | 7 | 13 | 17 | 22 |
| 50-64 | 76 | 7.2 | 77 | 1 | 1 | 3 | 10 | 18 | 21 | 48 |
| 65+ | 95 | 12.0 | 95 | 1 | 2 | 10 | 17 | 24 | 31 | 47 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 200 | 9.3 | 126 | 1 | 1 | 5 | 12 | 25 | 33 | 55 |
| 20-34 | 285 | 11.3 | 168 | 1 | 1 | 8 | 15 | 25 | 30 | 83 |
| 35-49 | 185 | 15.1 | 255 | 1 | 2 | 10 | 21 | 35 | 48 | 85 |
| 50-64 | 195 | 16.2 | 225 | 1 | 2 | 12 | 23 | 38 | 45 | 67 |
| 65+ | 327 | 21.6 | 321 | 2 | 4 | 18 | 29 | 49 | 62 | 88 |
| **TOT. SINGLE DX** | 526 | 6.9 | 56 | <1 | 1 | 4 | 9 | 16 | 21 | 37 |
| **" MULTIPLE DX** | 1192 | 15.2 | 246 | 1 | 2 | 10 | 21 | 35 | 48 | 83 |
| TOTAL 0-19 YRS | 332 | 7.7 | 93 | 1 | 1 | 4 | 9 | 20 | 28 | 44 |
| 20-34 | 432 | 9.5 | 129 | 1 | 1 | 6 | 12 | 20 | 28 | 56 |
| 35-49 | 261 | 12.2 | 210 | 1 | 1 | 7 | 16 | 30 | 40 | 78 |
| 50-64 | 271 | 13.7 | 200 | 1 | 1 | 9 | 20 | 34 | 43 | 59 |
| 65+ | 422 | 19.4 | 287 | 2 | 3 | 15 | 26 | 43 | 56 | 85 |
| **GRAND TOTAL** | **1718** | **12.7** | **203** | **1** | **1** | **8** | **18** | **30** | **42** | **73** |

## 750: Pedicle graft of skin (86.7)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 57 | 4.5 | 51 | <1 | 1 | 2 | 4 | 8 | 20 | 44 |
| 20-34 | 83 | 4.3 | 40 | <1 | <1 | 2 | 3 | 11 | 18 | 31 |
| 35-49 | 44 | 8.2 | 169 | <1 | 1 | 2 | 8 | 26 | 34 | 66 |
| 50-64 | 35 | 3.6 | 15 | <1 | 1 | 2 | 4 | 7 | 15 | 18 |
| 65+ | 20 | 7.8 | 75 | <1 | 1 | 4 | 16 | 26 | 60 | 99+ |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 68 | 7.8 | 277 | <1 | 1 | 2 | 4 | 17 | 45 | 91 |
| 20-34 | 104 | 15.9 | 393 | 1 | 1 | 9 | 23 | 47 | 78 | 99+ |
| 35-49 | 92 | 9.5 | 263 | 1 | 1 | 3 | 7 | 25 | 43 | 84 |
| 50-64 | 71 | 10.3 | 212 | 1 | 1 | 5 | 13 | 25 | 33 | 83 |
| 65+ | 60 | 21.0 | 544 | 1 | 1 | 10 | 36 | 61 | 82 | 98 |
| **TOT. SINGLE DX** | 239 | 5.3 | 68 | <1 | <1 | 2 | 4 | 15 | 25 | 48 |
| **" MULTIPLE DX** | 395 | 12.8 | 351 | 1 | 1 | 4 | 16 | 38 | 56 | 92 |
| TOTAL 0-19 YRS | 125 | 6.3 | 176 | <1 | 1 | 2 | 4 | 15 | 28 | 76 |
| 20-34 | 187 | 10.8 | 269 | <1 | 1 | 3 | 14 | 29 | 55 | 93 |
| 35-49 | 136 | 9.0 | 232 | <1 | 1 | 3 | 8 | 25 | 39 | 83 |
| 50-64 | 106 | 8.1 | 157 | 1 | 1 | 4 | 10 | 18 | 30 | 75 |
| 65+ | 80 | 17.7 | 458 | 1 | 1 | 9 | 27 | 56 | 82 | 99+ |
| **GRAND TOTAL** | **634** | **9.9** | **257** | **<1** | **1** | **3** | **11** | **27** | **45** | **85** |

## 751: Miscellaneous operations on skin and subcutaneous tissue (86.02; 86.23-86.25; 86.8; 86.99)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 251 | 1.8 | 1 | <1 | <1 | 1 | 2 | 3 | 4 | 9 |
| 20-34 | 167 | 2.8 | 6 | <1 | <1 | 2 | 3 | 5 | 8 | 14 |
| 35-49 | 134 | 2.2 | 2 | <1 | <1 | 2 | 3 | 5 | 6 | 8 |
| 50-64 | 164 | 2.2 | 3 | <1 | <1 | 2 | 3 | 4 | 5 | 10 |
| 65+ | 67 | 2.3 | 3 | <1 | <1 | 2 | 3 | 4 | 6 | 11 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 156 | 3.3 | 17 | <1 | <1 | 2 | 4 | 7 | 10 | 25 |
| 20-34 | 215 | 4.8 | 29 | <1 | 1 | 3 | 6 | 10 | 14 | 28 |
| 35-49 | 189 | 4.3 | 38 | <1 | 1 | 3 | 5 | 7 | 9 | 36 |
| 50-64 | 215 | 4.5 | 35 | <1 | 1 | 2 | 5 | 10 | 14 | 38 |
| 65+ | 156 | 6.9 | 57 | 1 | 1 | 4 | 11 | 15 | 22 | 40 |
| **TOT. SINGLE DX** | 783 | 2.2 | 4 | <1 | <1 | 2 | 3 | 4 | 6 | 11 |
| **" MULTIPLE DX** | 931 | 4.7 | 37 | <1 | 1 | 3 | 5 | 11 | 14 | 33 |
| TOTAL 0-19 YRS | 407 | 2.4 | 8 | <1 | <1 | 2 | 2 | 4 | 7 | 21 |
| 20-34 | 382 | 3.9 | 20 | <1 | <1 | 2 | 5 | 8 | 12 | 27 |
| 35-49 | 323 | 3.4 | 25 | <1 | <1 | 2 | 4 | 6 | 8 | 15 |
| 50-64 | 379 | 3.5 | 23 | <1 | <1 | 2 | 4 | 7 | 12 | 26 |
| 65+ | 223 | 5.5 | 46 | <1 | 1 | 3 | 7 | 14 | 18 | 34 |
| **GRAND TOTAL** | **1714** | **3.6** | **23** | **<1** | **<1** | **2** | **4** | **7** | **12** | **26** |

## 752: Miscellaneous diagnostic and therapeutic procedures on breast, skin, and subcutaneous tissue (85.19; 85.91-85.92; 86.01; 86.19; 86.26; 86.92)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 753: Encephalography and ventriculography (87.01-87.02)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 10 | 4.0 | 10 | 1 | 1 | 3 | 6 | 9 | 10 | 10 |
| 20-34 | 17 | 2.2 | 3 | 1 | 1 | 2 | 3 | 3 | 7 | 9 |
| 35-49 | 12 | 1.9 | 2 | <1 | <1 | 1 | 2 | 5 | 6 | 6 |
| 50-64 | 18 | 2.3 | 1 | 1 | 1 | 2 | 2 | 4 | 5 | 5 |
| 65+ | 20 | 6.7 | 36 | 1 | 1 | 4 | 10 | 17 | 19 | 19 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 19 | 9.3 | 215 | 1 | 1 | 3 | 11 | 30 | 48 | 57 |
| 20-34 | 17 | 6.4 | 44 | 1 | 2 | 3 | 8 | 19 | 21 | 22 |
| 35-49 | 22 | 9.8 | 205 | 1 | 2 | 5 | 8 | 30 | 51 | 53 |
| 50-64 | 45 | 6.9 | 53 | 1 | 2 | 5 | 8 | 14 | 20 | 40 |
| 65+ | 83 | 9.1 | 119 | 2 | 2 | 6 | 12 | 15 | 23 | 72 |
| **TOT. SINGLE DX** | 77 | 3.6 | 15 | 1 | 1 | 2 | 4 | 9 | 12 | 19 |
| **" MULTIPLE DX** | 186 | 8.4 | 115 | 1 | 2 | 5 | 11 | 16 | 24 | 56 |
| TOTAL 0-19 YRS | 29 | 7.5 | 148 | 1 | 1 | 3 | 9 | 16 | 39 | 57 |
| 20-34 | 34 | 4.3 | 28 | 1 | 1 | 3 | 4 | 11 | 19 | 22 |
| 35-49 | 34 | 7.0 | 146 | 1 | 1 | 2 | 7 | 15 | 44 | 53 |
| 50-64 | 63 | 5.6 | 42 | 1 | 1 | 3 | 7 | 12 | 16 | 38 |
| 65+ | 103 | 8.6 | 104 | 2 | 2 | 6 | 12 | 16 | 22 | 61 |
| **GRAND TOTAL** | **263** | **7.0** | **91** | **1** | **1** | **4** | **8** | **15** | **21** | **53** |

## 754: Myelography (87.21)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 138 | 3.0 | 11 | 1 | 1 | 2 | 3 | 8 | 10 | 17 |
| 20-34 | 2104 | 3.4 | 13 | 1 | 1 | 2 | 4 | 8 | 11 | 17 |
| 35-49 | 2686 | 3.4 | 15 | 1 | 1 | 2 | 4 | 8 | 11 | 19 |
| 50-64 | 1575 | 3.3 | 13 | 1 | 1 | 2 | 4 | 8 | 10 | 17 |
| 65+ | 499 | 4.1 | 20 | 1 | 1 | 2 | 5 | 10 | 14 | 23 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 114 | 4.6 | 22 | 1 | 1 | 3 | 6 | 11 | 14 | 22 |
| 20-34 | 1317 | 5.6 | 51 | 1 | 1 | 3 | 8 | 13 | 17 | 26 |
| 35-49 | 1912 | 5.7 | 44 | 1 | 1 | 3 | 8 | 12 | 17 | 30 |
| 50-64 | 1683 | 5.6 | 48 | 1 | 1 | 3 | 8 | 13 | 17 | 34 |
| 65+ | 1090 | 8.0 | 93 | 1 | 1 | 5 | 10 | 18 | 26 | 42 |
| **TOT. SINGLE DX** | 7002 | 3.4 | 15 | 1 | 1 | 2 | 4 | 8 | 11 | 18 |
| **" MULTIPLE DX** | 6116 | 6.0 | 56 | 1 | 1 | 3 | 8 | 14 | 18 | 35 |
| TOTAL 0-19 YRS | 252 | 3.7 | 17 | 1 | 1 | 2 | 4 | 10 | 14 | 20 |
| 20-34 | 3421 | 4.3 | 29 | 1 | 1 | 2 | 6 | 10 | 13 | 22 |
| 35-49 | 4598 | 4.3 | 29 | 1 | 1 | 2 | 5 | 10 | 14 | 24 |
| 50-64 | 3258 | 4.5 | 33 | 1 | 1 | 2 | 5 | 10 | 14 | 26 |
| 65+ | 1589 | 6.8 | 74 | 1 | 1 | 3 | 9 | 16 | 22 | 39 |
| **GRAND TOTAL** | **13118** | **4.6** | **36** | **1** | **1** | **2** | **6** | **11** | **15** | **27** |

## 755: Cerebral arteriography (88.41)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 46 | 2.7 | 3 | <1 | 1 | 2 | 3 | 6 | 7 | 9 |
| 20-34 | 186 | 4.3 | 70 | 1 | 1 | 2 | 4 | 8 | 14 | 46 |
| 35-49 | 260 | 3.7 | 15 | 1 | 1 | 2 | 4 | 8 | 13 | 21 |
| 50-64 | 510 | 3.0 | 9 | 1 | 1 | 2 | 3 | 5 | 9 | 16 |
| 65+ | 607 | 3.3 | 18 | 1 | 1 | 2 | 3 | 7 | 10 | 21 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 52 | 7.5 | 106 | 1 | 1 | 3 | 9 | 17 | 36 | 49 |
| 20-34 | 188 | 7.7 | 108 | 1 | 1 | 4 | 9 | 17 | 29 | 58 |
| 35-49 | 347 | 6.2 | 61 | 1 | 1 | 3 | 7 | 12 | 21 | 39 |
| 50-64 | 1116 | 5.9 | 70 | 1 | 1 | 3 | 7 | 13 | 19 | 49 |
| 65+ | 1719 | 6.7 | 57 | 1 | 2 | 4 | 8 | 14 | 20 | 40 |
| **TOT. SINGLE DX** | 1609 | 3.4 | 21 | 1 | 1 | 2 | 3 | 7 | 10 | 20 |
| **" MULTIPLE DX** | 3422 | 6.5 | 66 | 1 | 1 | 4 | 8 | 14 | 20 | 42 |
| TOTAL 0-19 YRS | 98 | 5.2 | 63 | 1 | 1 | 3 | 5 | 10 | 17 | 48 |
| 20-34 | 374 | 6.0 | 92 | 1 | 1 | 3 | 6 | 14 | 19 | 52 |
| 35-49 | 607 | 5.1 | 43 | 1 | 1 | 3 | 6 | 11 | 15 | 35 |
| 50-64 | 1626 | 5.0 | 53 | 1 | 1 | 3 | 6 | 10 | 16 | 37 |
| 65+ | 2326 | 5.8 | 50 | 1 | 1 | 3 | 7 | 13 | 18 | 38 |
| **GRAND TOTAL** | **5031** | **5.5** | **54** | **1** | **1** | **3** | **6** | **12** | **17** | **38** |

## 756: Coronary arteriography (88.55-88.57)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | 6 | 3.2 | 15 | 1 | 1 | 2 | 2 | 10 | 11 | 11 |
| 35-49 | 35 | 2.5 | 4 | <1 | 1 | 2 | 2 | 5 | 6 | 11 |
| 50-64 | 66 | 2.8 | 6 | 1 | 1 | 2 | 2 | 6 | 9 | 12 |
| 65+ | 33 | 3.5 | 12 | <1 | 1 | 2 | 3 | 10 | 12 | 15 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 20-34 | 10 | 3.4 | 12 | 1 | 1 | 2 | 4 | 9 | 13 | 13 |
| 35-49 | 126 | 4.6 | 20 | 1 | 1 | 2 | 6 | 10 | 14 | 21 |
| 50-64 | 333 | 4.1 | 13 | 1 | 1 | 2 | 5 | 10 | 11 | 17 |
| 65+ | 234 | 4.8 | 28 | <1 | 1 | 3 | 6 | 10 | 14 | 30 |
| **TOT. SINGLE DX** | 140 | 2.9 | 8 | 1 | 1 | 2 | 2 | 6 | 10 | 13 |
| **" MULTIPLE DX** | 706 | 4.4 | 20 | 1 | 1 | 2 | 6 | 10 | 12 | 21 |
| TOTAL 0-19 YRS | 3 | 2.0 | | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 20-34 | 16 | 3.3 | 13 | 1 | 1 | 2 | 3 | 10 | 12 | 13 |
| 35-49 | 161 | 4.1 | 17 | 1 | 1 | 2 | 5 | 9 | 13 | 18 |
| 50-64 | 399 | 3.9 | 12 | 1 | 1 | 2 | 5 | 9 | 11 | 17 |
| 65+ | 267 | 4.6 | 27 | <1 | 1 | 2 | 6 | 10 | 13 | 29 |
| **GRAND TOTAL** | **846** | **4.1** | **18** | **1** | **1** | **2** | **5** | **10** | **12** | **18** |

## 757: Other arteriography and phlebography (88.40; 88.42-88.49; 88.50-88.54; 88.58; 88.6)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 150 | 3.3 | 15 | 1 | 1 | 2 | 3 | 9 | 12 | 19 |
| 20-34 | 392 | 4.4 | 16 | 1 | 1 | 2 | 7 | 10 | 12 | 20 |
| 35-49 | 471 | 4.0 | 15 | 1 | 1 | 2 | 5 | 10 | 11 | 19 |
| 50-64 | 768 | 3.9 | 20 | 1 | 1 | 2 | 5 | 9 | 12 | 21 |
| 65+ | 770 | 3.7 | 20 | 1 | 1 | 2 | 4 | 9 | 12 | 19 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 199 | 4.2 | 56 | 1 | 1 | 2 | 4 | 10 | 13 | 66 |
| 20-34 | 501 | 6.4 | 68 | 1 | 1 | 4 | 8 | 13 | 16 | 31 |
| 35-49 | 821 | 6.8 | 57 | 1 | 1 | 4 | 9 | 14 | 18 | 45 |
| 50-64 | 2053 | 6.3 | 57 | 1 | 1 | 4 | 8 | 13 | 18 | 35 |
| 65+ | 3023 | 7.4 | 63 | 1 | 1 | 5 | 10 | 16 | 21 | 40 |
| **TOT. SINGLE DX** | 2551 | 3.9 | 19 | 1 | 1 | 2 | 5 | 10 | 12 | 19 |
| **" MULTIPLE DX** | 6597 | 6.8 | 61 | 1 | 1 | 4 | 9 | 14 | 19 | 38 |
| TOTAL 0-19 YRS | 349 | 3.9 | 39 | 1 | 1 | 2 | 3 | 10 | 13 | 34 |
| 20-34 | 893 | 5.5 | 47 | 1 | 1 | 3 | 7 | 11 | 15 | 26 |
| 35-49 | 1292 | 5.8 | 44 | 1 | 1 | 3 | 8 | 12 | 16 | 34 |
| 50-64 | 2821 | 5.6 | 49 | 1 | 1 | 3 | 8 | 12 | 16 | 31 |
| 65+ | 3793 | 6.0 | 57 | 1 | 1 | 4 | 9 | 15 | 20 | 37 |
| **GRAND TOTAL** | **9148** | **6.0** | **51** | **1** | **1** | **3** | **8** | **13** | **18** | **34** |

## 758: Other radiography and special diagnostic procedures (87.03-87.17; 87.22-87.99; 88.0-88.3; 88.7-88.8; 92.0-92.1)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 4737 | 3.3 | 17 | 1 | 1 | 2 | 4 | 6 | 8 | 19 |
| 20-34 | 7982 | 3.7 | 15 | 1 | 1 | 3 | 4 | 7 | 10 | 20 |
| 35-49 | 5852 | 4.2 | 19 | 1 | 1 | 3 | 5 | 8 | 11 | 21 |
| 50-64 | 5264 | 4.7 | 21 | 1 | 1 | 3 | 6 | 9 | 13 | 22 |
| 65+ | 5504 | 5.9 | 33 | 1 | 2 | 4 | 7 | 12 | 16 | 30 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 7574 | 4.7 | 47 | 1 | 1 | 3 | 5 | 9 | 14 | 40 |
| 20-34 | 12438 | 5.5 | 42 | 1 | 1 | 4 | 6 | 11 | 17 | 32 |
| 35-49 | 12641 | 6.3 | 44 | 1 | 2 | 4 | 7 | 12 | 19 | 32 |
| 50-64 | 21540 | 7.1 | 48 | 1 | 2 | 5 | 9 | 14 | 19 | 35 |
| 65+ | 47978 | 8.6 | 61 | 2 | 2 | 6 | 11 | 17 | 23 | 42 |
| **TOT. SINGLE DX** | 29339 | 4.3 | 22 | 1 | 1 | 3 | 5 | 9 | 12 | 23 |
| **" MULTIPLE DX** | 102171 | 7.3 | 55 | 1 | 2 | 5 | 9 | 15 | 21 | 38 |
| TOTAL 0-19 YRS | 12311 | 4.1 | 36 | 1 | 1 | 3 | 4 | 8 | 12 | 33 |
| 20-34 | 20420 | 4.8 | 33 | 1 | 1 | 3 | 5 | 9 | 14 | 29 |
| 35-49 | 18493 | 5.6 | 37 | 1 | 1 | 4 | 7 | 11 | 16 | 29 |
| 50-64 | 26804 | 6.6 | 44 | 1 | 2 | 5 | 8 | 13 | 18 | 33 |
| 65+ | 53482 | 8.3 | 60 | 2 | 2 | 6 | 10 | 16 | 22 | 41 |
| **GRAND TOTAL** | **131510** | **6.7** | **49** | **1** | **2** | **5** | **8** | **14** | **19** | **36** |

## 759: Therapeutic radiology and nuclear therapy (92.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **TOT. SINGLE DX** | | | | | | | | | | |
| **" MULTIPLE DX** | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| **GRAND TOTAL** | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 760: Special therapies and rehabilitation (93.0-93.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| **1. SINGLE DX** | | | | | | | | | | |
| 0-19 YRS | 2937 | 4.1 | 26 | 1 | 1 | 3 | 4 | 7 | 11 | 27 |
| 20-34 | 1577 | 6.1 | 33 | 1 | 2 | 4 | 7 | 12 | 18 | 29 |
| 35-49 | 1661 | 6.1 | 24 | 1 | 2 | 5 | 8 | 11 | 15 | 26 |
| 50-64 | 1490 | 6.0 | 23 | 1 | 2 | 5 | 8 | 11 | 14 | 24 |
| 65+ | 1538 | 7.6 | 52 | 2 | 2 | 6 | 9 | 14 | 19 | 34 |
| **2. MULTIPLE DX** | | | | | | | | | | |
| 0-19 YRS | 3530 | 6.0 | 66 | 1 | 1 | 4 | 6 | 13 | 21 | 44 |
| 20-34 | 2849 | 7.7 | 79 | 1 | 2 | 5 | 9 | 16 | 25 | 53 |
| 35-49 | 3212 | 8.1 | 61 | 2 | 2 | 6 | 10 | 17 | 22 | 39 |
| 50-64 | 5322 | 8.6 | 76 | 2 | 2 | 6 | 10 | 17 | 25 | 49 |
| 65+ | 12815 | 9.7 | 68 | 2 | 3 | 8 | 12 | 18 | 24 | 45 |
| **TOT. SINGLE DX** | 9203 | 5.7 | 33 | 1 | 1 | 4 | 7 | 11 | 15 | 28 |
| **" MULTIPLE DX** | 27728 | 8.6 | 72 | 2 | 2 | 6 | 11 | 17 | 24 | 46 |
| TOTAL 0-19 YRS | 6467 | 5.1 | 49 | 1 | 1 | 3 | 5 | 10 | 17 | 38 |
| 20-34 | 4426 | 7.1 | 64 | 1 | 2 | 5 | 8 | 15 | 22 | 46 |
| 35-49 | 4873 | 7.4 | 50 | 1 | 2 | 6 | 9 | 15 | 20 | 34 |
| 50-64 | 6812 | 8.0 | 66 | 1 | 2 | 6 | 9 | 15 | 22 | 44 |
| 65+ | 14353 | 9.5 | 68 | 2 | 3 | 7 | 12 | 18 | 24 | 45 |
| **GRAND TOTAL** | **36931** | **7.9** | **64** | **1** | **2** | **6** | **10** | **16** | **22** | **43** |

# Appendix

## *ICD-9-CM* Codes for Selected Operation Groups

## *ICD-9-CM* Codes for Selected Operation Groups

*ICD-9-CM* codes for operation groups 552, 557, 576, 619, 686, 713, and 741

| | | |
|---|---|---|
| 552 | Operations on tongue, oral cavity, and facial region except palatoplasty and operations on salivary glands | 25.01-25.02; 25.1-25.5; 25.92-25.99; 27.0-27.24; 27.3-27.5; 27.7; 27.92-27.99 |
| 557 | Other diagnostic and therapeutic procedures on nose, mouth, and pharynx | 21.00-21.01; 21.21; 21.29; 22.19; 24.19; 25.09; 25.91; 26.19; 27.29; 27.91; 28.19; 29.11; 29.19 |
| 576 | Other diagnostic and therapeutic procedures on larynx, trachea, bronchus, lungs, and heart | 31.41-31.42; 31.48-31.49; 33.21; 33.28-33.29; 34.28-34.29; 34.91-34.92; 37.29 |
| 619 | Miscellaneous diagnostic and therapeutic procedures on digestive tract, abdominal cavity, and abdominal wall | 42.22; 42.29; 44.12; 44.19; 45.12; 45.19; 45.22; 45.24; 45.28-45.29; 48.22-48.23; 48.29; 49.21; 49.29; 49.41-49.42; 50.19; 51.19; 52.19; 54.29 |
| 686 | Miscellaneous diagnostic and therapeutic procedures on female genital system | 65.19; 66.19; 67.19; 68.11; 68.19; 69.6; 69.7; 69.92; 70.21; 70.29; 71.19 |
| 713 | Other reduction of fracture without internal fixation, except maxillofacial and femur | 79.00-79.04; 79.06-79.09; 79.20-79.24; 79.26-79.29; 79.40-79.44; 79.46-79.49 |
| 741 | Other diagnostic and therapeutic procedures on musculoskeletal structure | 76.19; 76.96; 81.91-81.92; 81.98; 82.92-82.96; 83.29; 83.94-83.98; 84.41-84.43; 84.45-84.47 |

# Index to the Length of Stay Tables

If an operation does not appear in the Index, its corresponding *ICD-9-CM* code can be found in the *International Classification of Diseases, 9th Revision, Clinical Modification,* Volume 3, Procedures: Tabular List and Alphabetic Index, Commission on Professional and Hospital Activities, Ann Arbor, Michigan, 1978. The operation group containing the *ICD-9-CM* code can then be located in the length of stay tables.

# OPERATION INDEX

This operation index lists common operations and gives the number of the *ICD-9-CM List B* operation group to which each operation belongs.

| Group Number | Title |
|---|---|
| 555 | Adenoidectomy, except with tonsillectomy |
| 554 | With tonsillectomy |
| 515 | Adrenalectomy |
| 549 | Alveolectomy |
| 549 | Alveoloplasty |
| 695 | Amniocentesis (diagnostic) |
| 694 | For termination of pregnancy |
| | Amputation |
| 668 | Cervix |
| 732 | Finger |
| 737 | Foot |
| 733 | Forearm |
| 733 | Hand |
| 737 | Hip |
| 736 | Leg, above knee |
| 735 | Leg, below knee |
| 739 | Revision (any site) |
| 732 | Thumb |
| 734 | Toe |
| 733 | Upper arm |
| | Anastomosis |
| 617 | Bile duct |
| 581 | Blood vessel |
| 617 | Gallbladder |
| 601 | Intestine |
| 609 | Rectum |
| 757 | Angiocardiography, except coronary vessels |
| 756 | Coronary vessels |
| 757 | Angiography, except cardiac and cerebral vessels |
| 756 | Cardiac vessels |
| 755 | Cerebral vessels |
| 546 | Antrotomy, except radical maxillary |
| 545 | Radical maxillary |
| 572 | Aortocoronary bypass |
| 757 | Aortography |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| 603 | Appendectomy, except incidental |
| 604 | Incidental |
| | Arteriography |
| 757 | Aorta |
| 755 | Cerebral |
| 756 | Coronary |
| 757 | Femoral |
| 755 | Head |
| 756 | Heart |
| 757 | Renal |
| 741 | Arthrocentesis |
| 721 | Arthrodesis, except spinal |
| 720 | Spinal |
| 725 | Arthroplasty, except hip, knee, and temporomandibular |
| 723 | Hip, except total replacement |
| 722 | Total replacement |
| 724 | Knee |
| 699 | Temporomandibular |
| 719 | Arthroscopy |
| 716 | Arthrotomy |
| | Artificial |
| 686 | Insemination |
| 690 | Rupture of membranes for delivery |
| | Aspiration |
| 752 | Breast, except for biopsy |
| 742 | Biopsy |
| 741 | Bursa |
| 682 | Cul-de-sac |
| 677 | Curettage, uterus, except for termination of pregnancy |
| 674 | Termination of pregnancy |
| 651 | Hydrocele, except of spermatic cord |
| 656 | Spermatic cord |
| 632 | Kidney |
| 614 | Liver, percutaneous, except for biopsy |
| 615 | Biopsy |
| 537 | Middle ear |
| 546 | Nasal sinus |
| 663 | Ovary |
| 752 | Skin |
| 656 | Spermatocele |
| 529 | Vitreous |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Augmentation |
| 745 | Breast |
| 699 | Chin |
| | Biopsy |
| 515 | Adrenal gland |
| 611 | Anus |
| 638 | Bladder, except transurethral approach |
| 637 | Transurethral approach |
| 705 | Bone, except facial |
| 698 | Facial |
| 586 | Bone marrow |
| 501 | Brain |
| 742 | Breast |
| 563 | Bronchus |
| 501 | Cerebral meninges |
| 668 | Cervix |
| 597 | Colon |
| 597 | Duodenum |
| 668 | Endocervical |
| 588 | Esophagus |
| 532 | Eyeball |
| 517 | Eyelid |
| 666 | Fallopian tubes |
| 597 | Intestine (large) (small) |
| 719 | Joint structure |
| 632 | Kidney |
| 597 | Large intestine |
| 558 | Larynx |
| 552 | Lip |
| 615 | Liver |
| 565 | Lung |
| 582 | Lymph nodes |
| 570 | Mediastinum |
| | Meninges |
| 501 | Cerebral |
| 507 | Spinal |
| 727 | Muscle |
| 543 | Nose |
| 532 | Orbit |
| 660 | Ovary, except by aspiration |
| 663 | Aspiration |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Biopsy (continued) |
| 618 | Pancreas |
| 658 | Penis |
| 508 | Peripheral nerve |
| 626 | Peritoneum |
| 556 | Pharynx |
| 570 | Pleura |
| 650 | Prostate |
| 606 | Rectum |
| 651 | Scrotum |
| 747 | Skin |
| 597 | Small intestine |
| 507 | Spinal meninges |
| 654 | Testes |
| 514 | Thyroid |
| 552 | Tongue |
| 559 | Trachea |
| 670 | Uterus |
| 682 | Vagina |
| 558 | Vocal cord |
| 683 | Vulva |
| 518 | Blepharorrhaphy |
| 763 | Blood transfusion, except intrauterine exchange |
| 696 | Intrauterine exchange |
| 685 | Bone marrow transplant |
| 745 | Breast augmentation |
| 688 | Breech extraction for delivery |
| 567 | Bronchoscopy |
| 704 | Bunionectomy |
| 730 | Bursectomy, except of hand |
| 728 | Hand |
| 726 | Bursotomy |
| | Bypass, graft or shunt |
| 572 | Aortocoronary |
| 578 | Blood vessels (systemic) |
| 572 | Coronary |
| 596 | Gastric |
| 572 | Heart |
| 579 | Heart-lung |
| 601 | Intestine |
| 596 | Stomach |
| 634 | Urinary |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| 763 | Cancer chemotherapy |
| | Cardiac |
| 573 | Catheterization |
| 764 | Function tests |
| | Massage |
| 763 | Closed |
| 575 | Open |
| 574 | Pacemaker insertion or replacement |
| 575 | Revascularization, except coronary bypass |
| 572 | Coronary bypass |
| 764 | Stress tests |
| 509 | Carpal tunnel decompression or release |
| 760 | Cast application |
| 758 | C.A.T. scan |
| 526 | Cataract extraction, except intracapsular |
| 525 | Intracapsular |
| | Catheterization |
| 647 | Bladder, except suprapubic approach |
| 638 | Suprapubic approach |
| 581 | Blood vessel |
| 573 | Cardiac |
| 645 | Ureteral |
| 647 | Urethral, indwelling |
| 587 | Vein, except umbilical |
| 581 | Umbilical |
| 669 | Cerclage, cervix |
| 692 | Cesarean section, except low cervical |
| 691 | Low cervical |
| 763 | Chemotherapy |
| 758 | Cholangiogram (intravenous) (percutaneous) |
| 616 | Cholecystectomy |
| 758 | Cholecystogram |
| 617 | Cholecystotomy |
| 764 | Circulatory monitoring |
| 657 | Circumcision, except newborn |
| | Closure |
| 617 | Cholecystostomy |
| 601 | Colostomy |
| | Fistula – *see Fistulectomy* |
| 596 | Gastric anastomosis |
| 601 | Sigmoidostomy |
| 570 | Thoracostomy |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Closure (continued) |
| 635 | Ureterostomy |
| 640 | Urethrostomy |
| 599 | Colectomy |
| 568 | Collapse of lung, surgical |
| 619 | Colonoscopy, except surgical approach |
| 609 | Surgical approach (transabdominal) |
| 602 | Colostomy, except loop |
| 600 | Loop |
| 681 | Colporrhaphy, except for cystocele or rectocele repair |
| 680 | Cystocele or rectocele repair |
| 758 | Computerized axial tomography (C.A.T. scan) |
| 667 | Conization of cervix |
| 520 | Conjunctivoplasty |
| 764 | Consultation |
| | Control of hemorrhage |
| 638 | Bladder |
| 581 | Blood vessel |
| 569 | Chest |
| 627 | Laparotomy site |
| 544 | Nose, except by anterior packing |
| 557 | Anterior packing |
| 569 | Pleura |
| 581 | Postoperative, not elsewhere classified |
| 650 | Prostate |
| 590 | Stomach |
| 556 | Tonsil |
| 521 | Corneal transplant |
| 572 | Coronary bypass |
| 503 | Cranioplasty |
| 502 | Craniotomy |
| 668 | Cryoconization, cervix |
| 682 | Culdocentesis |
| 682 | Culdoscopy, except for tubal ligation |
| 664 | Tubal ligation |
| 523 | Cyclocryotherapy (eye) |
| 523 | Cyclodialysis |
| 638 | Cystectomy (bladder) |
| 680 | Cystocele repair |
| 636 | Cystoscopy |
| 638 | Cystostomy |
| 638 | Cystotomy |

## 761: Psychiatric diagnostic and therapeutic procedures (94.0-94.5)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| TOT. SINGLE DX | | | | | | | | | | |
| " MULTIPLE DX | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| GRAND TOTAL | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 762: Non-specific removal of foreign body (98.0-98.2)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| TOT. SINGLE DX | | | | | | | | | | |
| " MULTIPLE DX | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| GRAND TOTAL | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 763: Other non-operative therapeutic procedures (99.0-99.9)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| TOT. SINGLE DX | | | | | | | | | | |
| " MULTIPLE DX | | | | | | | | | | |
| TOTAL 0-19 YRS | | | | | | | | | | |
| 20-34 | | | | | | | | | | |
| 35-49 | | | | | | | | | | |
| 50-64 | | | | | | | | | | |
| 65+ | | | | | | | | | | |
| GRAND TOTAL | | | | | | | | | | |

This group consists of procedures that are coded for indexing, quality review, and purposes other than their effect on length of stay. For the purpose of operation length of stay tables, patients with only these procedures are not considered "operated" patients.

## 764: Miscellaneous diagnostic and therapeutic procedures (89.0-91.9; 95.0-97.8)

| TYPE OF PATIENT (1) | TOTAL PATIENTS (2) | AVG. STAY (3) | VARIANCE (4) | PERCENTILES 5th (5) | 10th (6) | 50th (7) | 75th (8) | 90th (9) | 95th (10) | 99th (11) |
|---|---|---|---|---|---|---|---|---|---|---|
| 1. SINGLE DX | | | | | | | | | | |
| 0-19 YRS | 5038 | 2.8 | 11 | 1 | 1 | 2 | 3 | 5 | 6 | 13 |
| 20-34 | 2500 | 3.7 | 16 | <1 | 1 | 3 | 4 | 7 | 10 | 21 |
| 35-49 | 1665 | 4.3 | 21 | 1 | 1 | 3 | 5 | 9 | 12 | 24 |
| 50-64 | 1904 | 5.0 | 19 | 1 | 1 | 4 | 7 | 10 | 13 | 21 |
| 65+ | 2141 | 5.9 | 30 | 1 | 2 | 4 | 7 | 11 | 15 | 28 |
| 2. MULTIPLE DX | | | | | | | | | | |
| 0-19 YRS | 6782 | 4.7 | 41 | 1 | 1 | 3 | 5 | 9 | 14 | 36 |
| 20-34 | 4549 | 5.2 | 39 | 1 | 1 | 3 | 6 | 11 | 16 | 33 |
| 35-49 | 4855 | 6.0 | 37 | 1 | 1 | 4 | 7 | 12 | 17 | 30 |
| 50-64 | 9863 | 6.9 | 45 | 1 | 2 | 5 | 9 | 14 | 18 | 35 |
| 65+ | 21640 | 8.3 | 60 | 2 | 2 | 6 | 10 | 16 | 22 | 41 |
| TOT. SINGLE DX | 13248 | 4.0 | 19 | 1 | 1 | 3 | 5 | 8 | 11 | 21 |
| " MULTIPLE DX | 47689 | 7.0 | 52 | 1 | 2 | 5 | 9 | 14 | 20 | 37 |
| TOTAL 0-19 YRS | 11820 | 3.9 | 30 | 1 | 1 | 3 | 4 | 7 | 10 | 29 |
| 20-34 | 7049 | 4.7 | 32 | 1 | 1 | 3 | 5 | 9 | 14 | 30 |
| 35-49 | 6520 | 5.5 | 34 | 1 | 1 | 4 | 7 | 11 | 16 | 29 |
| 50-64 | 11767 | 6.6 | 41 | 1 | 2 | 5 | 8 | 13 | 18 | 32 |
| 65+ | 23781 | 8.1 | 58 | 2 | 2 | 6 | 10 | 16 | 21 | 40 |
| GRAND TOTAL | 60937 | 6.3 | 46 | 1 | 1 | 4 | 8 | 13 | 18 | 35 |

| Group Number | Title |
|---|---|
| 675 | D and C (uterus), except for termination of pregnancy |
| 674 | Termination of pregnancy |
| 519 | Dacryocystorhinostomy |
| | Debridement |
| 748 | Burn (skin) |
| 705 | Fracture, except of skull |
| 503 | Skull |
| 748 | Skin (infection) (wound) |
| | Decompression |
| 509 | Carpal tunnel |
| 502 | Cranial |
| 506 | Laminectomy |
| 506 | Spinal |
| | Delivery |
| 690 | Artificial rupture of membranes |
| 688 | Breech extraction |
| 692 | Cesarean section, except low cervical |
| 691 | Low cervical |
| 689 | Forceps, except low forceps |
| 687 | Low forceps |
| 689 | Internal and combined version |
| 751 | Dermabrasion |
| 601 | Derotation, intestine |
| 580 | Dialysis, except peritoneal |
| 629 | Peritoneal |
| | Dilation |
| 638 | Bladder (neck) |
| 635 | Ureteral meatus |
| 645 | Ureterovesical orifice |
| 641 | Urethra, urethrovesical junction |
| 675 | Dilation and curettage (D and C), uterus, except for termination of pregnancy |
| 674 | Termination of pregnancy |
| | Dislocation-reduction, except temporomandibular |
| 714 | Closed |
| 715 | Open |
| 702 | Temporomandibular (joint) |
| | Division |
| | Adhesions – *see Lysis, adhesions* |
| 613 | Anal sphincter |
| 726 | Fascia, exept of hand |
| 728 | Hand |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Division (continued) |
| 508 | Peripheral nerve |
| 511 | Sympathetic nerve or ganglion |
| 659 | Dorsal or lateral slit of prepuce |
| | Drainage — *see also Incision (and drainage)* |
| 647 | Bladder, except surgical approach |
| 638 | Surgical approach (suprapubic) |
| 569 | Chest |
| 537 | Middle ear |
| 763 | Drug therapy |
| 598 | Duodenectomy |
| 764 | ECG (electrocardiogram) |
| 758 | Echoencephalography |
| 764 | EEG (electroencephalogram) |
| 764 | EKG (electrocardiogram) |
| 764 | Electrocardiogram (ECG) (EKG) |
| 764 | Electroencephalogram (EEG) |
| 752 | Electrolysis, skin |
| 753 | Encephalography |
| | Endoscopy |
| 636 | Bladder |
| 567 | Bronchus |
| 619 | Colon, except surgical approach |
| 609 | Surgical approach (transabdominal) |
| 682 | Cul-de-sac, except for tubal ligation |
| 664 | Tubal ligation |
| 589 | Esophagus |
| 589 | Gastric, except surgical approach |
| 596 | Surgical approach (transabdominal) |
| 576 | Larynx |
| 628 | Peritoneum |
| 557 | Pharynx |
| 609 | Small intestine |
| 589 | Stomach, except surgical approach |
| 596 | Surgical approach (transabdominal) |
| 570 | Thorax, transpleural |
| 636 | Urethra |
| 679 | Uterus |
| 686 | Vagina |
| 602 | Enterostomy |
| 532 | Enucleation, eyeball |

| Group Number | Title |
|---|---|
| 533 | Epilation, eyelid |
| 697 | Episioperinorrhaphy |
| 697 | Episiorrhaphy |
| 690 | Episiotomy, except with forceps delivery |
| 687 | With low forceps delivery |
| 689 | With other forceps delivery |
| 544 | Epistaxis control, except by anterior packing |
| 557 | Anterior packing |
| 588 | Esophagectomy |
| 589 | Esophagoscopy, except surgical approach |
| 619 | Surgical approach |
| 588 | Esophagotomy |
| 613 | Evacuation, hemorrhoids |
| | Evisceration |
| 532 | Eyeball |
| 673 | Pelvic |
| | Examination |
| 764 | Eye |
| 764 | General physical |
| 764 | Gynecologic |
| 764 | Pelvic |
| 761 | Psychiatric |
| 763 | Exchange transfusion, except intrauterine |
| 696 | Intrauterine |
| | Excision — *see also Excision of lesion (and destruction of tissue)* |
| 610 | Anal fistula |
| 727 | Baker's cyst |
| 705 | Bone, except facial |
| 699 | Facial, except partial |
| 698 | Partial |
| 556 | Branchial cleft cyst |
| 704 | Bunion |
| 517 | Chalazion, eyelid |
| 599 | Colon |
| 665 | Fallopian tubes, bilateral, except endoscopic approach |
| 664 | Endoscopic approach |
| 670 | Fibroid, uterine |
| 616 | Gallbladder |
| 508 | Ganglion, except of tendon |
| 727 | Tendon (of hand) |
| 612 | Hemorrhoids |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Excision (continued) |
| | Hydrocele |
| 679 | Round ligament |
| 656 | Spermatic cord |
| 651 | Tunica vaginalis |
| 717 | Intervertebral disc or cartilage |
| 719 | Joint, except intervertebral disc and cartilage of knee |
| 717 | Intervertebral disc |
| 718 | Knee cartilage |
| 631 | Kidney, except partial |
| 632 | Partial |
| 718 | Knee cartilage (semilunar) |
| 582 | Lymph nodes, except radical |
| 583 | Radical |
| 508 | Neuroma, peripheral nerve |
| | Ovary |
| 662 | Bilateral |
| 661 | Unilateral |
| 550 | Parotid gland |
| 746 | Pilonidal cyst or sinus |
| 515 | Pituitary gland |
| 649 | Prostate, except transurethral approach |
| 648 | Transurethral approach |
| 522 | Pterygium |
| 747 | Sebaceous cyst, except of eyelid |
| 517 | Eyelid, eyebrow |
| 718 | Semilunar cartilage, knee |
| 586 | Spleen |
| 514 | Thyroglossal duct or tract |
| 513 | Thyroid |
| 751 | Toenail |
| 552 | Tongue |
| 553 | Tonsil, except with adenoids |
| 554 | With adenoids |
| 556 | Tonsil tag |
| | Tubes and ovaries |
| 662 | Bilateral |
| 661 | Unilateral |
| 670 | Uterine fibroid |
| 577 | Varicocele, except of spermatic cord |
| 656 | Spermatic cord |

| Group Number | Title |
|---|---|
| | Excision of lesion (and destruction of tissue) |
| 626 | Abdominal wall |
| 611 | Anus |
| 617 | Bile duct |
| 638 | Bladder, except transurethral approach |
| 637 | Transurethral approach |
| 705 | Bone, except facial |
| 698 | Facial |
| 501 | Brain |
| 742 | Breast |
| 563 | Bronchus |
| 501 | Cerebral meninges |
| 668 | Cervix, except by conization |
| 667 | Conization |
| 597 | Colon |
| 520 | Conjunctiva |
| 522 | Cornea |
| 597 | Duodenum |
| 534 | Ear, external |
| 588 | Esophagus |
| 532 | Eye |
| 517 | Eyelid |
| 666 | Fallopian tube |
| 549 | Gingiva, gums |
| 597 | Intestine |
| 524 | Iris |
| 632 | Kidney |
| 597 | Large intestine |
| 558 | Larynx |
| 552 | Lip |
| 565 | Lung |
| | Meninges |
| 501 | Cerebral |
| 507 | Spinal |
| 727 | Muscle |
| | Nerve |
| 508 | Peripheral |
| 511 | Sympathetic |
| 543 | Nose, except submucous resection |
| 541 | Submucous resection |

| Group Number | Title |
|---|---|
| | Excision of lesion (continued) |
| 660 | Ovary |
| 550 | Parotid gland |
| 658 | Penis |
| | Perineum |
| 683 | Female |
| 747 | Male |
| 508 | Peripheral nerve |
| 607 | Perirectal tissue |
| 626 | Peritoneum |
| 606 | Rectum |
| 550 | Salivary gland |
| 524 | Sclera |
| 651 | Scrotum |
| 747 | Skin, except radical and of pilonidal cyst or sinus |
| 746 | Pilonidal cyst or sinus |
| 748 | Radical |
| 501 | Skull |
| 597 | Small intestine |
| 507 | Spinal meninges |
| 592 | Stomach |
| 727 | Tendon |
| 654 | Testis |
| 552 | Tongue |
| 559 | Trachea |
| 635 | Ureter |
| 640 | Urethra |
| 679 | Uterine ligament |
| 670 | Uterus |
| 682 | Vagina |
| 559 | Vocal cord |
| 683 | Vulva |
| | Exenteration |
| 532 | Eyeball, orbit |
| 673 | Pelvic |
| | Exploration |
| 627 | Abdominal cavity |
| 617 | Bile duct |
| 632 | Kidney |
| 627 | Exploratory laparotomy |
| 600 | Exteriorization, intestine |
| 579 | Extracorporeal circulation |

| Group Number | Title |
|---|---|
| | Extraction |
| 688 | Breech, for delivery |
| 547 | Dental |
| 686 | Menstrual |
| 547 | Teeth |
| 689 | Vacuum, for delivery |
| 751 | Face lift |
| 730 | Fasciectomy, except of hand |
| 728 | Hand |
| 726 | Fasciotomy, except of hand |
| 728 | Hand |
| | Fistulectomy |
| 610 | Anal |
| 638 | Bladder |
| 556 | Branchial cleft |
| 564 | Bronchus |
| 601 | Intestine |
| 542 | Nasal |
| 542 | Nasolabial |
| 546 | Oroantral |
| 609 | Perirectal |
| 556 | Pharynx |
| 640 | Rectourethral |
| 679 | Rectouterine |
| 681 | Rectovaginal |
| 638 | Rectovesical |
| 609 | Rectum |
| 562 | Tracheoesophageal |
| 523 | Fistulization, sclera (for glaucoma) |
| 741 | Fitting, limb prosthesis |
| 706 | Fixation, internal, of bone, except with fracture-reduction |
| | With fracture-reduction — *see Fracture-reduction* |
| 689 | Forceps delivery, except low forceps |
| 687 | Low forceps |
| | Formation |
| 581 | Arteriovenous shunt for dialysis |
| 634 | Ileal conduit or bladder |
| | Fracture-reduction, except facial bone, femur, and nose |
| 713 | Closed, except with internal fixation |
| 711 | With internal fixation |
| 713 | Open, except with internal fixation |
| 712 | With internal fixation |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Fracture-reduction, except facial bone, femur, and nose (continued) |
| 701 | Facial bone, except closed reduction |
| 700 | Closed reduction |
| | Femur |
| 710 | Closed, except with internal fixation |
| 708 | With internal fixation |
| 710 | Open, except with internal fixation |
| 709 | With internal fixation |
| 701 | Mandible, except closed reduction |
| 700 | Closed reduction |
| 701 | Maxillofacial, except closed reduction |
| 700 | Closed reduction |
| 544 | Nose |
| 552 | Frenectomy (labial) (tongue) |
| 557 | Frenotomy (labial) (tongue) |
| 721 | Fusion, joint, except spinal |
| 720 | Spinal |
| 508 | Ganglionectomy, except of tendon |
| 727 | Tendon (of hand) |
| 508 | Gasserian ganglionectomy |
| 593 | Gastrectomy, except total |
| 594 | Total |
| 589 | Gastroscopy, except surgical approach |
| 596 | Surgical approach (transabdominal) |
| 591 | Gastrostomy |
| 590 | Gastrotomy |
| 699 | Genioplasty |
| 758 | G.I. series x-ray |
| 549 | Gingivectomy, gingivoplasty |
| 523 | Glaucoma operation |
| 552 | Glossectomy |
| | Graft |
| 706 | Bone, except facial and skull |
| 702 | Facial |
| 503 | Skull |
| 749 | Free, of skin |
| 552 | Lip |
| 731 | Muscle, except of hand |
| 729 | Hand |
| 750 | Pedicle, of skin |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Graft (continued) |
| | Skin |
| 749 | Free |
| 750 | Pedicle |
| 731 | Tendon, except of hand |
| 729 | Hand |
| 573 | Heart catheterization |
| 579 | Heart-lung bypass |
| 599 | Hemicolectomy |
| 580 | Hemodialysis |
| 612 | Hemorrhoidectomy |
| 615 | Hepatectomy |
| | Herniorrhaphy |
| 625 | Diaphragmatic |
| 624 | Epigastric |
| 622 | Femoral |
| 624 | Incisional |
| | Inguinal |
| 621 | Bilateral |
| 620 | Unilateral |
| 623 | Umbilical |
| 624 | Ventral |
| 723 | Hip replacement, except total |
| 722 | Total |
| 576 | His bundle study |
| 515 | Hypophysectomy |
| 671 | Hysterectomy, except radical and vaginal |
| 673 | Radical |
| 672 | Vaginal |
| 678 | Hysteropexy |
| 679 | Hysteroscopy |
| 670 | Hysterotomy, except for termination of pregnancy |
| 693 | Termination of pregnancy |
| 602 | Ileostomy, except loop |
| 600 | Loop |
| 634 | Ileoureterostomy |
| | Implant, implantation |
| 745 | Breast |
| 740 | Limb prosthesis |

| Group Number | Title |
|---|---|
| | Incision (and drainage) |
| 626 | Abdominal wall |
| 610 | Anal fistula |
| 605 | Appendix |
| 683 | Bartholin's gland or abscess |
| 703 | Bone, except facial |
| 702 | Facial |
| 501 | Brain |
| 745 | Breast |
| 501 | Cerebral meninges |
| 569 | Chest (wall) |
| 522 | Cornea |
| 502 | Cranial sinus |
| 534 | Ear, external |
| 518 | Eyelid |
| 552 | Floor of mouth |
| 617 | Gallbladder |
| 549 | Gingiva, gums |
| 600 | Intestine |
| 716 | Joint |
| 519 | Lacrimal gland or sac |
| 614 | Liver |
| 582 | Lymph nodes |
| 539 | Mastoid |
| | Meninges |
| 501 | Cerebral |
| 506 | Spinal |
| 537 | Middle ear |
| 726 | Muscle |
| | Nasal sinus — *see Sinusotomy* |
| 726 | Palmar space (hand) |
| 618 | Pancreas |
| 613 | Perianal tissue |
| | Perineum |
| 683 | Female |
| 748 | Male |
| 607 | Perirectal tissue |
| 627 | Peritoneum, except via cul-de-sac |
| 682 | Via cul-de-sac |
| 746 | Pilonidal cyst or sinus |
| 569 | Pleura |
| 607 | Rectovaginal septum |
| 607 | Rectum |

| Group Number | Title |
|---|---|
| | Incision (and drainage) (continued) |
| 626 | Retroperitoneal abscess |
| 651 | Scrotum |
| | Sinus (nasal) — *see Sinusotomy* |
| 748 | Skin, except pilonidal cyst or sinus |
| 746 | Pilonidal cyst or sinus |
| 506 | Spinal meninges |
| 726 | Tendon |
| 569 | Thorax |
| 633 | Ureter |
| 682 | Vagina |
| 683 | Vulva |
| 690 | Induction of labor (medical or surgical) for delivery |
| | Injection |
| 694 | Intra-amniotic, for termination of pregnancy |
| 763 | Therapeutic agent (drugs) |
| 745 | Breast |
| 741 | Bursa |
| 741 | Joint |
| 741 | Tendon |
| 686 | Insemination, artificial |
| | Insertion |
| 574 | Cardiac pacemaker |
| 686 | Intrauterine contraceptive device |
| 679 | Laminaria, uterus |
| 666 | Insufflation, fallopian tubes |
| 694 | Intra-amniotic injection for termination of pregnancy |
| 758 | Intravenous pyelogram (IVP) |
| 764 | Intubation, nonoperative |
| 764 | Irrigation, nonoperative |
| 758 | IVP (intravenous pyelogram) |
| | Laceration repair (suture) — *see Repair, by site* |
| 506 | Laminectomy, except for excision of intervertebral disc |
| 717 | Excision of intervertebral disc |
| 628 | Laparoscopy, except for tubal ligation |
| 664 | Tubal ligation |
| 627 | Laparotomy, exploratory |
| 560 | Laryngectomy, except partial |
| 559 | Partial |
| 576 | Laryngoscopy |

| Group Number | Title |
|---|---|
| 562 | Laryngotomy, except permanent |
| 561 | Permanent |
| 706 | Lengthening of bone |
| 526 | Lens extraction, except intracapsular |
| 525 | Intracapsular |
| | Ligation |
| 665 | Fallopian tubes (bilateral), except endoscopic approach |
| 664 | Endoscopic approach |
| 612 | Hemorrhoids |
| 577 | Varicose veins |
| 655 | Vas deferens |
| 577 | Ligation and stripping, varicose veins |
| | Lobectomy |
| 501 | Brain |
| 615 | Liver |
| 566 | Lung, except partial |
| 565 | Partial |
| 687 | Low forceps delivery |
| 512 | Lumbar puncture |
| | Lysis, adhesions |
| 630 | Abdominal |
| 741 | Hand |
| 524 | Iris |
| 716 | Joint |
| 731 | Muscle, except of hand |
| 741 | Hand |
| 630 | Pelvis |
| 630 | Peritoneum |
| 731 | Tendon, except of hand |
| 741 | Hand |
| 745 | Mammary augmentation |
| 745 | Mammectomy |
| 758 | Mammography |
| 745 | Mammoplasty, reduction |
| 699 | Mandibulectomy, except partial |
| 698 | Partial |
| 699 | Mandibulotomy |
| | Mastectomy |
| 743 | Complete |
| 743 | Simple, except extended |
| 744 | Extended |

| Group Number | Title |
|---|---|
| | Mastectomy (continued) |
| 742 | Subtotal |
| 744 | Radical |
| 743 | Partial |
| 538 | Mastoidectomy, except revision |
| 539 | Revision |
| 745 | Mastotomy |
| | Meatotomy |
| 635 | Ureteral |
| 639 | Urethral |
| 690 | Medical induction of labor |
| 718 | Meniscectomy, knee |
| 686 | Menstrual extraction or regulation |
| 699 | Mentoplasty |
| 764 | Monitoring, circulatory (CVP) (PCW) |
| 754 | Myelography |
| 670 | Myomectomy, uterus |
| 537 | Myringotomy |
| 631 | Nephrectomy (complete) |
| 632 | Nephropexy |
| 682 | Obliteration, vagina or vaginal vault |
| 751 | Onychectomy |
| 751 | Onychoplasty |
| | Oophorectomy |
| 662 | Bilateral |
| 661 | Unilateral |
| 533 | Ophthalmoscopy |
| 652 | Orchiectomy |
| 653 | Orchiopexy |
| 654 | Orchioplasty, except suspension |
| 653 | Suspension |
| 705 | Ostectomy, except facial bone |
| 699 | Facial bone, except partial |
| 698 | Partial |
| 706 | Osteoplasty, except facial bone |
| 699 | Facial bone |
| 703 | Osteotomy, except facial bone |
| 699 | Facial bone |
| 540 | Otoscopy |

| Group Number | Title |
|---|---|
| 551 | Palatoplasty |
| 618 | Pancreatectomy |
| 618 | Pancreatotomy |
| | Paracentesis |
| 626 | Abdominal |
| 524 | Anterior chamber (eye) |
| 576 | Thoracic |
| 514 | Parathyroidectomy |
| 550 | Parotidectomy |
| 673 | Pelvic exenteration |
| 575 | Pericardiocentesis |
| 629 | Peritoneal dialysis |
| 628 | Peritoneoscopy |
| 557 | Pharyngoscopy |
| 556 | Pharyngotomy |
| 757 | Phlebography |
| 540 | Piercing, ear |
| 576 | Pleurocentesis |
| | Plication |
| 644 | Urethral, urethrovesical |
| 581 | Vena cava |
| 753 | Pneumoencephalography |
| 566 | Pneumonectomy |
| 659 | Preputiotomy |
| 608 | Proctectomy |
| 609 | Proctopexy |
| 619 | Proctosigmoidoscopy, except surgical approach |
| 609 | Surgical approach (transabdominal) |
| 607 | Proctotomy |
| 649 | Prostatectomy, except transurethral approach |
| 648 | Transurethral approach |
| 761 | Psychiatric testing and evaluation |
| 761 | Psychotherapy |
| | Puncture |
| 501 | Cranial |
| 512 | Lumbar |
| 501 | Skull, for drainage |
| 512 | Spinal |
| 758 | Pyelogram (intravenous) (retrograde) |
| 636 | Pyeloscopy |
| 632 | Pyelotomy |
| 590 | Pyloromyotomy |
| 596 | Pyloroplasty |

| Group Number | Title |
|---|---|
| 759 | Radiation therapy |
| | Radiography |
| 757 | Arteries, except cerebral and coronary |
| 755 | Cerebral |
| 756 | Coronary |
| 758 | Breast |
| 755 | Cerebral |
| 755 | Arteries |
| 753 | Ventricles |
| 758 | Chest |
| 758 | Gallbladder |
| 758 | Kidney |
| 758 | Skeletal |
| 758 | Spine, except myelogram |
| 754 | Myelogram |
| 757 | Veins |
| | Reattachment |
| 738 | Extremity |
| 528 | Retina |
| 731 | Tendon, except of hand |
| 729 | Hand |
| 530 | Recession, ocular muscle |
| | Reconstruction |
| 535 | Ear ossicles |
| 699 | Mandible |
| 729 | Thumb |
| 680 | Rectocele repair |
| | Reduction |
| | Dislocation, except temporomandibular |
| 714 | Closed |
| 715 | Open |
| 702 | Temporomandibular (joint) |
| | Fracture, except facial bone, femur, and nose |
| 713 | Closed, except with internal fixation |
| 711 | With internal fixation |
| 713 | Open, except with internal fixation |
| 712 | With internal fixation |
| 701 | Facial bone, except closed reduction |
| 700 | Closed reduction |

| Group Number | Title |
|---|---|
| | Reduction (continued) |
| | Femur |
| 710 | Closed, except with internal fixation |
| 708 | With internal fixation |
| 710 | Open, except with internal fixation |
| 709 | With internal fixation |
| 701 | Mandible, except closed reduction |
| 700 | Closed reduction |
| 701 | Maxillofacial, except closed reduction |
| 700 | Closed reduction |
| 544 | Nose |
| 619 | Hemorrhoids |
| | Hernia – *see also Herniorrhaphy* |
| 764 | Manual |
| 601 | Intestinal torsion |
| 601 | Intussusception, intestine |
| 745 | Mammoplasty |
| 601 | Volvulus, intestine |
| | Release |
| | Adhesions – *see Lysis, adhesions* |
| 728 | Dupuytren's contracture |
| 749 | Scar contracture, skin |
| 642 | Urethral stricture |
| | Removal |
| | Calculus |
| 617 | Biliary tract |
| 638 | Bladder |
| 632 | Kidney, except transurethral approach |
| 635 | Transurethral approach |
| 519 | Lacrimal tract (by incision) |
| 533 | Without incision |
| 633 | Ureter, except transurethral approach |
| 635 | Transurethral approach |
| 764 | Catheter, nonoperatively |
| 764 | Cholecystotomy tube |
| 693 | Ectopic fetus |
| 707 | Fixation device, internal |
| | Foreign body |
| 762 | Bronchus, except with incision |
| 563 | With incision |
| 532 | Eye, except as below |
| 524 | Anterior chamber |
| 520 | Conjunctiva |

| Group Number | Title |
|---|---|
| | Removal (continued) |
| 524 | Iris |
| 527 | Lens |
| | Prosthesis |
| 745 | Breast |
| 716 | Joint |
| 697 | Retained placenta, manually |
| 764 | Sutures |
| 764 | T-tube, gallbladder |
| 764 | Therapeutic device, nonoperatively |
| 529 | Vitreous |
| 580 | Renal dialysis |
| | Repair |
| 630 | Abdominal wall |
| 613 | Anal fistula, except by excision |
| 610 | Excision |
| 581 | Aneurysm (arteriovenous) |
| 638 | Bladder, except obstetric laceration |
| 697 | Obstetric laceration |
| 518 | Blepharoptosis |
| 581 | Blood vessel |
| 706 | Bone, except facial |
| 699 | Facial |
| 745 | Breast |
| 564 | Bronchial fistula |
| 704 | Bunion |
| 571 | Cardiac septal defect |
| 669 | Cervix, except obstetric laceration |
| 697 | Obstetric laceration |
| 570 | Chest wall |
| 551 | Cleft palate |
| 520 | Conjunctiva |
| 522 | Cornea, except transplant |
| 521 | Transplant |
| 680 | Cystocele, vagina |
| 730 | Diastasis recti |
| 534 | Ear, external |
| 518 | Ectropion or entropion, eyelid |
| 697 | Episiotomy |
| 518 | Eyelid |
| | Fistula — *see Fistulectomy* |
| 704 | Hallux valgus |
| 729 | Hand |

| Group Number | Title |
|---|---|
| | Repair (continued) |
| | Hernia — *see Herniorrhaphy* |
| 723 | Hip, except total replacement |
| 722 | Total replacement |
| | Hydrocephalus, by ventricular shunt |
| 504 | Extracranial |
| 505 | Intracranial (ventriculostomy) |
| 601 | Intestine |
| 725 | Joint, except hip and knee |
| 723 | Hip, except total replacement |
| 722 | Total replacement |
| 724 | Knee |
| 724 | Knee (joint) |
| | Laceration — *see also Repair, by site* |
| 697 | Obstetric |
| 562 | Larynx |
| 552 | Lip |
| 615 | Liver |
| 568 | Lung |
| 731 | Muscle, except extraocular and of hand |
| 531 | Extraocular |
| 729 | Hand |
| 556 | Nasopharyngeal atresia |
| 542 | Nose |
| 663 | Ovary |
| 551 | Palate |
| 658 | Penis |
| 685 | Perineum, female, except obstetric laceration |
| 697 | Obstetric laceration |
| 509 | Peripheral nerve |
| 596 | Pylorus, stomach |
| 680 | Rectocele, vagina |
| 609 | Rectum |
| 528 | Retinal detachment or tear |
| 730 | Rotator-cuff (shoulder) |
| 503 | Skull |
| 507 | Spina bifida |
| 507 | Spinal meningocele |
| 596 | Stomach |
| 644 | Stress incontinence, except urethral suspension |
| 643 | Urethral suspension |
| 731 | Tendon, except of hand |
| 729 | Hand |

| Group Number | Title |
|---|---|
| | Repair (continued) |
| 654 | Testis |
| 562 | Trachea |
| 635 | Ureter |
| 640 | Urethra |
| 679 | Uterus, except obstetric laceration and supporting structures |
| 697 | Obstetric laceration |
| 678 | Supporting structures |
| 681 | Vagina, except for cystocele and rectocele, and obstetric laceration |
| 680 | Cystocele and rectocele |
| 697 | Obstetric laceration |
| 552 | Vestibular arch (mouth) |
| 685 | Vulva, except obstetric laceration |
| 697 | Obstetric laceration |
| | Replacement |
| 574 | Cardiac pacemaker |
| 571 | Heart valve |
| 725 | Joint, except hip and knee |
| 723 | Hip, except total replacement |
| 722 | Total replacement |
| 724 | Knee |
| 647 | Urinary catheter |
| | Resection |
| 638 | Bladder |
| 581 | Blood vessel |
| 564 | Bronchus |
| 599 | Large intestine |
| 730 | Muscle, except of hand |
| 728 | Hand |
| 541 | Nasal septum, submucous |
| 530 | Ocular muscle |
| 660 | Ovary |
| 608 | Rectum (abdominoperineal) (pull-through) |
| 549 | Root of tooth |
| 598 | Small intestine |
| 730 | Tendon, except of hand |
| 728 | Hand |
| 548 | Restoration, tooth |
| 758 | Retrograde pyelogram |
| | Revision |
| 739 | Amputation stump |
| 581 | Arteriovenous shunt for dialysis |
| 596 | Gastric anastomosis |

| Group Number | Title |
|---|---|
| | Revision (continued) |
| 601 | Intestinal anastomosis |
| 539 | Mastoidectomy |
| 562 | Tracheostomy |
| 539 | Tympanoplasty |
| 505 | Ventricular shunt |
| 542 | Rhinoplasty |
| 751 | Rhytidectomy, facial |
| 549 | Root canal therapy |
| 549 | Root resection (tooth) |
| 690 | Rupture of membranes for delivery |
| | Salpingectomy |
| 665 | Bilateral |
| 666 | Unilateral |
| | Salpingo-oophorectomy |
| 662 | Bilateral |
| 661 | Unilateral |
| 730 | Scalenectomy |
| 726 | Scalenotomy |
| | Scan |
| 758 | C.A.T. (computerized axial tomography) |
| 758 | Radioisotope |
| 528 | Scleral buckling, for retinal detachment |
| 706 | Shortening, bone |
| | Shunt — *see also Bypass* |
| 504 | Cerebral ventricles, except intracranial and revision |
| 505 | Intracranial |
| 505 | Revision |
| 578 | Peripheral vessels |
| 630 | Peritoneovenous |
| 578 | Systemic vessels |
| 550 | Sialoadenectomy |
| 599 | Sigmoidectomy |
| 546 | Sinusectomy, nasal |
| 546 | Sinusotomy, except radical maxillary |
| 545 | Radical maxillary |
| 751 | Size-reduction, skin tissue except breast |
| 745 | Breast |
| | Sphincterotomy |
| 617 | Ampullary (of Vater) (pancreas) |
| 613 | Anus |

## OPERATION INDEX (continued)


| Group Number | Title |
|---|---|
| | Sphincterotomy (continued) |
| 638 | Bladder neck |
| | Spinal |
| 720 | Fusion |
| 512 | Tap or puncture |
| 586 | Splenectomy |
| 721 | Stabilization of joint, except foot and knee |
| 725 | Foot |
| 724 | Knee |
| 535 | Stapedectomy |
| 535 | Stapes mobilization |
| | Stress incontinence repair, except by |
| 644 | urethral suspension |
| 643 | Urethral suspension |
| | Stress test |
| 764 | Cardiac |
| 764 | Fetal |
| 764 | Uterine |
| 541 | Submucous nasal resection |
| 690 | Surgical induction of labor for delivery |
| | Suspension |
| 643 | Urethra, except with bladder neck |
| 644 | With bladder neck |
| 644 | Urethrovesical |
| | Suture – *see also Repair, by site* |
| 730 | Muscle, except of hand |
| 729 | Hand |
| 748 | Skin |
| 730 | Tendon, except of hand |
| 729 | Hand |
| 510 | Sympathectomy |
| 719 | Synovectomy |
| | |
| 554 | T and A |
| | Takedown – *see Closure* |
| 730 | Tenosynovectomy, except of hand |
| 728 | Hand |
| 726 | Tenotomy, except of hand |
| 728 | Hand |
| | Termination of pregnancy |
| 676 | Aspiration curettage |
| 674 | Dilation and curettage |

| Group Number | Title |
|---|---|
| | Termination of pregnancy (continued) |
| 691 | Hysterotomy |
| 694 | Intra-amniotic injection |
| | Therapy |
| 761 | Behavior |
| 763 | Cancer (chemical) |
| 761 | Electroconvulsive (ECT) |
| 760 | Occupational |
| 760 | Physical |
| 761 | Psychiatric |
| 759 | Radiation |
| 760 | Rehabilitation |
| 761 | Shock (EST) |
| 760 | Speech |
| 758 | Thermography |
| 576 | Thoracentesis |
| 570 | Thoracoscopy, transpleural |
| 569 | Thoracotomy |
| 515 | Thymectomy |
| 513 | Thyroidectomy |
| 552 | Tongue-tie operation |
| 553 | Tonsillectomy, except with adenoidectomy |
| 554 | With adenoidectomy |
| 548 | Tooth restoration |
| 562 | Tracheostomy, except permanent |
| 561 | Permanent |
| 760 | Traction, skeletal, except with fracture-reduction |
| | With fracture-reduction — *see Reduction, fracture* |
| | Transfer — *see Transplant* |
| | Transfusion |
| 763 | Blood |
| 763 | Exchange, except intrauterine |
| 696 | Intrauterine |
| 585 | Bone marrow |
| 763 | Coagulation factor |
| 763 | Gamma globulin |
| 763 | Plasma |
| | Transplant, transplantation |
| 585 | Bone marrow |
| 521 | Corneal |
| 571 | Heart |
| 632 | Kidney |

| Group Number | Title |
|---|---|
| | Transplant, transplantation (continued) |
| 731 | Muscle, except of hand |
| 729 | Hand |
| 731 | Tendon, except of hand |
| 729 | Hand |
| 648 | TUR of prostate |
| 543 | Turbinectomy, nasal |
| 536 | Tympanoplasty, except revision |
| 539 | Revision |
| 537 | Tympanotomy |
| 758 | Ultrasound, diagnostic |
| 635 | Ureterectomy |
| 634 | Ureterostomy |
| 633 | Ureterotomy |
| 636 | Urethroscopy |
| 642 | Urethrotomy |
| 634 | Urinary diversion |
| 764 | Uroflowmetry, uroflometry |
| 763 | Vaccination |
| 689 | Vacuum extraction for delivery |
| 686 | Vaginoscopy |
| 595 | Vagotomy |
| 577 | Varicose vein operation |
| 655 | Vasectomy |
| 656 | Vasoplasty |
| 655 | Vasotomy |
| 656 | Vasovasostomy |
| 757 | Venography |
| 504 | Ventricular shunt, except revision |
| 505 | Revision |
| 753 | Ventriculography |
| 684 | Vulvectomy |
| | X-ray — *see Radiography* |
| 751 | Z-plasty, skin |